




Denali National Park and Preserve

Geologic Resources Inventory Report

Natural Resource Report NPS/NRPC/GRD/NRR—2010/244





THIS PAGE:
Mountaineering rangers on backcountry patrol in a land of giants. Ruth Glacier, Denali National Park and Preserve.

National Park Service photograph by Tucker Chenoweth, Denali National Park and Preserve.

ON THE COVER:
View of the terminus of Buckskin Glacier with Mount McKinley (20,320 ft) in the background, Denali National Park and Preserve, Alaska, August 6, 2004.

Photo courtesy of Ronald D. Karpilo, Jr., Colorado State University.

Denali National Park and Preserve

Geologic Resources Inventory Report

Natural Resource Report NPS/NRPC/GRD/NRR—2010/244

Geologic Resources Division
Natural Resource Program Center
P.O. Box 25287
Denver, Colorado 80225

September 2010

U.S. Department of the Interior
National Park Service
Natural Resource Program Center
Ft. Collins, Colorado

The National Park Service, Natural Resource Program Center publishes a range of reports that address natural resource topics of interest and applicability to a broad audience in the National Park Service and others in natural resource management, including scientists, conservation and environmental constituencies, and the public.

The Natural Resource Report Series is used to disseminate high-priority, current natural resource management information with managerial application. The series targets a general, diverse audience, and may contain NPS policy considerations or address sensitive issues of management applicability.

All manuscripts in the series receive the appropriate level of peer review to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and designed and published in a professional manner. This report received informal peer review by subject-matter experts who were not directly involved in the collection, analysis, or reporting of the data.

Views, statements, findings, conclusions, recommendations, and data in this report do not necessarily reflect views and policies of the National Park Service, U.S. Department of the Interior. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the U. S. Government.

Printed copies of this report are produced in a limited quantity and they are only available as long as the supply lasts. This report is available from the Geologic Resources Inventory website (http://www.nature.nps.gov/geology/inventory/gre_publications.cfm) and the Natural Resource Publications Management website (<http://www.nature.nps.gov/publications/NRPM>).

Please cite this publication as:

Thornberry-Ehrlich, T. 2010. Denali National Park and Preserve: geologic resources inventory report. Natural Resource Report NPS/NRPC/GRD/NRR—2010/244. National Park Service, Ft. Collins, Colorado.

Contents

Lists of Figures and Tables	iv
Executive Summary	v
Acknowledgements.....	vi
Introduction	1
<i>Purpose of the Geologic Resources Inventory</i>	<i>1</i>
<i>Park Setting and Establishment</i>	<i>1</i>
<i>Geologic Setting</i>	<i>1</i>
Geologic Issues	5
<i>Introduction.....</i>	<i>5</i>
<i>Glacial Issues.....</i>	<i>5</i>
<i>Earthquakes and Earthquake Hazards.....</i>	<i>6</i>
<i>Mining and Disturbed Lands.....</i>	<i>8</i>
<i>Permafrost Issues</i>	<i>9</i>
<i>Surface Water Issues</i>	<i>10</i>
<i>Paleontological Resources</i>	<i>11</i>
<i>Geothermal Activity</i>	<i>12</i>
<i>Slope Processes and Erosion.....</i>	<i>12</i>
<i>General Geology Issues and Research Possibilities.....</i>	<i>13</i>
Geologic Features and Processes.....	20
<i>Mt. McKinley (Denali).....</i>	<i>20</i>
<i>Denali Fault System.....</i>	<i>20</i>
<i>Kantishna Hills.....</i>	<i>21</i>
<i>Accreted Terranes.....</i>	<i>22</i>
<i>Glacial Features.....</i>	<i>22</i>
<i>Glaciers</i>	<i>24</i>
<i>Glacial Retreat and Surging Glaciers.....</i>	<i>25</i>
<i>Other Notable Geologic Features.....</i>	<i>25</i>
Map Unit Properties	31
<i>Geologic Units of Denali National Park and Preserve.....</i>	<i>32</i>
Geologic History	51
<i>Precambrian (prior to 542 million years ago).....</i>	<i>51</i>
<i>Paleozoic Era (542 to 251 million years ago).....</i>	<i>51</i>
<i>Mesozoic Era (251 to 65.5 million years ago).....</i>	<i>51</i>
<i>Cenozoic Era (the past 65.5 million years).....</i>	<i>52</i>
Glossary.....	58
Literature Cited.....	64
Appendix A: Overview of Digital Geologic Data.....	69
Appendix B: Scoping Session Participants	73
Attachment 1: Geologic Resources Inventory Products CD	

List of Figures

Figure 1: Map of Alaska Showing major mountain ranges and rivers.	2
Figure 2: Generalized cross section through central Alaska showing the juxtaposition of accreted terranes	3
Figure 3: Map of Denali National Park and Preserve.....	4
Figure 4: The Straightaway Glacier is one of many surge-type glaciers which issue from Mt McKinley.....	14
Figure 5: Glacial retreat photography pairing at the head of Hidden Creek.....	15
Figure 6: Glacial retreat photography pairing in the Teklanika River valley.....	16
Figure 7: Abandoned mined land reseteration	17
Figure 8: Thermokarst development.....	17
Figure 9: Toklat River.....	18
Figure 10: Theropod discovery in the Cantwell Formation.....	18
Figure 11: Plant fossils provide insight to the paleoecology of Denali.....	18
Figure 12: View of landslide debris on Buckskin Glacier.....	19
Figure 13: Researchers investigate dinosaur fossils in Denali's Cantwell Formation.....	19
Figure 14: Denali (Mt. McKinley) dominates the ridgeline of the Alaska Range.	26
Figure 15: Denali has a complex geologic history which includes active plate dynamics.....	26
Figure 16: Map of Alaska showing major faults and other regional tectonic structures.....	27
Figure 17: Cross section of subduction zone between the Talkeetna Superterrane and the Yukon-Tanana Terrane	28
Figure 18: Diagrammatic view of a retreating glacier.....	29
Figure 19: Short-lived features occurring on the surface of glaciers.....	29
Figure 20: Ice and crevasses of Eldridge Glacier.....	30
Figure 21: Geologic timescale.	54
Figure 22: Three-dimensional model of an active margin setting	55
Figure 23: Reversal of stream direction from southward to northward accompanying uplift of the Alaska Range.	55
Figure 24: Map of Alaska during the Paleocene and Eocene epochs	56

List of Tables

Table 1: Summary diagram of geologic events in the Denali area	57
--	----

Executive Summary

This report accompanies the digital geologic map for Denali National Park and Preserve in Alaska, produced by the Geologic Resources Division in collaboration with its partners. It contains information relevant to resource management and scientific research. This document incorporates preexisting geologic information and does not include new data or additional fieldwork.

Denali National Park and Preserve covers more than 6 million acres and features a vast and varied geological landscape. The park protects large portions of the long, arcuate Alaska Range—including Mount McKinley (known as “Denali” [“The High One”] in the native Athabaskan language), which has the greatest vertical relief of any mountain on Earth. The active tectonic processes of southern Alaska are still uplifting the mountains along the Denali fault system. Countering this uplift are weathering and erosion of the highlands by wind and water. Like much of Alaska, the whole of Denali National Park and Preserve is composed of accreted terranes—where the Pacific plate, acting like a conveyor belt, has been bringing bits of islands, the ocean floor, and slivers of other continents northward to for hundreds of millions of years forming a “jigsaw puzzle” of these terranes. The compositions, structures, metamorphic grades, and fossils of these terranes set them apart from neighboring rocks separated by discrete faults. The terranes are covered with more recent sedimentary deposits, and are studded with igneous intrusive rocks. Glacial ice covers a significant amount of the surface area of Denali National Park and Preserve; glaciers carve through thousands of meters of sedimentary, metamorphic, and igneous rocks exposed within the park, creating wide U-shaped valleys and transporting vast amounts of sediment. These glaciers flow over 40 km (25 mi), descending more than 4,500 vertical meters (14,800 ft) from the highest peaks of the range to the lowland hills below. The size, climate, isolation, and complexity of the park’s geologic setting create both a challenge to scientists hoping to understand the environment and a valuable target for preservation of a pristine ecosystem.

Understanding the geology in south-central Alaska leads to an appreciation for the unique relationship between geology and the environment. Geologic processes give rise to rock formations, topographic expression, surface and subsurface fluid movement, and soils; thus, geologic units hold clues to the history of the area. Man-made disturbances at the park are also significant. River terraces and channel deposits were reworked during mining periods, locally altering the topographic expression of the landscape and threatening riparian zone health. Human developments may have altered the fragile permafrost regime; as well, glaciers are shrinking due to climate change.

Denali National Park and Preserve’s Resource Stewardship Strategy for 2008-2027 identified several geologic processes, values, and components fundamental to achieving the park’s goals and maintaining its significance. The following are critical geologic issues for park management:

- **Glacier Issues.** Today, glacial ice covers approximately 17% of the park. Glacier processes are incredibly dynamic and have far-reaching effects. Resource management concerns include interactions with the changing climate (manifested as glacial thinning or surging glaciers) to visitor safety on glacial surfaces and unstable glacial deposits. The aerial photographic record of glaciers at Denali is extensive, providing an important tool for evaluating glacial changes. Glaciers carved vast valleys and deposited thick mantles of unconsolidated sediment on park slopes, which are vulnerable to slope processes and susceptible to intense erosion. Continual monitoring of the “state of health” of the park’s glaciers would help resource managers understand both the past and future conditions of the glaciers.
- **Seismicity.** Active faults are prevalent on the park’s landscape. The large-scale Denali fault system runs through the entire park, extending more than 2,100 km (1,300 mi). Fault processes are active; many fresh fault scarps reveal recent movement, and more than 600 seismic events are measured in the area each year. More than 70% of these events are relatively small—magnitude 1.5 to 2.5—and lie beneath the Kantishna Hills, which is a growing anticline. However, even small-magnitude earthquakes can damage park infrastructure, including buildings and roads, and undermine slope stability throughout the park. On November 3, 2002, a magnitude 7.9 earthquake centered east of the park caused severe local shaking. This event was followed by increased seismic activity, which will require close to 14 years for conditions to return to background level. Seismic and geodetic monitoring provides important knowledge of the tectonic setting, geologic structure, and activity throughout the area. Studies of ancient earthquake history on important faults can yield data on recurrence intervals and past magnitudes.

- **Mining and Disturbed Lands.** Alaska has long been associated with vast mineral wealth. Gold was discovered in the park area in 1903. Most ore-bearing veins were intruded into brittle faults and fractures within metamorphosed pelitic rocks. Much mining activity focused on the placer gold deposits of the Kantishna Hills region in the northern foothills area of the park. Extensive terrace areas were processed during extraction efforts. Massive tailing piles remain, exposing heavy metal-laden material to weathering and causing acid mine drainage. Concerns include: water quality; long-term riparian zone health; channel and floodplain morphology; and visitor safety. Mining continues in the Kantishna Hills area with strict environmental protection regulations. Studies of the change in the chemistry of the waters near mined or disturbed areas should be considered.

Other geologic issues at the park include permafrost, surface water issues, paleontological resources, geothermal energy, and slope processes. Permafrost is

present across the northern portion of the park. Permafrost in Alaska has been warming since the 1970s. Melting permafrost can have dramatic effects on the landscape of the park, including the development of thermokarst. Surface water quality in the park is influenced by the underlying geology. The significant paleontological resources of the park are diverse and span hundreds of millions of years. A Paleontology Management Plan is currently being developed by the park. Geothermal resources are present within the park; however a comprehensive inventory or monitoring plan has not yet been completed. Slope processes (mass wasting) present a geologic hazard in many areas of the park. Seismic activity increases the risk for mass wasting within the park.

The glossary contains definitions of many of the technical terms used in the report. A geologic timescale is provided as figure 21.

Acknowledgements

The Geologic Resources Inventory (GRI) is one of 12 inventories funded by the National Park Service Inventory and Monitoring Program. The GRI is administered by the Geologic Resources Division of the Natural Resource Program Center.

The Geologic Resources Division relies heavily on partnerships with institutions such as the U.S. Geological Survey, Colorado State University, state geologic surveys, local museums, and universities in developing GRI products.

Special thanks to: Guy Adema (Denali National Park and Preserve) for providing information and images for the report. John Burghardt (NPS Geologic Resources Division) provided substantial comments on the mining and disturbed lands section. Kerry Moss and Julia Brunner of the NPS Geologic Resources Division also provided information.

Credits

Author

Trista Thornberry-Ehrlich (Colorado State University)

Review

Peter Haeussler (U.S. Geological Survey)
Will Elder (National Park Service)
Tim Connors (NPS Geologic Resources Division)

Editing

Bonnie Dash (Envirocal)

Digital Geologic Data Production

Jim Chappell (Colorado State University)

Digital Geologic Data Overview Layout Design

Phil Reiker (NPS Geologic Resources Division)



Dedication

This report is dedicated to the memory of Phil Brease, Denali park geologist from 1986 until his death in 2010. His passion for geology and geology education continues to be an inspiration to many. Photo courtesy Guy Adema (Denali National Park and Preserve).

Introduction

The following section briefly describes the National Park Service Geologic Resources Inventory and the regional geologic setting of Denali National Park and Preserve.

Purpose of the Geologic Resources Inventory

The Geologic Resources Inventory (GRI) is one of 12 inventories funded by the National Park Service (NPS) Inventory and Monitoring Program. The GRI, administered by the Geologic Resources Division of the Natural Resource Program Center, is designed to provide and enhance baseline information available to park managers. The GRI team relies heavily on partnerships with institutions such as the U.S. Geological Survey, Colorado State University, state geologic surveys, local museums, and universities in developing GRI products.

The goals of the GRI are to increase understanding of the geologic processes at work in parks and to provide sound geologic information for use in park decision making. Sound park stewardship requires an understanding of the natural resources and their role in the ecosystem. Park ecosystems are fundamentally shaped by geology. The compilation and use of natural resource information by park managers is called for in section 204 of the National Parks Omnibus Management Act of 1998 and in NPS-75, Natural Resources Inventory and Monitoring Guideline.

To realize these goals, the GRI team is systematically conducting a scoping meeting for each of the 270 identified natural area parks and providing a park-specific digital geologic map and geologic report. These products support the stewardship of park resources and are designed for nongeoscientists. Scoping meetings bring together park staff and geologic experts to review available geologic maps and discuss specific geologic issues, features, and processes.

The GRI mapping team converts the geologic maps identified for park use at the scoping meeting into digital geologic data in accordance with their Geographic Information Systems (GIS) Data Model. These digital data sets bring an interactive dimension to traditional paper maps. The digital data sets provide geologic data for use in park GIS and facilitate the incorporation of geologic considerations into a wide range of resource management applications. The newest maps contain interactive help files. This geologic report assists park managers in the use of the map and provides an overview of park geology and geologic resource management issues.

For additional information regarding the content of this report and current GRI contact information please refer to the Geologic Resources Inventory web site (<http://www.nature.nps.gov/geology/inventory/>).

Park Setting and Establishment

Mount McKinley National Park was established on February 26, 1917 by an act of Congress. The park was intended as "... a public park for the benefit and enjoyment of the people... for recreation purposes by the public and for the preservation of animals, birds, and fish and for the preservation of the natural curiosities and scenic beauties thereof ... said park shall be, and is hereby established as a game refuge." (39 Stat. 938). In 1976, the area was declared an International Biosphere Reserve. In 1980, the Alaska National Interest Lands Conservation Act added approximately 3.8 million acres to Mount McKinley National Park (about 1.9 million acres of which were to become the Denali Wilderness); at this point, the park was renamed as Denali National Park and Preserve. Today, the park covers more than 6 million acres, preserving and protecting large portions of the Alaska Range, including the 6,194-m- (20,320- ft)-tall Mount McKinley—known as "Denali" ("The High One")—in the native Athabaskan language). The park is situated approximately 200 km (125 mi) north of Anchorage, Alaska (fig. 1).

Geologic Setting

Mount McKinley's two peaks tower over neighboring summits, ultimately rising from 61 m (200 ft) at the lowest point in the park (on the Yentna River) to 6,194 m (20,320 ft). The peaks form an east-west-trending 1,000-km- (600-mi)-long line of mountains known as the Alaska Range. This major regional feature forms a topographic barrier and drainage divide between the coastal lowlands around Cook Inlet and the Yukon lowlands of the interior of Alaska (Brease 2004). The Denali fault runs parallel to the length of the central and eastern Alaska Range, and it crosses most of Alaska. Approximately 375 km of right-lateral offset have been documented along the fault since Cretaceous time (Lowey 1998), and the fault remains seismically active.

The Alaska Range has permanent snow cover on northern exposures above approximately 2,100 m (7,000 ft). This snowpack supports several large glaciers. Over 17% of the park's land (more than 1 million acres) is covered with glacial ice. The largest glacier is the 55-km- (34-mi)-long Muldrow Glacier. This glacier flows northward toward the developed road corridor west of the Eielson Visitor Center. Kahiltna Glacier is the longest in the park, at 71 km (44 mi). Ruth Glacier has a thickness of up to 1,160 m (3,850 ft), and moves about 0.95 m (3.1 ft) per day through the deepest gorge in North America (National Park Service 2009).

To the north of the Alaska Range is a series of east-west-trending foothill ridges ranging from approximately 610 to 1,372 m (2,000 and 4,500 ft) in height; the widths of these foothills range from 5 to 11 km (3 to 7 mi). The

parallel foothills extend eastward from the Kantishna Hills north of Wonder Lake. They are separated by broad, flat, sediment-filled glacial valleys that drain the region from south to north. Northward-draining rivers include the Foraker, Herron, McKinley, Bearpaw, Toklat, Savage, Teklanika, and Nenana Rivers. To the south, across the divide posed by the Alaska Range, the Chulitna, Susitna, Tokositna, Kahiltna, Yentna, and Kichatna Rivers drain.

Like most of Alaska, the Denali area is composed of a series of accreted terranes (fig. 2). These terranes include bits of islands, oceanic crust, flysch basin deposits, and continental fragments that have been transported from elsewhere by large strike-slip faults and ancient oceanic plates. Because the terranes were too thick to subduct beneath the North American continent into the Aleutian trench, they collided with the edge of the continent, squeezing ocean basins, island arcs, and miniterranes between them in a complex wedge of thickened crust.

Volcanism, mountain building, and orogenesis often accompany terrane accretion. The terranes typically follow an arching pattern that trends roughly east-west,

paralleling the Gulf of Alaska coastline. In the park, the terranes are identified as packages of rocks bounded by faults, with distinctly different rock types, fossils, and other physical properties than surrounding terranes.

In the area of Denali, the oldest major accreted terrane is the Yukon-Tanana Terrane. This large mass was added to the continent about 225 million years ago. The Talkeetna Superterrane (Wrangellia, Alexander, and Peninsular Terranes)—also referred to as the Wrangellia composite terrane—slammed into the continent about 110 to 85 million years ago. These two terranes, as well as several smaller terranes (such as the McKinley, Pingston, Windy, and Dillinger Terranes), compose the bulk of the landforms in Denali National Park and Preserve (fig. 3). Smaller accretionary prisms, such as the Chugach, accreted about 67 million years ago; the Prince William accreted by 50 million years ago. The youngest regional-scale terrane, the Yakutat Terrane, is still attached to the Pacific Plate, traveling approximately 5 cm (2 in.) northward per year; thus, accretion is an ongoing process along the coast of southern Alaska.



Figure 1: Map of Alaska Showing major mountain ranges and rivers. Graphic compiled by Phil Reiker (NPS Geologic Resources Division).




Figure 2: Generalized cross section from north (south of the Brooks Range) to south (Gulf of Alaska) through central Alaska showing the juxtaposition of accreted terranes over the subduction zone between the Pacific and North American Plates. Note the location of Denali National Park and Preserve. Graphic by Trista L. Thornberry-Ehrlich (Colorado State University), adapted from Lillie (2005) fig. 11.7.





Figure 3: Map of Denali National Park and Preserve. See yellow box on park map for location of detail map. NPS graphics.

Geologic Issues

The Geologic Resources Division held a Geologic Resources Inventory scoping session for Denali National Park and Preserve on February 24–26, 2004, to discuss geologic resources, address the status of geologic mapping, and assess resource management issues and needs. This section synthesizes the scoping results, in particular those issues that may require attention from resource managers. Contact the Geologic Resources Division for technical assistance.

Introduction

Resource management issues are discussed in relative order of significance, with the most critical listed first. Potential research projects and topics of scientific interest are presented at the end of this section.

Glacial Issues

Glaciers and Glacial Geology

Glaciers are a major feature of the Denali National Park and Preserve landscape. Over 1 million acres (17% of the park's area) is covered with ice. This ice includes more than 400 glaciers, some 40 of which are named. Famed glaciers include the Kahiltna Glacier, which is 71 km (44 mi) long, making it the longest glacier in the Alaska Range. Ruth Glacier is locally the thickest in the park, filling a deep gorge. Eldridge, Traleika, Tokositna, Yentna, and Muldrow Glaciers are among the other well-known glaciers at Denali. Permanent snowpack above 2,100 m (7,000 ft) supports the flow of glaciers from elevations above 6,100 m (20,000 ft) down glacial valleys to elevations as low as 300 m (1,000 ft).

Some of the smaller glaciers at Denali display interesting characteristics. Most of the smaller valley or hanging glaciers in the park are unique because terminal areas are insulated by deep deposits of rock debris and dust. This lessens melting and allows the smaller glaciers to remain stagnant for longer periods of time than larger glaciers. Portions of the larger Muldrow, Ruth, Kahiltna, and Tokositna glaciers are also insulated by deep deposits of rock debris and dust (P. Haeussler, USGS, geologist, written communication, November 2009).

Glaciers have been, and continue to be, powerful agents of landscape change. They carve valleys and gorges out of solid bedrock much faster than rivers and streams. Glaciers leave vast deposits of till and moraines. Glaciers feed outwash flows and streams, and can dam streams to form temporary lakes. At Denali National Park and Preserve, glacially carved features include valley cirques, arête ridges, drumlins, expansive U-shaped valleys, and horns. The widespread glacial deposits include: lateral, medial, and terminal moraines (less than 100 m [325 ft] thick); ablation and lodgement till (less than 10 m [32 ft] thick); eskers and advance and recessional outwash; and rock glaciers (potentially 100s of meters thick).

Rock glaciers are masses of talus that accumulate locally to the point where they retain moisture, freeze, and move

slowly downslope (Brease 2004). Blocks of glacial ice left stranded by glacial retreat can become buried by outwash deposits. The ice then melts, forming small, round kettle lakes such as Chilchukabena Lake near the northwest boundary of the park. Wonder Lake, at over 82 m (268 ft) deep, is the largest kettle lake at Denali. Pro-glacial Lake Moody, which formed by the damming action of ice and moraines in front of melting glacial ice, is also a glacial remnant at Denali.

Historical glacial activity in the area is recorded in the rocks, deposits, and vegetation patterns at Denali National Park and Preserve. Four distinct periods of widespread glaciation are recognized in the region of the park. Evidence of former glacial maxima exists on the north side of the Alaska Range, where, beyond the existing glaciers, moraine and outwash deposits extend into the foothills belt and cover large areas of bedrock, possibly filling vast areas underlying present-day river valleys through the unglaciated foothills section. The Wisconsin glaciation (Wisconsinan stage) was the last major advance of glacial ice in North America. Pork Chop Pond, a Wisconsinan kettle lake, contains thick layers of sediment (cored in 1996) that include aeolian sediments, peat, microlaminated muds, tephra (volcanic ash), and other organic deposits. These sediments record late Wisconsin-glacial and following interglacial climate (Axford and Werner 1997).

Glaciers and Climate Change

As stated by the Intergovernmental Panel on Climate Change: "Warming of the climate system is unequivocal, as is now evident from observations of increases in global average air and ocean temperatures, widespread melting of snow and ice, and rising global sea level." (IPCC 2007). This warming is very likely (more than 90% certain) related to anthropogenic greenhouse gas emissions (IPCC 2007).

Alaska has warmed at more than twice the rate of the continental United States. Alaska's annual average temperature has increased 2.0°C (3.4°F) while winters have warmed by 3.5°C (6.3°F) over the past 50 years (Karl et al. 2009). Projections reported by Karl et al. (2009) suggest Alaska's annual average temperature will rise between 2.8 and 7.2°C (5 and 13°F) by 2100, depending on emission scenarios. For more information regarding climate change effects and NPS response, visit the NPS Climate Change Response Program web site (<http://www.nature.nps.gov/climatechange/index.cfm>).

In response to the warming climate, glaciers are shrinking at the park. Most of the glaciers are consistently thinning, and only about 10% are surging (fig. 4) (P. Haeussler, USGS, geologist, written communication, November 2009). Surging refers to periods of relatively high-speed glacial flow following periods of slow flow. Most surging occurs on glaciers with northern exposure. The photographic record of the glaciers of Alaska spans more than 110 years. The earliest aerial photographs date from 1926 as part of a cooperative arrangement between the U.S. Geological Survey and the U.S. Navy. Systematic vertical aerial photography of Alaskan glaciers began in the 1940s and continues today (Molina and Sfraga 1999). These records provide a comprehensive data set and could be utilized to determine fine-scale glacial changes at Denali.

Glacial monitoring at the park began in earnest in 1991 as part of the National Park Service's Long-term Ecological Monitoring Program. This program initially focused on an "index" method—i.e., single point mass balance monitoring—but has since been supplemented with more detailed mass balance measurements on smaller glaciers, terminus monitoring on multiple glaciers, selected movement monitoring, depth measurements, and extensive photo documentation. Two index monitoring sites exist on Kahiltna and Muldrow Glaciers (Adema et al. 2003).

In 2004, data from past survey and research expeditions were revisited, new digital photographs were taken, and new stations were established for future comparisons of glacial positions (figs. 5, 6). Geographic coordinates, elevation, modern analog aerial images (using helicopters), and other accurate geographic information were incorporated into a parkwide GIS coverage of glacial extent from surveys during the 1950s through 2000s. Information included bidecadal coverages for individual glaciers (Karpilo et al. 2004). Efforts such as this could be expanded on a parkwide scale, incorporating more historic information, photographs, and modern technology to study glacial change at Denali National Park and Preserve. Karpilo (2009) suggested glacier monitoring techniques based on four "vital signs:" glacier mass balance, terminus position, area, and velocity.

Glacial Hazards

Glacial issues at the park extend to geologic hazards caused by glacial movement and glacial remnants. Relief in glaciated areas is often very steep and prone to erosion and mass movement depending on rock type. Glacial till often maintains very steep slopes (P. Haeussler, USGS, geologist, written communication, November 2009). Many deformed and altered rock units are present in glaciated areas of the park; these are prone to mass movements when exposed on slopes because of pervasive zones of weakness. When glaciers retreat up a valley, near-vertical walls are left with no structural support or stability. Deep, unstable, unconsolidated glacial deposits are also prone to mass wasting and intense erosion. Glacier terminal areas are attracting increasing numbers of visitors, and these areas could

pose icefall and rockfall hazards, as well as unstable trail bases. The vast extent of glaciers and icefields affect the local climate. Glaciers produce microclimates that cause local temperature cells to push moisture-rich air upward, creating localized weather aberrations.

Resource Management Suggestions for Glacial Issues

- Perform comprehensive studies on the history of glaciation at Denali. Potential research targets could include glacial debris and deposits, lichen growth, vegetation patterns, striations, and glacial lakes and lake deposits.
- Study glacial change at Denali by developing a glacier "state of health" program within the park (P. Haeussler, USGS, geologist, written communication, November 2009).
- To understand climatic effects on glaciers, conduct glacier dynamics research and perform mass balance studies. Promote contracting with the University of Alaska Fairbanks glaciology group (contact Chris Larsen) for airborne laser altimetry studies on a select group of glaciers to quantify glacier mass balance (P. Haeussler, USGS, geologist, written communication, November 2009).
- Study stability and safety issues related to glacial areas (W. Elder, NPS, geologist, written communication, December 2009).
- Compare glacial response at Denali with other known concentrations of glaciers in Alaska and worldwide to compare and contrast glacial response to global climatic shifts.
- Perform mass balance studies to determine reasons behind glacial surging.
- Perform glacial movement studies to determine differences between glaciers influenced to varying degrees by nonclimatic factors such as rock and dust coverage.
- Cooperate with other government agencies to obtain complete records of historical and ongoing glacial photography and measurement.
- Determine quantitative relationships between glacier size, orientation, location, depth, etc. and the degree of local microclimatic aberrations.
- Study Late Quaternary pollen, plant macrofossils, and insect fossils to determine past glacial and interglacial conditions (climate, vegetation, etc.) (Elias et al. 1996).
- Inventory the named glaciers in the park using 1957 and 1983 aerial and other photography and topographic maps to document spatial changes.
- Increase glacier profiling efforts, including monitoring terminal fronts, glacier thickness, and surge activity.

Earthquakes and Earthquake Hazards

Faults are prevalent features of the geologic structure at Denali National Park and Preserve. The large-scale Denali fault system runs through the entire park, parallel to the Alaska Range, extending more than 2,100 km

(1,300 mi) from the Yukon-Alaska border to the Bering Sea. The system includes the Denali fault (locally split into the Hines Creek and McKinley strands) and numerous subsidiary faults. Faulting at the park is active, with many fresh fault scarps showing evidence of recent movement. Fission track data indicate that Mt. McKinley is being uplifted along regional faults at rates of ~1 mm/year. More than 600 earthquakes per year are measured in the park area. More than 70% of these events are of magnitude 1.5 to 2.5; they commonly occur at very shallow depths, within 0 to 14 km (0 to 9 mi) below ground surface. The Alaska regional seismic network consisted of approximately 400 monitoring sites operated by various agencies. Seismic waveform data are now recorded digitally. The Alaska Earthquake Information Center (AEIC) is upgrading existing sites with digital broadband sensors (Ratchkovski et al. 2004).

Most earthquakes at Denali are relatively small (too small to be noticed by humans); however, even minor shaking can pose hazards of mass wasting, landslides, and debris flows on undercut, weakened, or unconsolidated geologic units exposed on slopes. On November 3, 2002, a magnitude 7.9 earthquake (consisting of multiple subevents), centered 48 km (30 mi) east of the park, caused severe local shaking. Reports of shaking and “unusual effects” from this earthquake were received from distances up to 3,500 km (2,200 mi) across western Canada (Cassidy and Rogers 2004). For the 30 years prior to this event, only four events greater than magnitude 3 were recorded. Leading up to and following this large earthquake, seismicity in the area increased. It is estimated that close to 14 years will be required for local seismicity rates to return to background level (Ratchkovski et al. 2004). According to the AEIC website (<http://www.aeic.alaska.edu/>), its system of over 400 sites currently reports about 20,000 earthquakes each year (well over 100 earthquakes in a few days) across the network. The AEIC site lists the location, depth, distance from surrounding communities, and magnitude for each event. The website frequently notes central Alaska for earthquake activity.

The fault rupture associated with the 2002 event was complex, beginning on variably dipping fault segments and continuing along vertical strike-slip segments (Eberhart-Phillips et al. 2003; Aochi et al. 2005). An unusual pattern of landslides and liquefaction resulted from the earthquake. The landslides were concentrated as rockfalls and rockslides in a narrow (~30-km [19-mi]-wide) zone that straddled the more than 340-km (211-mi) rupture zone. This pattern is consistent with strong, low-frequency shaking with only moderate acceleration levels. These rockslides were most visible at Black Rapids Glacier, Gakona Glacier, West Fork Glacier, and McGinnis Glacier (Harp et al. 2003; Schwartz et al. 2005).

Conversely, liquefaction effects (sand blows, lateral spreads, and broad settlement) within alluvial deposits of streams in and adjacent to the central Alaska Range were much more widespread, extending for several hundred km east of Fairbanks (Harp et al. 2003). This pattern is consistent with a long duration of shaking. Most

structural damage was focused 160 km (100 mi) east of the park, but the Denali fault system is capable of producing a similar earthquake in or closer to the park. This could have catastrophic effects on visitor use facilities, park infrastructure, the Trans-Alaska pipeline, and geologic landforms.

Widespread aftershocks from the large 2002 earthquake east of Denali increased the number of events recorded by the AEIC—from 1,000 to over 16,000—for a similar period (Ratchkovski et al. 2004). Thus, even if earthquakes are not focused within park boundaries, regional seismic episodes have far-reaching effects that could impact park resources. Future development plans should take into account the earthquake-related hazards of liquefaction, lateral spreading, rockfalls, landslides, and fault ruptures. In addition to these potential problems, the subgrade soils and permafrost (possibility of melting) in the Denali area pose further geotechnical design concerns in the event of an earthquake, so more detailed research is needed to understand their response to intense ground shaking (Adamczak et al. 2004). Braile (2009) suggested the following “vital signs” for seismic monitoring: monitoring earthquakes, analysis and statistics of earthquake activity, analysis of historical and prehistoric earthquake activity, earthquake risk estimation, geodetic monitoring and ground deformation, and geomorphic and geologic indications of active tectonics.

Resource Management Suggestions for Earthquakes

- Continue to collaborate on studies by Alaska Earthquake Information Center, universities and the U.S. Geological Survey regarding seismic activity in the region.
- Add seismometers in the park, to augment those put in place by the Geophysical Institute at the University of Alaska, Fairbanks. Present locations include Wickersham Dome in the Kantishna Hills, Thorofare Mountain near the Eielson Visitor Center, and Mount Healy. Cooperate with the AEIC to increase the number of digital broadband seismic stations within park boundaries.
- Continue seismic and geodetic monitoring in the park, as well as the study of ancient earthquakes on known important faults, to understand the implications of an event on a particular fault and to establish recurrence intervals (P. Haeussler, USGS, geologist, written communication, November 2009).
- Perform studies to determine the paleoearthquake history along the Denali fault in the ~161-km- (100-mi)-long region west of the 2002 event.
- Perform detailed mapping of the Denali fault and other active fault traces across the park to better understand deformation through time.
- Study paleoseismicity (most recent events) along sections of the park’s faults using samples of gouge and radiocarbon dating of offset organic material.
- Support efforts to determine fault slip rates (e.g., Matmon et al. 2006).

- Study geomorphology, tectonic history, and fault orientation relationships, and perform thermochronology studies (apatite fission track, U-Th/He) to determine timing and characteristics of landform development.
- Perform vulnerability assessments of areas in which visitor health and safety are likely to be affected in the event of a large earthquake.
- Incorporate earthquake epicenter data into a GIS, including geologic data, slope data, and trail data, to determine areas susceptible to failure during a moderate earthquake.
- Promote a better GPS geodetic network in the park to provide information about seismic hazards, where active structures lie, and the larger-scale tectonic setting (P. Haeussler, USGS, geologist, written communication, November 2009).
- Promote 3-D ionosphere tomography studies using GPS data to constrain source and propagation of seismic waves in the Denali area (Garcia et al. 2005).

Mining and Disturbed Lands

Since the gold rush era of the late 1800s, Alaska has been associated with vast mineral wealth, with approximately 30 million ounces of gold recovered to date. Most ore-bearing veins were intruded into brittle faults and fractures within metamorphosed pelitic rocks, and are composed of quartz with minor chlorite, carbonate minerals, and white mica (Goldfarb et al. 1997). In 1903, gold was discovered in the area of what is now Denali National Park and Preserve (Norris 1998). Past mining activity was focused in the Dunkle Mine area in the Chulitna Terrane of the Alaska Range, the Mt. Eielson/Copper Mountain district (in the 1940s), and the placer gold deposits of the Kantishna Hills region of the Yukon-Tanana Terrane in the northern foothills area of the park (Van Maanen and Solin 1988; Metz et al. 1989; see summary at www.mindat.org). Mining interests included precious metals such as gold and silver, base metals (copper, lead, antimony, zinc, tungsten, arsenic) and coal (Metz et al. 1989). Abandoned cabins, debris, small tailing piles, assay trenches, adits, and small shafts are historical markers of the hard rock mining history in the region. Most of the historical underground mine features have been closed or have since collapsed (GRI scoping notes 2004).

No new claims have been made since the establishment of Denali National Park and Preserve in 1980. In 1985, litigation was intended to halt mining inside the park, and no legal mining has occurred since (Burghardt 1997; J. Burghardt, NPS Geologic Resources Division, geologist, written communication, May 2010). According to the NPS Abandoned Mine Lands database (maintained at the GRD in Denver, Colorado), as of May 2010, there were 28 patented claims and 9 unpatented claims within the park. The patented claims are underground mine and placer mine sites for commodities such as gold, copper, coal, limestone, and antimony. Many of the claims have been used to establish lodges and cabins rather than mines (J. Burghardt, NPS Geologic Resources Division, geologist,

written communication, May 2010). The Abandoned Mineral Lands program also maintains a list of various identified mine-related features within the park, including adits, equipment, surface mines, tailings, buildings, and other structures. Currently (September 2010), this database has 37 listed features, at 28 sites, many of which require remediation. An online mineral database website, www.mindat.org, contains records of nearly 200 mine-related localities within or near the park.

Occasionally, legal disputes arise from the mining claims within the park. The Stampede antimony mine is owned jointly with the University of Alaska, Fairbanks. Several mine areas (Gold King placer claims and Red Top mill) are CERCLA (Comprehensive Environmental Response, Compensation, and Liability Act; “Superfund”) sites. This designation involves assessing risks to human health and the environment posed by hazardous waste sites, identifying potentially responsible parties (PRPs), and seeks to find funding for remediation efforts (GRI scoping notes 2004).

Denali National Park and Preserve also contains vast deposits of well-sorted sands and gravels. Small-scale quarries and gravel pits exist within park boundaries for construction and maintenance of park roads and other infrastructure and visitor use facilities. The park’s Resource Stewardship Strategy for 2008-2027 lists monitoring and mitigating impacts to fluvial morphology and water quality as a result of park gravel acquisition operations among its resource management goals (Denali National Park and Preserve 2009).

Negative effects from mining are far reaching and long lasting. Impacts include disturbed floodplain areas from extensive placer mining. Placer mining involved removing riparian zone vegetation and topsoil, then sieving valuable minerals from gravels in the stream channel, active floodplain, and old terraces. This process leaves behind barren gravel piles devoid of any soil that would promote natural revegetation. There are channels down to bedrock in some areas (Densmore 2005). A concern of park resource management is that past placer mining activity could result in changes in channel and floodplain morphology, streamflow, and streambed composition facilitated by erosion and/or deposition, that could affect aquatic and riparian ecosystem health (Van Maanen and Solin 1988). Recent regulations require that topsoil be stripped and stockpiled away from the placer mining area, then reincorporated with the sieved gravels so that planted material used in reclamation has a substrate in which to grow (J. Burghardt, NPS Geologic Resources Division, geologist, written communication May 2010). Remediation of old and/or abandoned mines is ongoing. Methods include rebuilding the floodplain areas, restructuring the stream channel (fig. 7), and recontouring terraces, as well as planting species such as *Alnus viridis* (green alder) for nitrogen fixation within new soils. Replanting along Glen Creek watershed (mined from 1906 to the mid 1970s) demonstrated the ability of the *Alnus viridis* species to accelerate the rate of succession by stimulating the growth of woody dominant riparian species such as *Salix alaxensis* (felleaf willow) (Densmore 2005). For many

areas, reclamation is limited by factors such as steep slopes, a lack of a local topsoil source from disturbed areas, an acid-generating sulfide substrate not amenable to revegetation, and remoteness (Burghardt 1997).

Preliminary studies suggest that differences in physical properties of mined and non-mined reaches of several streams were relatively minor; however, the streambed composition below mined areas tended to be enriched in fine-grained material (34 mm average mean particle size vs. 65 mm) and exhibited less variation in mean particle size (Van Maanen and Solin 1988). Along Eldorado Creek, mining and the operation of heavy equipment throughout the area, and particularly at the antimony mine at the headwaters of Slate Creek, have enhanced the flow of natural iron-rich springs (Burghardt 1997).

Placer mining forces larger particles to the streambed surface, which is contrary to natural settling. This could affect spawning habitat for fish as well as aquatic plant distributions. Along Rainy Creek in the Kantishna Hills region, the entire stream had been diverted from its natural channel during mining operations involving the construction of settling ponds. This resulted in extreme erosion of soil and fine-grained floodplain material as well as damage to pre-existing vegetation (Van Maanen and Solin 1988).

Associated with mining are exposures of hazardous materials in mine tailings. Mine tailings and other ground disturbances expose bedrock and crushed rock material to weathering. This leads to water and soil contamination by heavy metals and sulfides due to acid mine drainage. The Banjo Mine (a small-scale gold mine) has 500 cubic yards of mine tailings that contain leachable arsenic and lead, as well as antimony and cadmium that exceed drinking water standards and hazardous waste toxicity limits. Stampede Mine, a 20-acre site, maintains a fine-grained mine tailings dump adjacent to Stampede Creek. This mine is associated with dissolved antimony concentrations in the creek that exceed federal drinking water standards (ADEC 2006).

The Slate Creek antimony deposit, associated with historical mining operations within the park between Moose Creek and McKinley River, has drainage with pH values of 2.7 to 5.8, as well as concentrations of sulfate ions, arsenic, iron, manganese, nickel, antimony, and total dissolved solids that exceed State of Alaska drinking water standards. At the mouth of Slate Creek, some 3 km (2 mi) downstream, manganese and antimony contents exceeded drinking water standards. Stream sediment, soil, and rock samples around the mining area were enriched in arsenic, manganese, and antimony. In addition, sparse to no vegetation exists in the active stream channel and aquatic algae are conspicuously absent (Eppinger et al. 2000).

Mining operations introduce foreign objects and substances to the landscape, including petroleum drums, mercury, and other chemicals for gold extraction. Several areas in the Kantishna Hills region (Gold King #1-15-Area#1; Comstock #1-8-Area#5; Discovery #1-3-Area#7; Trommel Wash Plant-Area#8; Antimony Lode

Mining Area#9; Caribou Howtay #11b-24-Area#10; Glacier Assoc.#1-5-Area#12; Caribou Howtay #1-6-Area#13; Rainy Creek #1-3&6-8-Area#2; Liberty #53-Area#3; and Stampede Mine) have petroleum hydrocarbon contamination arising from miscellaneous drums, spills, and soil stains associated with the operations. The Glen Creek area contains mercury and diesel contamination from placer mine operations (ADEC 2006). A project to close a hazardous mine opening within the park was included in the 2009 American Recovery and Reinvestment Act (ARRA).

Resource Management Suggestions for Mining and Disturbed Lands

- Develop a mining history interpretive program focusing on the geology responsible for the valuable mineral resources in the Denali area.
- Continue reclamation efforts, including revegetation and remediation of streams and riparian (floodplain) environments. Increase plantings of *Alnus viridis*, which accelerates the rate of succession by stimulating growth of woody dominants (Densmore 2005).
- Collect continuous records of streamflow that relate changes in channel configuration and bed elevation to sediment transport, erosion, and deposition in mined and non-mined stream reaches for comparison.
- Monitor suspended sediment concentrations and fine bed material in mined and non-mined streams to help determine ecological effects of placer mining.
- Monitor water quality in mine-affected areas to determine degree of improvement, and target areas in need of further restoration. Tracking water chemistry changes through time would help resource managers understand the long-term effects of disturbed lands (P. Haeussler, USGS, geologist, written communication, November 2009).
- Continue to support environmental-geochemical studies of different mining areas, especially those located near springs and/or streams.
- Contact the NPS Geologic Resources Division regarding the history of mining at the park; the current status of mining operations, ore contents, reclamation efforts, and claim litigation and acquisition; and other mine-related queries.

Permafrost Issues

Unique to high altitudes and latitudes, permafrost is present discontinuously in the arctic climate of Denali National Park and Preserve. This frozen layer can persist to depths of up to 610 m (2,000 ft) below the surface. At Denali, the permafrost is defined as containing soil and rock that have been at a temperature of 0°C (32°F) or colder for two or more years.

Osterkamp and Jorgenson (2009) described the rationale for monitoring permafrost conditions and processes, including examples and case studies from Denali National Park and Preserve. Clark and Duffy (2003) mapped permafrost distribution within the park. Their

map shows that permafrost is present across the entire northern side of the park, but only locally south of the Alaska Range. Presenting a more regional perspective, the International Permafrost Association has published a digital map that summarizes the distribution and properties of permafrost and ground ice in the circum-Arctic region (Zhang et al. 1999). Permafrost is a major controlling factor on surface water flow, and therefore on vegetation and habitat. Permafrost exerts an influence on the fluxes of carbon and nitrogen from terrestrial to aquatic ecosystems coincident with fire disturbances (Petrone et al. 1999). It appears that leaching of carbon and nutrients from the burned ash layer is greatest in the active layer over permafrost, and this is marked as increases in these chemical components within the watershed (Petrone et al. 1999). Abundant water is available in the active layer above the permafrost because it cannot seep into the impermeable permafrost below (Brease 2004).

Permafrost also acts as a natural barrier to the downward flow of water, as well as against potentially harmful contaminants such as oil or diesel spilled from fuel storage tanks. Industrial users are interested in this potential use of natural permafrost in lieu of synthetic secondary liners (McCauley et al. 1999). Contaminants could pool atop the surface of the permafrost, causing a local long-standing concentration.

A delicate heat balance exists between permafrost and the layer above it, which freezes and thaws with seasonal temperature change. The balance of this system is sensitive to changes in the vegetative cover and mat and snow depth, and is dependent on the structural and compositional characteristics of the freeze-thaw layer (Brease 2004). These parameters can significantly affect the local thermal regime, resulting in changes at ground level such as frost heaving, polygonal ground patterns, and sags. Extreme conditions or anthropogenic alterations in the right geologic setting can cause melting of permafrost. Geologists are particularly concerned about the effects of global climate change (global warming) on permafrost in subarctic ecosystems. Permafrost is warming throughout Alaska; temperatures have increased throughout the state since the late 1970s as summarized by Karl et al. (2009). Ongoing monitoring is necessary to determine the extent to which this is occurring at the park. Osterkamp and Jorgenson (2009) suggested methods for monitoring permafrost conditions and processes utilizing thermal state and physical conditions “vital signs.”

Melting of permafrost can have dramatic effects on the landscape at Denali. The term “thermokarst” describes a specific landform that develops when permafrost is partially or totally melted. Thermokarst is a highly irregular surface expression that forms variously shaped polygonal depressions (patterned ground), altoplanation (long, smooth, flat ridges), and ice wedges (fig. 8). If melting of ice-rich permafrost (ground ice content greater than 20% by volume) occurs, widespread irregular subsidence of up to 2.0 m (6 ft) of the ground surface could result in significant impacts on the hydrological and ecosystem processes throughout the

region. In areas of abundant ice, thawing would result in river bank collapse, creation of thaw lakes, and other thermokarst features (Zhang et al. 1999).

Permafrost is very sensitive to human disturbance. Thermokarst has been created by humans in other parts of Alaska during construction projects. Accompanying thermokarst development, increases in solifluction, or soil movement, atop the frozen layer are more pronounced when permafrost is melted. This situation can cause heaving, sagging, slope creep, soil slumping, and deep erosion at the surface during successive periods of freeze-thaw heaving in the active layer (periglacial milieu) (Brease 2004). Hillslope stability becomes a safety concern. Like other subsurface movements or disturbances, solifluction can be detrimental to buried cables, wastewater treatment pipes, sewers, septic systems, utility poles, paved surfaces, trails, buildings, and roadbed foundations. Five large areas of roadway have been affected. Extensive knowledge of permafrost extent, properties, and processes will help resource managers decide where to develop road areas in the future and how best to maintain existing roads.

Resource Management Suggestions for Permafrost Issues

- Increase monitoring and develop a soil temperature monitoring network.
- Work with the National Resource Conservation Service soils mapping program to create a derivative map that includes different types of permafrost. A soil survey geographic database has been completed for the park as part of the NPS Soil Resources Inventory program (National Park Service 2006). Clark and Duffy (2003) produced a permafrost distribution map for the park.
- Increase the installation of thermistors (installed 35 m [115 ft] deep) along road corridors for permafrost monitoring.
- Investigate permafrost and ground ice conditions as well as their relation to the present climatic conditions and potential response to climatic change. Create a model of ground surface response to permafrost thaw to better predict ensuing ecosystem changes.
- Install additional inclinometers and piezometers along roadway slopes, as well as on slope areas near visitor use facilities or other identified vulnerable reaches.

Surface Water Issues

Snow and rainfall are channeled into three major systems at Denali National Park and Preserve. The first system is the series of glaciers, ice, and perennial snowfields. The second is the groundwater aquifers. The third includes all the streams, rivers, creeks, lakes, and ponds present on the landscape in Denali, including surface ice on lakes, which is a significant ecosystem factor.

There are approximately 400 streams and rivers in the park. Major waterways include the Foraker, McKinley, Kantishna, Toklat, and Teklanika Rivers on the northern slopes, and the Yentna, Kahiltna, and Chulitna Rivers

flowing south. The Nenana River marks the eastern boundary of the park. Due to continued regional uplift and plentiful water and sediment supply, these streams are aggrading in some areas and downcutting in the upland areas.

Ecosystem processes on the river floodplains are closely linked to fluvial processes and controlled by climate. Fluvial processes include flooding, sedimentation, and erosion, which interact with biotic processes such as seed dispersal and seedling establishment, and thus determine temporal and spatial development of riparian zones (Adams 1999). Floodplains with regular influxes of fine-grained silt and well-drained river terrace environments are rich ecosystems at Denali, supporting vast forests and lowland species. It is important that resource managers understand the fluvial processes that influence vegetation succession, including silt deposition, erosion, and river flow patterns that regulate flood events, as these processes are intimately linked with climate (Adams 1999).

Studies within the Rock Creek watershed targeted for long-term ecological monitoring revealed that streamwater chemical compositions, including elevated levels of Ca^{2+} , Mg^{2+} , and SO_4^{2-} , are directly influenced by underlying geology. In contrast, streamwater concentrations of nutrients and dissolved organic carbon are regulated by biological activity. Vegetation patterns are controlled by soil development, which is strongly related to geologic processes of weathering and alteration in addition to fluvial processes such as erosion and flooding. The productivity of the Rock Creek watershed is limited by physical factors such as unstable channel morphology, increased stream discharge, and organic matter retention (Popovics et al. 1999). Studies of this nature could be applied to many other watersheds at Denali to begin to quantify geologic controls on streamwater quality and riparian zone ecosystem health.

The park contains classic examples of braided streams (fig. 9). These streams typically (but not always) flow from glaciers, and contain choking amounts of sediment, causing them to split and crisscross the landscape. Not all streams at Denali are cloudy glacier-fed streams; clear streams that develop from springs and precipitation contain much lower levels of sediment. Turbidity and suspended sediment are greater in glacier-fed streams from the northern slope than from the southern slope. Levels of pH, alkalinity, conductivity, and ionic concentrations are also elevated in northern streams. These differences in chemical and sediment characteristics of streams at Denali are controlled by bedrock geology; marine sedimentary bedrock yields higher dissolved ion concentrations than resistant granitic bedrock (Edwards et al. 2000).

The wet northern slope area contains a unique wetland hydrology with shallow glacial lakes, bogs, ponds, and taiga. This flat, boggy, tundra area comprises approximately 20% of the park's land area. Some areas developed over millennial timeframes (Mann 1999). Long-term peat development records climatic change

over time through pollen, plant debris, and other remains.

Resource Management Suggestions for Surface Water Issues

- Contact the NPS Water Resources Division for assistance with surface water issues. The USGS Water Resources Discipline is an additional source of technical assistance.
- Inventory lakes, ponds, and other small water bodies, including bogs and wetlands.
- Conduct a comprehensive survey determining a vulnerability index for flooding in areas used by backcountry campers, park infrastructure, and other visitor use facilities.
- Identify floodplain areas, such as those around Kantishna lodges, for remediation.
- Perform studies linking spatial variability, climate, and fluvial processes with vegetation patterns and succession.
- Perform studies combining soils characteristics and detailed geologic mapping, soil water flow rate and chemistry, stream retention of organic matter, and streamwater chemistry to understand the relationships between soil, underlying bedrock, channel morphology, and streamwater quality for various watersheds.
- Monitor stream morphology, focusing on areas near gravel mines and damaged floodplain areas (from placer mining). Lord et al. (2009) have suggested techniques for monitoring stream systems.

Paleontological Resources

According to the park's Resource Stewardship Strategy for 2008-2027, paleontological resources are important resources for park management and visitors, and the park has a legal mandate to protect such important, nonrenewable resources. The 2009 Paleontological Resources Preservation Act directs the Secretaries of Interior and Agriculture to implement comprehensive paleontological resource management programs. The NPS and other federal land managing agencies are developing joint regulations associated with the Act (J. Brunner, NPS Geologic Resources Division, policy and regulatory specialist, personal communication, May 2010).

Denali National Park and Preserve contains many fossiliferous geologic units. Most of these are from the warm shallow seas present throughout much of the late Paleozoic and Mesozoic eras. Some of the accreted terranes traveled north from lower latitudes where marine life flourished. Fossils include invertebrates (as the dominant type), marine mollusks, insects, plant debris, and abundant trace fossils. A brachiopod, *Myrospirifer breasei*, was named after park geologist Phil Brease. Fossils found outside the park, in the same geologic units that exist in the park, that might also be present at Denali, include dinosaur remains of hadrosaurs, pachycephalosaurs, theropods, tyrannosaurids, ceratopsians, and a variety of vertebrate

trace fossils (Gangloff 1999; W. Elder, written communication, December 2009). Paleontological work by Brease et al. (2009) revealed the presence of dinosaur, plants and trace fossils in abundance within certain geologic units in the park (figs. 10, 11). Thousands of trace fossils of fish, pterosaurs, theropods, hadrosaurs, birds, and terrestrial and aquatic invertebrates are preserved in the lower Cantwell Formation, making this one of the best-preserved Late Cretaceous polar continental ecosystems in the global paleontological record (Brease et al. 2009).

At Denali, fossils occur in rocks which span the Paleozoic, Mesozoic, and Cenozoic eras. The boundary between the Cretaceous and Tertiary periods, commonly referred to as the “K-T boundary,” is present in the park. This yields a unique opportunity to determine the high-latitude effects on the paleoecosystem during the mass extinction that occurred at the end of the Cretaceous Period.

Paleontology work in the park has been mostly limited to the Healy quadrangle, with some additional fossil finds in the Talkeetna quadrangle on the south side of the range crest by Reed and Nelson (1977) (P. Haeussler, USGS, geologist, written communication, November 2009). Much of the park area remains to be surveyed and inventoried for paleontological resources. Field-based inventories are ongoing. A literature-based paleontological resource summary for the central Alaska Inventory and Monitoring Network is currently underway with expected completion in late 2010 or early 2011. The network summary is part of a national effort to systematically research fossil occurrences in NPS areas Servicewide. Conducting additional large-scale geologic mapping would inevitably reveal additional fossils. The park is currently developing a Paleontology Management Plan to address this vast resource (Brease et al. 2009; G. Adema, NPS, Denali National Park and Preserve, personal communication, July 2010). Santucci et al. (2009) identified “vital signs” for monitoring in situ paleontological resources: geologic and climatic variables affecting natural erosion rates, catastrophic geologic processes or geohazards, hydrology and bathymetry (i.e., changes in water level affecting resources near water bodies), and human impacts.

Resource Management Suggestions for Paleontological Resources

- Perform a comprehensive paleontological inventory at the park, including documented locations, abundance, ease of access, risk factors and disturbance, baseline conditions, fragility, and any necessary protection measures (Denali National Park and Preserve 2009).
- Study timing relationships of various deposits across the park using index fossils and locality studies. Focus on the Cantwell Formation for bio-stratigraphic work.
- Cooperate with researchers in creating a paleogeographic atlas of Alaska using invertebrate fossil records, for example.

- Perform palynology studies to research the Cretaceous-Tertiary stratigraphic boundary in the park area.
- Use paleontology to understand the spatial and temporal relationships between the different terranes and subterraneans present in the park region.
- Create paleontology interpretive programs to increase visitor awareness of the fossil resources at Denali.
- Increase efforts started by the geoscientist in the parks program to produce a comprehensive paleontological database (using Microsoft Access), expanding on the more than 200 sites already entered. Obtain GPS data for sites, as well as ground truthing, and further description and characterization. Information should be incorporated into the park’s GIS.

Geothermal Activity

Denali National Park and Preserve is located near an active plate margin, above a deep subduction zone, making it a prime area for geothermal activity. It was one of the original 22 national parks considered for geothermal resource management, but was not one of the 16 selected for extensive research (K. Moss, NPS Geologic Resources Division, environmental specialist, written communication, December 2009). Hot and warm springs are present throughout the park area, including Windy Creek and Wigane Creek. Cold meteoric groundwater trickling through the subsurface is heated by rock deep below the earth’s surface. This heating causes the water to expand; it is then usually forced upward under intense pressure to emerge as heated pools. There is no inventory of, or monitoring plan for, geothermal features within the park. These are needed data sets for park resource managers.

Resource Management Suggestions for Geothermal Activity

- Obtain water chemistry data for geothermal features along the south side of the park, focusing on the Windy Creek area.
- Inventory travertine deposits.
- Perform studies on the hydrogeology of Wigane Creek. This upwelling water feature could be geothermal, but the nature of the system is unknown.
- Inventory all hot spring features in the park. Map locations digitally for incorporation in the park’s GIS.

Slope Processes and Erosion

The geologic processes of erosion are prevalent at Denali National Park and Preserve. Glaciers are remarkable agents of erosion, and sheet runoff, streams, and rivers all carve channels and valleys into the landscape. Topography provides opportunities for erosion (P. Haeussler, USGS, geologist, written communication, November 2009). When unconsolidated sediments—such as slope deposits, glacial till and moraines, and alluvium, as well as altered and/or deformed bedrock—are exposed on moderate slopes, the potential for erosion and mass wasting increases. Some volcanic and sedimentary units are quickly altered to shrink-and-swell clays (minerals that swell when water-saturated and

shrink upon drying). This constant change in volume undermines the integrity of the rocks.

The area's earthquake activity increases the risk of catastrophic failure on park slopes. As described above in the seismicity section, ground shaking related to a single large earthquake event can produce widespread effects, including an increase in the intensity and quantity of smaller-scale seismic events that can induce slope failure.

An earthquake in 1912, located near the intersection of the Denali fault and the Richardson Highway, destroyed a climbing route on Mt. McKinley used in the 1910 Sourdough Expedition that allowed quick ascension of the summit of North Peak (Carver et al. 2004; Brease 2004; P. Haeussler, USGS, geologist, written communication, November 2009). On steep, unconsolidated slopes, as well as slopes of deformed rocks with cleavage oriented parallel to the slope, large slumps and slides can be triggered by small seismic events. Vulnerable areas can be assessed and identified based on the composition of the slope forming material, degree of slope, and orientation to known active faults.

Along the entrance road to the park, where it parallels the Hines Creek fault, landslides are relatively common after heavy rainfall. The water-saturated and heavy rocks and soil slide down the slope in slumps of ice-cored, lobe-shaped masses called solifluction lobes (Gilbert 1979). Much of the park road is built on unconsolidated glacial and fluvial deposits that erode easily and slump.

Large landslides often display hummocky topography, which is a very irregular and hazardous trail base for visitor use. Several slides—one older and vegetated, and three younger and unvegetated (not visible on 1949 era aerial photographs)—exist along the trail to Triple Lakes. Another slide, only 1 km (0.6 mi) above the park road, pushed the course of Tattler Creek southward. In 1953, massive landslides of sheared basalt flows dammed Stony Creek, creating Bergh Lake (Gilbert 1979). These slides attest to longstanding and continuous slide activity in the area.

Rockfalls are prominent components to mass wasting at the park. Large rockfalls exist on the south side of the upper Buckskin Glacier and off the east face of the Eye Tooth (fig. 12). Rockfall events are inherent in areas of steep topography, freeze-and-thaw cycles, bare slopes, and rock faces (P. Haeussler, USGS, geologist, written communication, November 2009).

Snow and ice avalanches are a year-round threat to visitor safety, park infrastructure, and backcountry areas. Snow avalanches are a significant hazard at the park. Earthquakes can trigger snow avalanches, much in the same way as rockfalls or landslides. Larger avalanches with longer runouts can be generated with a larger trigger mechanism such as an earthquake (P. Haeussler, USGS, geologist, written communication, November 2009).

Other features at risk for erosion and mass wasting include rock glaciers, glacier fronts, terminal moraine

slopes, and eskers (figs. 4, 19). Park resource management attempts to mitigate the effects of erosion and mass wasting along the most vulnerable and exposed areas. Engineering along Lake Moody attempts to alleviate some of the geologic hazards in the area. Areas along park roads and trails subjected to repeated subsurface sliding are targeted for reinforcement. Wieczorek and Snyder (2009) suggested the following “vital signs” for monitoring slope processes: types of landslides, landslide triggers and causes, geologic materials in landslides, measurement of landslide movement, and assessing landslide hazards and risks.

Resource Management Suggestions for Slope Processes and Erosion

- Catalog mass failures. As new failures occur, their location and type should be noted and incorporated into the park's GIS. Past mass failures should also be categorized, catalogued, and incorporated in the park's GIS (P. Haeussler, USGS, geologist, written communication, November 2009).
- Perform slope vulnerability studies to identify reaches within the park susceptible to massive landslides in the event of an earthquake. Target unconsolidated deposits and heavily deformed geologic units.
- Determine what effects, if any, social trail development has had on erosion and seasonal runoff in the park.
- Identify areas where park infrastructure is threatened by erosion and mass wasting.
- Incorporate spatial slope data into the park's GIS to correlate with underlying geologic units and structure, as well as vegetation patterns and hydrologic information to determine areas vulnerable to extreme erosion and mass wasting.

General Geology Issues and Research Possibilities

The remoteness, arctic climate, ruggedness, and sheer size of Denali National Park and Preserve poses significant challenges to geologists and other researchers seeking to study the landscape. However, these same characteristics also serve to preserve a living laboratory and a unique opportunity to study pristine alpine-arctic environments, active glaciers, active seismicity, permafrost, paleontology (fig. 13) and active margin terrane accretion.

Resource Management Suggestions for General Geology Issues and Research Possibilities

- Perform larger scale, detailed geologic mapping in a digital format for incorporation in the park's GIS.
- Perform regular surveys and repeated monitoring of soil conditions to determine the lasting impacts of off-road vehicle and snowmobile use on soil compaction and wetland/bog ecosystem health.
- Identify and photograph the park's rare or unique geological features and type sections. Periodically monitor their condition (Denali National Park and Preserve 2009).
- Perform an air quality and visibility change analysis, focusing on the introduction of industrial pollution,

arctic haze, and coal-fired electricity generation. Contact the NPS Air Resources Division (Denver, CO) for technical assistance.

- Cooperate with federal and local agencies for continuing geologic mapping efforts.
- Promote detailed studies of the geologic history and origin of the smaller terranes located along the Denali Fault and throughout the Kahiltna Glacier area.
- Develop additional geologic-themed interpretive programs to increase visitor awareness of the role

geology plays in the landscape evolution and ecosystem at the park.

- Map, describe, and characterize the significant loess deposits (windblown, glacially-derived silt) at Denali.
- Determine if caves exist in the Devonian and Silurian age limestones. If caves are present, perform a comprehensive inventory, digitally mapping locations to incorporate into the park's GIS.
- If needed, develop a plan to protect any existing cave resources as well as ensure visitor safety.



Figure 4: The Straightaway Glacier (shown) is one of many surge-type glaciers which issue from Mt. McKinley. Surging glaciers can move 10-100 times the normal flow rates. NPS Photo/Adema.



Figure 5: Repeat photography pair showing retreat of a glacier at the head of Hidden Creek in Denali National Park and Preserve. Upper photo was taken in 1911 by USGS geologist Steven R. Capps, lower image was taken on August 7, 2004 by Ronald D. Karpilo, Jr. Photos courtesy of Ronald D. Karpilo, Jr. (Colorado State University).



Figure 6: Repeat photography pair showing retreat of a glacier in the Teklanika River valley in Denali National Park and Preserve, Alaska. Upper photo was taken in 1919 by USGS geologist Steven R. Capps, lower image was taken on August 9, 2004 by Ronald D. Karpilo, Jr. Photos courtesy of Ronald D. Karpilo, Jr. (Colorado State University).



Figure 7: Denali has been steadily restoring abandoned mine lands like this project on Slate Creek. Encapsulated soil lifts guide a new channel away from exposed ore to improve water quality and aquatic habitat. NPS Photo/Adema.




Figure 8: Thermokarst development, like this example in the Toklat Basin on the north side of Denali, is an indicator of the dramatic and quick change occurring on the base permafrost landscape. NPS Photo/Yocum.



Figure 9: One of Denali's most dramatic features are the large, braided rivers like the Toklat river. Not only geologically striking, the Toklat provides a renewable source of gravel for maintenance of the Denali Park Road. NPS Photo/Adema.



Figure 10: Theropod dinosaur discoveries in the Cantwell Formation have excited considerable paleontological research. A theropod footprint trace fossil is shown here. NPS Photo/Brease.



Figure 11: Plant fossils, such as this fern, provide insight to the paleoecology of Denali. NPS Photo/Brease.



Figure 12: View of landslide debris on Buckskin Glacier with granite peak named The Moose's Tooth (3,150 m; 10,335 ft) visible in the background, Denali National Park and Preserve, Alaska, August 7, 2004. Photo courtesy of Ronald D. Karpilo, Jr. (Colorado State University).



Figure 13: Researchers investigate dinosaur fossils in Denali's Cantwell Formation. NPS Photo/Brease.

Geologic Features and Processes

This section describes the most prominent and distinctive geologic features and processes in Denali National Park and Preserve.

Mt. McKinley (Denali)

Mt. McKinley (known as “Denali” [“The Great One”] in the native Athabaska language) dominates the landscape of Denali National Park and Preserve, towering some 915 m (3,000 ft) above the neighboring peaks in the Alaska Range (figs. 1, 3, 14). Its highest point is 6,194 m (20,320 ft) above sea level, making it the tallest mountain on the North American continent. Mt. McKinley features greater relief than most mountains on Earth. For example, the Wickersham Wall on the northern slope of the mountain has approximately 5,500 m (18,000 ft) of vertical relief. This peak was first scaled in 1913 by a party led by Hudson Stuck. Today, features such as Browne Tower, Karstens Ridge, McGonagall Pass, and Denali Pass, not to mention the numerous icefalls and crevasses posed by the large glaciers in the area, still challenge climbers (Greenwood and Rowell 1980). The American Alpine Journal frequently describes climbing endeavors and routes within Denali National Park and Preserve.

Uplift of the Alaska Range began during the Jurassic and culminated in the Tertiary, some 60 to 30 million years ago. Recent studies suggest that the modern mountain mass in the immediate area of Denali was uplifted rapidly (~1 km/million years [0.6 mi/million years]) beginning ~6 million years ago (Plafker et al. 1989). This uplift was a result of interactions between the right lateral Denali fault (with a local morphology that includes a 22-degree bend) and a series of northeast-striking thrust faults, and changes in motion between the Pacific and North American plates (Haeussler 2005). This caused a space problem that forced crustal blocks upward (Fitzgerald et al. 1993).

Mt. McKinley is still rising approximately 1 mm (0.04 in) per year. Uplift of the range hinged on the convergence of several geologic factors, including subduction, faulting, terrane accretion, and plutonism (Fitzgerald et al. 1993). The emplacement of the Chugach and Prince William accretionary wedges and the Yakutat terranes, as well as assorted smaller terranes, involves extreme shortening (fig. 15) as well as thrust faulting, which thickens the crust and shoves large crustal blocks upward (Dumoulin et al. 1998). The buoyancy of this thickened crust leads to further uplift as it adjusts to approach isostatic equilibrium.

Subduction of the Pacific Plate beneath southern Alaska introduces water into the mantle, which then leads to partial melting and generation of magmas. At Denali, this led to plutonism and volcanism until about 38 million years ago. Since that time, subduction has continued, but there is no magmatism in Denali National Park and Preserve (P. Haeussler, USGS, geologist, written communication, November 2009).

Relatively buoyant and resistant granitic plutons form much of the core of the Alaska Range. These rocks are not easily worn away by erosion and weathering, thus allowing elevations to remain relatively high compared with other mountains of similar ages. The dominant peaks of the range—Mt. McKinley, Mt. Foraker, and Mt. Hunter—are composed of resistant granites. Differential erosion of less competent metasedimentary rocks of the surrounding mountains accentuates the height of these peaks (Fitzgerald et al. 1993). The biotite granite of the so-called McKinley pluton comprises most of the Mt. McKinley massif. It contains quartz, plagioclase, potassium feldspar, biotite, white mica, tourmaline, and other accessory minerals. Nearly 4 km (2.5 mi) of this igneous intrusive body is exposed within Denali National Park and Preserve, rendering it an ideal research target for petrologic and geochemical variations within a large continuous intrusion (West and Swanson 1995).

Denali Fault System

The Denali fault system is a 2,100-km- (1,300-mi)-long structure, trending from the Yukon border southwest toward the Bering Sea. The entire system connects with the Queen Charlotte fault and the Fairweather fault of southeastern Alaska toward British Columbia, forming a significant piece of the transform fault boundary between the Pacific and North American plates. The great arcuate system of northward-convex transcurrent faults that sweeps across Alaska is one of the most distinctive geologic features within the state (Moore et al. 1994). The Denali fault was first named and described by St. Amand in 1957. The surface trace of the fault is concave to the south, roughly paralleling the shape of the Gulf of Alaska. The curvature of the fault is about 70 degrees between the eastern and western portions. It is one of the largest and topographically best-expressed geologic features of south-central Alaska (Csejtey et al. 1982). This system now accommodates counter-clockwise rotation of accreting terranes by absorbing some of the relative motion between the Pacific and North American plates. The Tintina-Kaltag fault and Iditarod-Nixon Fork fault systems (fig. 16) (other large-scale, right-lateral, strike-slip systems), located north of Denali, accommodate some of this motion; as well, some faults to the south accommodate offsets today (Miller et al. 2002; Ratchkovski et al. 2004; W. Elder, written communication, December 2009). These margin-parallel, strike-slip faults have experienced multiple episodes of dextral motion since approximately 100 million years ago (Miller et al. 2002).

Estimates of right lateral offset along the Denali fault system are as little as several tens of km to as much as 600 km (370 mi) (Csejtey et al. 1982; Redfield and Fitzgerald 1993; Csejtey et al. 1995; Redfield and Fitzgerald 2000). The fault splits into two strands within

the Denali region. This split might have been due to a change in relative plate direction during the last ~50 million years from north-south to a more lateral east-west (Gilbert 1979; Cole et al. 1996). The northern branch is the Hines Creek fault; the McKinley fault is the southern strand of the Denali fault. Both faults are right-lateral motion, near vertical strike-slip faults. The northern side of each fault moves eastward relative to the southern block.

The Hines Creek fault traverses the northern reaches of the park. It roughly separates 350-million year old schist, as well as lower and middle Paleozoic metasedimentary rocks of the Yukon-Tanana Terrane from the Pingston and McKinley Terranes. It also bisects portions of the Paleozoic Farewell Terrane (a name used for the combined Dillinger, Nixon Fork, and Mystic Terranes) as well as granites of Mt. McKinley to the south that are Tertiary age (56 million years ago) (Csjetey et al. 1995; Dumoulin et al. 1998; Brease 2004). This fault passes near park headquarters and crosses the Alaska Railroad, extending westward, usually forming valley floors, for 80 km (50 mi). This fault is not considered to be active.

The McKinley strand of the Denali fault is active, and it passes through the central section of the park. It enters the park near Cantwell and trends eastward toward Anderson Pass and then beneath the Muldrow Glacier. (St. Amand 1957). This fault has displaced a pluton of 38 million year old Foraker granite (named for the second highest peak in the park) that is horizontally more than 40 km (25 mi). This corresponds to an average slip rate of 1 mm (0.04 in.) per year for the past 38 million years.

Both strands of the Denali fault are north of Mt. McKinley. The McKinley fault has significant reverse thrust motion associated with it—i.e., the fault surface dips slightly toward the south, with Mt. McKinley being thrust upward and northward along this surface (Ford et al. 2003). The Wickersham Wall on the northwest side of Mt. McKinley reflects the recent vertical component of movement along the fault (Gilbert 1979). Fault scarps along the McKinley strand, as well as local features such as angular grabens, en echelon (overlapped or staggered) ruptures, and “mole tracks,” exhibit a reliable history of Holocene movement across the fault system (Redfield and Fitzgerald 1993; Schwartz et al. 2005). These features also demonstrate the partitioned nature of movement between normal, thrust, and lateral (strike-slip) sense throughout such a major structure. The two fault strands coalesce east and west of Denali National Park and Preserve.

As the Alaska Range is being uplifted, reverse and transform motion is accommodated by the Denali fault system and smaller-scale normal faults and thrust faults. On the trail to Triple Lakes, several igneous dikes, intruding the Cantwell Formation, have been offset along several small faults. The hike along Tattler Creek shows evidence of at least 450 m (1,500 ft) of uplift within the Cantwell Formation (Gilbert 1979).

The intensity of deformation within the rocks through which the Denali and other faults pass is variable, but

penetrative foliations, well-developed lineations, and strained grains suggest extensive transposition along the strands of the Denali fault, in particular. Locally, the zone of observable fault deformation reaches a width of 5 km (3 mi). Cataclasites and zones of fault gouge from brittle deformation regimes cut mylonitic fabrics, recording earlier ductile deformation along the fault (White 1997). Rocks trapped between shifting fault blocks are pulverized, deformed, and crushed. This renders them easily eroded; thus, most faults are traceable on the surface as linear topographic depressions (Gilbert 1979). Much of the actual fault trace is covered in Quaternary age sediments and/or alpine tundra (White 1997), and is visible east of Anderson Pass (P. Haeussler, USGS, geologist, written communication, November 2009).

Kantishna Hills

The Kantishna Hills are tectonically active and among the most interesting geologic features of Alaska (P. Haeussler, USGS, geologist, written communication, November 2009). Precambrian to early Paleozoic crystalline metamorphic rocks (mica schist), intruded by swarms of Cretaceous to Tertiary dikes, underlie the hills (Prindle 1907; Ruppert et al. 2008). The northeast-trending Kantishna Anticline is a growing feature resulting from at least five deformational episodes. High-angle and thrust faults generally follow the trend of the anticlinorium, whereas conjugate folds and faults trend northwest (Bundtzen 1981; Salisbury and Dietz, Inc. 1984; Burghardt 1997). The southern end of the Kantishna Hills anticlinorium marks one of the most prominent areas of seismic activity in all of interior Alaska (Ruppert et al. 2008). It is growing at such a rate as to force rivers to divert around it (P. Haeussler, USGS, geologist, written communication, November 2009). The McKinley River flows around the southern end of the anticlinorium and through the seismic zone. The channel of this river changes from a well-developed braided channel to an incised meander morphology as the river approaches the southwestern part of the structure. North of the Kantishna Hills anticlinorium, the river abruptly reverts to a braided channel morphology (Ruppert et al. 2008). Active deformation, as the structure is propagating southwestward, is also responsible for reaches of convexity along the longitudinal stream profiles of McKinley River and Moose Creek near the Kantishna Hills (Ruppert et al. 2008).

The lode mineralization in the Kantishna Hills appears to be spatially related to the axis of the Kantishna Anticline. This happened where the structure (especially areas of brittle deformation) focused mineralization and the emplacement of locally high-grade mineralized veins (Bundtzen 1981; Salisbury and Dietz, Inc. 1984; Burghardt 1997). Superheated, saturated fluids preferentially flowed through the fracture zones, depositing mineralized veins that would later attract the interest of miners.

Accreted Terranes

As much as 90% of Alaska's current land mass is not part of the original North American craton, but was instead transported as a series of exotic terranes conveyed atop tectonic plates and accreted onto the continent (Stone 1984). These terranes include bits of islands, volcanic arcs, marine sediments, oceanic crust, and slivers of other continents that either obducted onto the continent or were too thick to subduct into the Aleutian trench. The central part of the Alaska Range in Denali National Park and Preserve is composed of three major separate terranes of varying scale (P. Haeussler, USGS, geologist, written communication, November 2009).

The terranes form long linear belts, and typically follow an arching pattern trending roughly east-west, paralleling the Gulf of Alaska coastline. Each separate terrane is bounded by faults, and is distinguished by differing rock assemblages, metamorphic grade, structural fabric, fossils, etc. Volcanism and mountain building (orogenesis) often accompany terrane accretion; at Denali, pervasive faulting (including strike-slip, thrust, and high-angle reverse faults) have jumbled the terranes, obscuring their chronologic order (Stone 1980). Several faults, including the West Fork, Talkeetna, Denali, Chitina Valley, Castle Mountain, and Totschunda faults, separate terranes in the Denali area (Nokleberg et al. 1994). The contacts between the various accreted terranes are also gave rise to hydrothermal metamorphism and magmatism, which in turn led to the development of gold veins in brittle faults and fractures (Goldfarb et al. 1997). Geologists use stratigraphic correlations, radiometric dating, paleomagnetism, detailed mapping, and other techniques to determine the relative sequence of accretion tectonic history. At least nine widespread magmatic episodes are represented by plutonic rocks scattered across Alaska (Miller 1994).

The names, designations, and numbers of identified terranes in the central Alaska Range have changed with more detailed geologic understanding of the area. Moore et al. (1994) provides one view of the tectonic assembly of the terranes of Alaska (beyond the scope of this report, which presents only a summary). The oldest terrane is the Yukon-Tanana terrane (and the Nixon Fork terrane, located west of Denali) that is composed of sequences of Precambrian through Paleozoic metasedimentary and metavolcanic rocks (Csejter et al. 1982). This terrane comprises much of central Alaska and is present within the northern areas of the park, north of the Hines Creek fault. Farther south, several smaller terranes were caught between the Yukon-Tanana terrane and the accreting Wrangellia composite terrane or Talkeetna superterrane (Wrangellia, Alexander, Maclaren, and Peninsular terranes) during the Cretaceous collision between the Talkeetna and the continent (figs. 2, 16, 17) (Csejter et al. 1982).

Farthest north among these smaller "miniterranes" is the Pingston terrane, which contains isoclinally folded, deep water mid-Paleozoic through Mesozoic, silty limestone, slate, and phyllite. The neighboring McKinley, Dillinger, and Windy terranes contain mixed Mesozoic flysch

rocks and pillow lavas (McKinley terrane), micaceous sandstone (turbidites), graptolitic shale, deep water limestone (Dillinger terrane), serpentinite, basalt, tuff, chert, limestone, and mixed flysch (Windy terrane).

Farther south, the Mystic terrane is composed of Mesozoic graptolitic shale, pillow basalt, shallow water limestone, sandstone, and chert. The Chulitna Terrane contains Mesozoic ophiolite rocks (oceanic crust), chert, volcanic conglomerate, limestone, and mixed flysch. The ophiolites of this belt were altered to serpentinite and contain clues to crustal suture (Roeske et al. 2005). The West Fork terrane contains chert, sandstone, conglomerate, and tuffaceous argillite. The nearby Broad Pass terrane contains chert, tuff, and blocks of limestone and serpentinite (Jones et al. 1980).

These small terranes are tectonically mixed with intervening Jurassic and Cretaceous flysch, deposited in deep basins that existed between the colliding terranes that collapsed and deformed upon collision (Csejter et al. 1982). The Kahiltina assemblage is one such flysch sequence. It likely depositionally overlapped the northern margin of the Wrangellia and Peninsular terranes, and is exposed in the northern Talkeetna Mountains and southern Alaska Range (Ridgway et al. 2002). Another noteworthy flysch sequence is the Gravina-Nutzotin belt, which overlies the Wrangellia and Alexander terranes (Nokleberg et al. 1994). These sedimentary basins might record the progressive development of a suture zone between the Talkeetna Superterrane (Wrangellia composite terrane) with the North American continental margin (Ridgway et al. 2002).

Younger large terranes and accretionary prisms, located to the south of Denali National Park and Preserve, include the Chugach and Prince William accretionary blocks and the Yakutat terrane (P. Haeussler, USGS, geologist, written communication, November 2009). The youngest regional-scale terrane, the Yakutat terrane, is still attached to the Pacific Plate, traveling approximately 5 cm (2 in.) northward per year; thus, accretion is an ongoing process along the coast of southern Alaska.

Glacial Features

Glacial processes carved much of the rugged topography at Denali National Park and Preserve. A glacier is a powerful agent of erosion, capable of profoundly altering the landscape over which it passes. Glaciers primarily erode by two distinct processes: plucking and abrasion (Dyson 1966). In plucking, the glacier actually quarries out distinct masses of rock, incorporates them within the ice, and carries them along; as the glacier moves forward, these blocks of rock are dragged or carried along with it (Dyson 1966). Abrasion involves glacial ice mixed with bits of rock and debris that actually scours the underlying bedrock as it passes over.

Cirques

By quarrying headward and downward at its mountain source, the glacier ultimately carves a steep-sided, bowl-shaped basin called a cirque or glacial amphitheatre. The

cirque is the first place that ice forms and the place from which it disappears last; thus, it is subjected to intense glacial erosion longer than any other part of the glacial valley. Once the glacial ice melts away, a body of water known as a cirque lake can form in the depression (Dyson 1966). Cirque lakes are not common at Denali because most of the glaciers are still active.

U-Shaped Valleys

Rock fragments of various sizes frozen into the bottom and sides of the glacier form a huge file or rasp, which abrades or wears away the bottom and sides of the valley down which the glacier flows. The valley thus attains a characteristic U-shaped cross-section, with steep sides and a broad bottom. This is in contrast to a stream-cut valley and its characteristic narrow, V-shaped profile. Many valleys in the park, once devoid of glacier ice, will possess this distinct U-shaped cross-section. The upper valleys of the Savage, Sanctuary, and Teklanika Rivers are U-shaped (Gilbert 1979).

The topography of the valley floor is not necessarily smooth, as it is usually contoured from erosion controlled by geologic properties of the bedrock. Steep drops or “steps” can mark valleys, between which the valley floor has a comparatively gentle slope. Such a valley floor is called a glacial stairway (Dyson 1966). These features result from the differences of erosional resistance between different rock types of the underlying geologic formations. A glacier will scour more deeply into a weaker rock, such as shale, schist, or altered volcanic rock, forming a “tread” (in contrast to the cliffs or “risers” formed by the erosion of stronger rocks such as massive sandstones, quartzites, or granites).

Hanging Valleys

The “tributaries” of glacial valleys, filled with smaller glaciers that feed into the larger ones, are known as hanging valleys. They form as a result of differences in erosional power between the smaller glacier and the larger valley glacier. The thicker the stream of ice, the more capable it is of erosion. Thus, a main valley deepens greatly, while the tributary valleys, with their smaller glaciers, cut down more slowly, leaving them hanging high above the floor of the main valley once the glacier ice melts (Dyson 1966).

Arêtes

Conspicuous throughout the Alaska Range are long, sharp ridges forming much of the backbone of the mountains poking above valleys filled with glacial ice. These features are known as arêtes, and owe their origin to glacial processes.

As the long valley glaciers enlarge their source cirques by cutting farther toward the axis of the mountain range, the mountain wall is finally reduced to a very narrow, steep-sided ridge, an arête. In the absence of glacier ice, knife-like ridges separate deep valleys at the park.

In some places, glaciers on opposite sides of a ridge can cut through the ridge, creating a low area known as a col (but usually called a pass) (Dyson 1966). At locations

where three or more glaciers have plucked their way back toward a common point, they leave at their heads a sharp-pointed peak known as a horn.

Braided Rivers

Most of the major rivers at Denali seem underfit for their valleys. An underfit stream appears to be too small to have eroded the valley in which it flows, commonly the result of drainage changes caused by glaciers. They flow, splitting and intertwining in wide, glacially eroded, U-shaped valleys. They are choked with excess sediment and follow steep gradients. Toklat River, visible from the park road, is a textbook example of such a river (fig. 9) (Gilbert 1979). When one stream channel fills with sediment, other channels form nearby. As streams drop vast piles of poorly sorted sediment, valley trains (long narrow accumulations of glacial outwash confined by valley walls) develop (Brease 2004).

Glacial Lakes

Most lakes throughout the park owe their existence directly or indirectly to glaciers and glacial processes. They can be divided into several main types, depending on their origin: cirque lakes, other rock-basin lakes, lakes dammed by glaciers and/or glacial deposits, kettle lakes, or glacial valley lakes.

Cirque lakes fill the depression plucked out of solid rock by a glacier at its source. These are not as common at Denali as in other glaciated parks because glacial activity is still so prevalent at the uppermost parts of most of the river drainages (Gilbert 1979). Other rock-basin lakes fill depressions created where glaciers moved over areas of comparatively weak rock. In all cases of cirque or rock-basin lakes, a bedrock dam contains the water.

Lakes held back by outwash deposits are dammed by mixed sediments and stratified gravel, which were contained in or washed out from glaciers when they advanced into the lower parts of the valleys. Moraine lakes are formed when a moraine deposit blocks a stream outlet. Some lakes at Denali are caused by glaciers acting as dams across pre-existing river valleys or outwash streams from another glacier. Glacial impoundment is recorded in the rock record as varved clay (distinctly laminated sediments including the upper, fine-grained “winter” layer reflecting quiet depositional setting) lacustrine deposits that formed in the transient glacial lakes. Glacial Lake Moody once existed in the Nenana River canyon when the river was dammed by glacial deposits following the Healy Glaciation and during the Riley Creek Glaciation (Gilbert 1979). Glacial valley lakes are often long and finger-like, filling the former U-shaped valleys once occupied by large glaciers. These are large-scale features that could be dammed by moraine deposits.

Kettle lakes and ponds form in response to glacier advance and retreat. Large pieces of ice often calve from the head of a glacier. These pieces are left stranded when the glacier retreats up the valley. When outwash streams deposit sand, clay, gravel, and other sediments around the block of stranded ice, the block is insulated and, once

melted, forms a small, typically round, pond in the midst of new thick glacial outwash. West of the Eielson Visitor Center, as well as north of the Teklanika Campground, numerous small lakes and ponds formed by melted blocks of ice in morainal deposits are visible (Gilbert 1979). Wonder Lake, near Kantishna, was formed by a combination of glacial erosion and ice block melting (Brease 2004).

Glacial Deposits

As the glacier ice moves, rock fragments continually break loose and can entrain into the ice itself. Some of the fragments are ground into powder as they move against each other and scour against the bedrock under the glacier. Many types of rock yield a milky gray powder when finely ground. Melt-water streams issuing from present-day glaciers are cloudy or milky from their load of this finely ground “rock flour.” Much of this silt is deposited in lakes where light refracting off the particles imparts a milky turquoise color to the water.

Glacial till is a term used to describe deposits of mixed rock left by glaciers. During glacier advance, huge amounts of rock debris are transported down glacial valleys. When the glacier stops advancing and begins melting and retreating up the valley, a large rock debris deposit, known as a terminal moraine, is dumped at the point of furthest advance (fig. 18). Terminal moraines help constrain the timing of glacial advance and retreat.

The McKinley River area contains one of the most complete glacial deposit sequences along the north flank of the Alaska Range at Denali National Park and Preserve. Included in this sequence are three pre-late Wisconsin age moraines. These are broad ridges formed when the Muldrow and Peters Glaciers coalesced and extended down the valley as a broad lobe approximately 50 km (31 mi) from present margins. Local deposits include a four-fold late Wisconsin moraine sequence (40 km [25 mi] down the valley) and four separate neoglacial moraines. The oldest of the neoglacial deposits is 3,000 years old; the most recent is dated within the past 100 years. These deposits display irregular topography, with some remnant ice present at depth within the deposit and little vegetation cover (Werner et al. 1990).

Terminal moraine deposits throughout the park are susceptible to intense erosion immediately following deposition, as well as the obscuring effects of a vegetative cover. Studies of moraine deposits at Denali reveal at least seven major periods of glacial advance beginning around 2 to 3 million years ago. Locally, these are called (from earliest to latest): the Teklanika (Late Tertiary, >2 million years before present [ybp]); Browne (150,000+ ybp); Bear Creek (125,000 to 150,000 ybp); Dry-Lignite Creek (125,000 ybp); Healy-McLeod Creek (65,000 to 75,000 ybp); and Riley Creek-Wonder Lake (9,000 to 25,000 ybp) glaciations. They are followed by the Carlo Creek readvance of 8,000 ybp (Gilbert 1979; Thorson 1986; Brease 2004). As the Alaska Range was uplifted, subsequent glaciations that did not extend out as far as the previous ones left a record of nested moraines

(Gilbert 1979; P. Haeussler, USGS, geologist, written communication, November 2009). Similarly, the various types of rock blocks in glacial deposits can record the distance and direction of the glacial transport from the stone’s provenance to present position.

Lateral moraines are the piles of loose fragments and blocks that fall from overhanging cliffs. These blocks accumulate as low ridges of glacial till riding along the edge of the glacier. If two glaciers intersect, adjacent lateral moraines join to form medial moraines (Brease 2004).

Glacier ice supporting large boulders and blocks of rock can be stranded or rafted on glacial lakes. When the ice melts, the largest blocks are known as glacial erratics. A remnant boulder, or erratic, left from the Browne Glaciation sits on a hillside at elevation 1,040 m (3,400 ft) south of park headquarters. A similar granitic erratic, pulled from the crest of the Alaska Range was stranded at 1,280 m (4,200 ft) on the summit of Mt. Fellows, east of park headquarters (Gilbert 1979).

Glacial deposits are cause for resource management concern because of the large proportions of weak rock flour and clays that form the matrix for larger, irregular blocks of rock. In many glacial deposits, the loose material continually slumps, sometimes sliding over a road or trail surface. Knowledge of the location of such deposits is critical for maintenance of park infrastructure and facilities.

Bergschrunds

Usually a large crevasse, the bergschrund develops in the ice at the head of a glacier as a result of the glacier moving away from the headwall (fig. 19). The size of the bergschrund of many active glaciers in Denali depends on the mass and area of glacier ice. Bergschrunds consist of an arcuate opening between the head of the glacier and the mountain (cirque) wall. The trend of the bergschrund usually parallels the wall. It is at this site that plucking is most active and dominant because water enters the cracks by day and freezes in the rock crevices at night.

Glaciers

Glaciers are major features on the landscape of Denali National Park and Preserve, covering 17% (~1 million acres) of the total park area. More than 400 glaciers exist at the park, the largest and/or most visible 40 of which are named. The Kahiltna Glacier is 71 km (44 mi) long, making it the longest glacier in the Alaska Range. The Ruth Glacier is the thickest glacier in the park, filling the very deep Ruth Gorge. The Eldridge, Tokositna, Yentna, Peters, Harper, and Muldrow Glaciers are among the other named glaciers at Denali.

Permanent snowpack and abundant precipitation above 2,100 m (7,000 ft) supports the flow of glaciers from elevations above 6,100 m (20,000 ft) down glacial valleys to ~305 m (1,000 ft). The most extensive glaciers and snowfields are concentrated on the southeastern side of the Alaska Range, due to the abundant snowfall from

moisture-bearing winds from the Gulf of Alaska. On the drier north side, glaciers are typically smaller and shorter, with the exception of Muldrow Glacier (Brease 2004). Ice within these glaciers moves at varying speeds depending on several factors: glacier thickness, underlying topography, bedrock type and structure, and the presence of water beneath the ice. The average rate in the smallest glaciers is 1.8 to 2.4 m (6 to 8 ft) a year, while the average in the largest glaciers is greater than 70 m (230 ft) a year. Glaciers are never motionless; however, their movement is somewhat slower in winter than in summer. Despite slow speeds, over a period of years, glacial ice transports immense quantities of rock material and debris ultimately to the ends of the glaciers.

Short-lived Glacial Surface Features

Countless interesting, short-lived surface features can be seen at various times on any glacier. These include crevasses, moulins (glacier wells), debris cones, and glacier tables (fig. 19). Crevasses are cracks that occur in the ice of all glaciers due to tensions caused by differences in ice velocity throughout the body of the glacier (fig. 20). They can be hidden by skiffs of snow or debris, and thus can pose a hazard problem to visitors.

Debris cones result from the insulating effect of rock debris, usually deposited by a stream running over the glacier's surface, which protects the ice underneath from the sun's rays. As the surface of the glacier is lowered by melting, cones or mounds form beneath the rock-insulated area and grow gradually higher until the debris slides from them. They are seldom higher than 3 to 4 feet, but can pose a hazard to hikers on the glacier. A glacier table is a mound of ice that is capped and insulated by a large boulder. Its evolution is similar to that of the debris cone or mound (Dyson 1966).

Snow that fills crevasses and wells during the winter often melts out from below, leaving thin snowbridges over the cracks in early summer. These snowbridges pose a very real danger to those traveling on glaciers because of their inherent weakness and instability.

Glacial Retreat and Surging Glaciers

As climate warms (Karl et al. 2009; IPCC 2007), glaciers are shrinking at Denali National Park and Preserve (figs. 5, 6, 18). Most of the glaciers are consistently thinning (see "Geologic Issues" section). Many glaciers appear underfit for their U-shaped valleys (Brease 2004). When the yearly snow accumulation decreases, the ice front of the glaciers seems to retreat, whereas the mass of the

glaciers is merely decreasing by melting on top and along the edges (analogous to an ice cube melting on a kitchen counter).

Surging Glaciers

When a glacier advances faster than average in one or more seasons, it is said to be surging. The Muldrow Glacier, approaching the park visitor center from the west, has advanced rapidly several times (Brease 2004). The last major surge occurred during the 1956-1957 season. Glacial surging can be the result of periodic water buildup beneath the glacier, lubricating the base of the glacier and facilitating rapid and potentially catastrophic movements of ice (Gilbert 1979). A major tributary to the Muldrow Glacier is the Traleika Glacier. This glacier has flowed at rates of 20 to 70 m (66 to 230 ft) per year since 1991 (Adema et al. 2003).

Presently, only about 10% of glaciers are surging at any one time (fig. 4) (P. Haeussler, USGS, geologist, written communication, November 2009). Most of surging glaciers are located on glaciers of northern exposure along the Alaska Range.

Other Notable Geologic Features

- Polychrome overlook's multi-colored rocks are the result of weathered rhyolite tuffs and interlayered flows of the Teklanika Formation, deposited 56 million years ago. Similar, colorful volcanics are visible on Cathedral, Double, and Igloo Mountains (Cole 2004). This volcanism was contemporaneous with the formation of Mt. McKinley, suggesting that the volcanic vent source was to the southwest.
- Evidence of more recent volcanism, ~38 million years ago, is conspicuous from the Eielson Visitor Center, and forms much of Mt. Galen with basalt, andesite, dacite, and rhyolite lava flows, tuffs, and breccias (Cole 2004).
- Near Teklanika Campground, the quartz-rich conglomerate of the Cantwell Formation rests atop much older metamorphic rocks, marking a 100- to 65-million year gap (unconformity) in the geologic record. Elsewhere, in the Tolkat River gorge, the Cantwell sits atop pillow basalts that are 135 million years older (Gilbert 1979).
- "Badlands" topography is developed along the gorge of the East Fork River within rhyolite pebble conglomerates of the Nenana Gravel.



Figure 14: Denali (Mt. McKinley) dominates the ridgeline of the Alaska Range in Denali National Park and Preserve. Wonder Lake (note canoe for scale) is in the foreground. NPS photo.



Figure 15: Denali has a rich variety of deformed bedrock, a tribute to the complex geologic history which includes active plate dynamics. NPS Photo/Adema.




Figure 16: Map of Alaska showing the Denali fault, Iditarod-Nixon fault (INF), Tintina-Kaltag fault, the Border Ranges fault (BRF - Talkeetna fault), and other regional structures in relation to the southern coast of Alaska and the Aleutian trench subduction zone. Note the location of the Yakutat Terrane. An orocline is an orogenic belt with an imposed curvature or sharp bend. Graphic by Phil Reiker (NPS Geologic Resources Division) after fig 1. from Miller et al. (2002).




Figure 17: Cross section of subduction zone between the Talkeetna Superterrane and the Yukon-Tanana Terrane of southern Alaska during early Cretaceous accretion events. Note the location of miniterranes and ocean basins between the two large terranes. Graphic by Trista L. Thornberry-Ehrlich (Colorado State University), adapted from Csejtey et al. (1982) fig. 9.




Figure 18: Diagrammatic view of a retreating glacier with a recent terminal moraine and a broad space between the moraine and the glacier front. Note the variety of glacial deposits. Graphic by Trista L. Thornberry-Ehrlich (Colorado State University).




Figure 19: Short-lived features occurring on the surface of glaciers. Graphic by Trista L. Thornberry-Ehrlich (Colorado State University).



Figure 20: Ice and crevasses of Eldridge Glacier in Denali National Park and Preserve, Alaska, August 7, 2004. Photo courtesy of Ronald D. Karpilo, Jr. (Colorado State University).

Map Unit Properties

This section identifies characteristics of map units that appear on the Geologic Resources Inventory digital geologic map of Denali National Park and Preserve. The accompanying table is highly generalized and for background purposes only. Ground-disturbing activities should not be permitted or denied on the basis of information in this table.

Geologic maps facilitate an understanding of Earth, its processes, and the geologic history responsible for its formation. Hence, the geologic map for Denali National Park and Preserve provided information for the “Geologic Issues,” “Geologic Features and Processes,” and “Geologic History” sections of this report. Geologic maps are two-dimensional representations of complex three-dimensional relationships; their color coding illustrates the distribution of rocks and unconsolidated deposits. Bold lines that cross or separate the color patterns mark structures such as faults and folds. Point symbols indicate features such as dipping strata, sample localities, mines, wells, and cave openings.

Incorporation of geologic data into a Geographic Information System (GIS) increases the usefulness of geologic maps by revealing the spatial relationships among geologic features, other natural resources, and anthropogenic features. Geologic maps are indicators of water resources because they show which rock units are potential aquifers and are useful for finding seeps and springs. Geologic maps are not soil maps, and do not show soil types, but they do show parent material—a key factor in soil formation. Furthermore, resource managers have used geologic maps to make connections between geology and biology; for instance, geologic maps have served as tools for locating sensitive, threatened, and endangered plant species, which may prefer a particular rock unit.

Although geologic maps do not show where earthquakes will occur, the presence of a fault indicates past movement and possible future seismic activity. Similarly, map units show areas that have been susceptible to hazards such as landslides, rockfalls, and volcanic eruptions. Geologic maps do not show archaeological or cultural resources, but past peoples may have inhabited or been influenced by depicted geomorphic features. For example, alluvial terraces may have been preferred use areas and formerly inhabited alcoves may occur at the contact between two rock units.

The geologic units listed in the following table correspond to the accompanying digital geologic data. Map units are listed in the table from youngest to oldest. Please refer to the geologic timescale (fig. 21) for the age associated with each time period. The table highlights characteristics of map units such as: susceptibility to erosion and hazards; the occurrence of paleontological resources (fossils), cultural resources, mineral resources, and caves or karst; and suitability as habitat or for recreational use. Some information on the table is

conjectural and meant to serve as suggestions for further investigation.

The GRI digital geologic maps reproduce essential elements of the source maps including the unit descriptions, legend, map notes, graphics, and report. The following references are source data for the GRI digital geologic data for Denali National Park and Preserve:

Clautice, K.H., Newberry, R.J., Blodgett, R.B., Bundtzen, T.K., Gage, B.G., Harris, E.E., Liss, S.A. Miller, M.L., Reifenhuth, R.R., and D.S. Pinney. 2001. *Bedrock Geologic Map of the Chulitna Region, Southcentral Alaska*. Scale 1:63,360. Report of Investigations RI2001-1A. Fairbanks, Alaska: State of Alaska Department of Natural Resources, Division of Geological and Geophysical Surveys.

Pinney, D.S. 2001. *Surficial-Geologic Map of the Chulitna Mining District, Southcentral Alaska*. Scale 1:63,360. Report of Investigations RI2001-1C. Fairbanks, Alaska: State of Alaska Department of Natural Resources, Division of Geological and Geophysical Surveys.

Pinney, D.S. 2001. *Engineering-Geologic Map of the Chulitna Mining District, Southcentral Alaska*. Scale 1:63,360. Report of Investigations RI2001-1D. Fairbanks, Alaska: State of Alaska Department of Natural Resources, Division of Geological and Geophysical Surveys.

Wahrhaftig, Clyde. 1970. *Geologic Map of the Healy D-4 Quadrangle, Alaska*. Scale 1:63,360. Geological Quadrangle Map GQ-806. Reston, VA: U.S. Geological Survey.

Wahrhaftig, Clyde. 1970. *Geologic Map of the Healy D-5 Quadrangle, Alaska*. Scale 1:63,360. Geological Quadrangle Map GQ-807. Reston, VA: U.S. Geological Survey.

Wahrhaftig, Clyde. 1952. *Geologic Map of Part of Healy C-4 Quadrangle, Alaska, Showing Pleistocene Deposits along the Nenana River*. Scale 1:63,360. Professional Paper 2933. Reston, VA: U.S. Geological Survey.

Wahrhaftig, Clyde, and R.E. Fellows. 1952. *Geologic Map of Part of Healy B-4 Quadrangle, Alaska, Showing Quaternary Deposits along the Nenana River*. Scale 1:63,360. Professional Paper 2934. Reston, VA: U.S. Geological Survey.

Wilson, F.H., Dover, J.H., Bradley, D.C., Weber, F.R., Bundtzen, T.K., and P.J. Haeussler, compilers. 1998. *Geologic Map of Central (Interior), Alaska*. Open-File Report OF 98-133. Reston, VA: U.S. Geological Survey.

The Map Unit Properties Table is divided into units that are present on larger scale maps (1:63,360; Chulitna region and Healy quadrangle maps) and those on the smaller scale map (1:250,000; Wilson et al. 1998 compilation). The overview of digital geologic data (Appendix A) illustrates the small scale digital geologic data.

The GRI team implements a geology-GIS data model that standardizes map deliverables. This data model dictates GIS data structure including data layer architecture, feature attribution, and data relationships within ESRI ArcGIS software, and increases the overall utility of the data. GRI digital geologic map products include data in ESRI personal geodatabase and shapefile GIS formats, layer files with feature symbology, Federal Geographic Data Committee (FGDC)-compliant metadata, a Windows help file that contains all of the ancillary map information and graphics, and an ESRI ArcMap map document file that easily displays the map and connects the help file directly to the map document. GRI digital geologic data are included on the attached CD and are available through the NPS Data Store (<http://science.nature.nps.gov/nrdata/>). Data will be available on the Natural Resource Information Portal when the portal goes online. As of August 2010, access is limited to NPS computers at <http://nrinfo/Home.mvc>.

Geologic Units of Denali National Park and Preserve

The rock units at Denali National Park and Preserve represent a vast span of geologic history. Because juxtaposed slices of accreted terranes characterize the geology in the Denali area, different rock types and ages form a puzzle of units on the landscape. The oldest rocks are metamorphosed Proterozoic basement rocks of the Nixon Fork sequence. These rocks contain zircon grains dated using uranium-lead isotopes (U-Pb) at $1,250 \pm 50$ million years ago (Wilson et al. 1998).

Most of the oldest rocks are part of the Yukon-Tanana Terrane, north of the Hines Creek fault (Gilbert 1979). These Late Precambrian to Early Paleozoic age rocks include metamorphosed pelitic and quartzose schists, amphibolite, gritty quartzite, marble, greenstone, phyllitic layers, and other metasedimentary and metaigneous rocks (Gilbert 1979; Wilson et al. 1998). Ordovician rocks include dark chert, slate, argillite, and some greenstone, limestone, and dolostone. Devonian limestone, chert, and volcanic ash fall and lava flows record prolific life and volcanic activity in a marine basin-island arc setting. The Mississippian Moose Creek Member contains metamorphosed basalts and other volcanic rocks (Clautice et al. 2001).

Pennsylvanian and Permian limestones, mudstones, and graywacke sandstones record marine deposition in open basins. During the Mesozoic, granitic magma formed irregular intrusive bodies throughout the region, and mixed schists and amphibolite of the MacLaren Metamorphic Belt underwent several grades of metamorphism and folding (Wilson et al. 1998; Clautice et al. 2001). Triassic greenstones, basalts flows, graywackes, tuffs, and limestones possibly record hotspot activity in a marine basin. These rocks record paleolatitudes of 5 degrees near the equator (Clautice et al. 2001).

Jurassic and Cretaceous age rocks in the map area range from calcareous sandstone and argillite to conglomerates and limestones to vast suites of volcanic rocks of varying compositions, granite-diorite-rhyolite intrusions, mixed mélanges, and mafic igneous rocks (Wilson et al. 1998; Clautice et al. 2001). The vast variety of rocks of this age attests to the tectonic history of the area. Many terranes were accreted with accompanying metamorphism, sedimentation, deformation, and volcanism during this time.

Tectonic processes continued throughout the Tertiary and Quaternary Periods. Tertiary age units include rhyolite, andesite, granite, granodiorite, tonalite, and monzonite intrusive igneous rocks. These units are mixed with fluvial sedimentary rocks, carbonaceous mudstone, siltstone, conglomeratic sandstone, olivine basalts, volcanics, and coal beds. Named Tertiary units include the Suntrana, Sanctuary, Healy Creek, Lignite Creek, Grubstake, and Cantwell Formations (Wahrhaftig 1970; Wilson et al. 1998).

The intrusive igneous bodies and plutons of hornblende dacite of Jumbo Dome have a relatively young potassium-argon (K-Ar) radiometric date of 2.72 ± 0.25 million years ago (Wahrhaftig 1970). Atop this and other map units at Denali National Park and Preserve are vast and varied unconsolidated Quaternary age rocks of fluvial, colluvial, glacial, lacustrine, and aeolian origin. Older deposits include: rubble from Jumbo Dome; gravels of Late Wisconsin glaciations; high terraces and pediment gravels; rock glaciers; older alluvium (sands, gravels, and silts); glacial erratics up to 2 m (6 ft) in diameter; some varved clays deposited in glacial lakes; and terminal glacial moraine deposits (Wahrhaftig 1970; Clautice et al. 2001).

Younger units include Holocene tills, alluvium, outwash, the end moraine of the Carlo Readvance, extensive terraces above current stream valleys, windblown loess, talus and landslide deposits, and other colluvium. The youngest units at Denali include: boggy peat; stream alluvium; alluvial fans; outwash gravel and drift; some localized debris flows; varved clay and silt in recently dried-up lakes and ponds; swamp deposits; and flood-plain deposits of sand, gravel, silt, mud, and scattered boulders (Wahrhaftig 1970; Clautice et al. 2001).

Map Unit Properties Table: Denali National Park and Preserve (Large Scale [1:63,360] Map Units)

Age	Unit Name (Symbol)	Features and Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
QUATERNARY	Peat (Qp) Artificial Fill and Excavation Sites (Qh) Undifferentiated Stream Alluvium (Qa) Stream Gravel (Qsg) Alluvial Fan Deposits (Qaf) Outwash Gravel (Qo) Floodplain Alluvium (Qfp)	Qp contains dense, dark organic material, often found in boggy areas. Qh is composed of pebbles, cobbles, gravels, sand, and silt beneath roads and former gravel pits. Unit is well- to poorly sorted. Qa includes stream gravel, pebbles, cobbles, sand, and silt, with some boulders in well-stratified and moderately sorted channels, floodplains, and low terraces. Stream gravel (Qsg) is present in active streambeds. Qaf contains fan-shaped, heterogeneous strata of gravels, sands, silts, and boulders. Some debris flow deposits present locally. Qo contains gravel in layers fluvially deposited and well sorted. Qfp consists of elongate deposits of fluvial sand, gravel, and silt with scattered boulders. Unit is well- to moderately sorted, well-stratified, often mantled by thin silt-clay layers. Some terrace deposits locally.	Low to very low.	Hummocky topography and scarps make Qfp unstable if water saturated.	Units are prone to rapid erosion during flood events, slope failure if undercut, and slope creep, especially when water saturated.	Modern remains possible; pollen, plant macrofossils, insects.	May contain campsites, settlements, and other artifacts. Historic mining?	Gravel, sand, silt, and clay. Placer deposits? Congeliturbate deposited along Alaska Railroad between Lagoon and Carlo.	None.	Qa supports extensive willow-alder thickets and other riparian habitats. Qfp supports bog environments and riparian zones.	Units are suitable for most recreation. Avoid development of use-areas on riparian zones and undercut floodplain areas.	Units record most recent geologic activity in the area.
	Undifferentiated Colluvium (Qc) Landslide Deposits (Qcl) Talus and Rubble Deposits (Qct) Rock Glacier Deposits (Qrg)	Units contain irregular masses, aprons, tongues, and fans of angular rock fragments, sand, and gravel. May include drift and outwash. Surface expression often mirrors underlying rock structure. Medium to thickly bedded layers of lobed and terraced deposits. Rubble includes blocks up to 15 m in diameter. Qct contains mostly large blocks, with little- to no matrix material such as silt and sand. Qrg includes mixtures of blocks, ice, gravel, sand, and silt in tongue- and fan-shaped masses with very irregular surfaces.	Low.	Units are often found at the base of unstable slopes. Surfaces are usually steep and irregular with large spaces between blocks; unsuitable for most forms of infrastructure.	Units are associated with mass movement processes, including sliding, rolling frost creep, gelifluction, and flowing. Open work rubble surfaces are dangerous. Extensive ground cracks present locally.	Modern remains possible; pollen, plant macrofossils, insects.	None documented.	Sand, gravel, and boulders.	None.	Lichen communities help determine relative ages of deposits. Units may support large trees if old enough.	Avoid for recreational use; dangerous surfaces associated with units.	Units record recent slope processes in the area; vegetation patterns can date slide activity.
	Swamp Deposits (Qs) Outwash Alluvium (Qao) Colluvial-Alluvial Valley Fill, Fan, and Apron Deposits (Qcf) Terrace Gravel (Qtg) Abandoned Channel Deposits (Qac) Terrace Alluvium (Qat)	Qs includes water/ice-saturated layered peat, and organic silt and sand of variable thickness. Qao and Qcf contain elongate, apron, tongue- and fan-shaped heterogeneous mixtures of boulder, pebbles, and cobbles with matrices composed of sand and silt in thick beds. Terrace and channel gravels (Qtg, Qac, Qat) are typically more rounded than angular cobbles, and are well sorted with local crossbeds and smooth surfaces.	Low.	Slopes and scarps associated with Qao and Qtg may be too unstable for heavy infrastructure. High permeability of these units are probably not suitable for waste-water treatment facilities.	Qcf is associated with debris flow mechanisms, and may include avalanche deposits.	Peat, plant fragments.	May contain campsites, settlements and other artifacts. Historic mining?	Sand, silt, and gravel. Placer deposits?	None.	Units contain bogs, wetlands, and riparian environments.	Fine for most light recreation. Avoid development of use-areas on riparian zones and undercut floodplain areas.	Units record Quaternary fluvial activity in the area.

Map Unit Properties Table: Denali National Park and Preserve (Large Scale [1:63,360] Map Units)

Age	Unit Name (Symbol)	Features and Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
QUATERNARY	Till of Latest Holocene Age (Qt2) Till of Early to Middle Holocene Age (Qt1) Alluvium (Qca) Outwash Gravel (Qco) Sand (Qcls) End Moraine of the Carlo Readvance (Qcm)	Units contain heterogeneous mixtures of pebbles, cobbles, sand, silt, and clay with scant boulders. Some ice may exist locally. Surfaces of till units are often hummocky with morainal topography. Qcls is deposited in a dammed lake behind the end of the Carlo Readvance.	Low to very low for sand-rich units	Heterogeneous nature of these unconsolidated units renders them unstable on slopes and too permeable for waste treatment.	Units are associated with slumping and mass movements; hummocky topography is difficult to traverse.	Modern remains possible; pollen, plant macrofossils, insects.	May contain campsites, settlements, and other artifacts.	Sand, silt, and gravel. Placer deposits?	None.	Qt1 supports some soil development with lichens, mosses, and tundra herbs.	Units are fine for most light recreation. Avoid areas of very large rubble.	Unit records latest glacial activity in the area.
	Alluvium and Pediment Gravel (Qra) Younger Alluvium and Pediment Gravel (Qra2) Older Alluvium and Pediment Gravel (Qra1) Varved Clay (Qrlc) Outwash Gravel (Qro) Rubble of Jumbo Dome (Qrr) Glacial Moraine Deposits (Qrm)	Units contain mixed pebbles, gravels, and sands in smooth surfaced lobes and pediments. Qrlc was deposited in glacial lakes prior to the Carlo Readvance. Qrr is composed of inactive (non-flowing) rock glaciers and solifluction sheets cored with ice. Qrm contains till with extremely heterogeneous mixtures of boulders, pebbles, and clay with some outwash deposits interlayered.	Low.	Heterogeneous nature of these unconsolidated units renders them unstable, especially if water-saturated and/or present on undercut slopes.	Units are susceptible to frost creep, erosion, and mass wasting. Open work rubble surfaces are dangerous. Extensive ground cracks present locally.	Modern remains possible; pollen, plant macrofossils, insects.	May contain campsites, settlements, and other artifacts.	Gravel and sand; igneous stone rubble.	None.	Unit supports valley floor vegetation, some riparian zones.	Units are fine for most light recreation. Avoid areas of very large rubble.	Varved clays record seasonal cycles of deposition associated with glacial lakes.
	Varved Clay and Silt (Qhlc) Alluvium and Pediment Gravel (Qha) Outwash Gravel (Qho) Rubble of Jumbo Dome (Qhr) Morainal Deposits (Qhm) Glacial Erratics (Qhe) Lacustrine Silt and Clay (Qhl)	Units contain layered clays, gravels, pebbles, and sands, as well as large boulders floated on ice rafts across glacial lakes (including Glacial Lake Moody). Morainal deposits include cobbles and boulders with a gravel-sand-silt matrix.	Low to very low for clay-rich units.	Heterogeneous nature of these unconsolidated units renders them unstable, especially if water-saturated and/or present on undercut slopes.	Boulders and other large rocks may weather out of finer-grained matrix and pose rockfall hazard; units are susceptible to erosion and gullyng.	Plant fragments.	None documented.	Clay, gravel, sand, and boulders.	None.	None documented.	Avoid areas of very large rubble for recreational opportunity development.	Units record glacial activity. Varved clays record seasonal cycles of deposition associated with glacial lakes.

Map Unit Properties Table: Denali National Park and Preserve (Large Scale [1:63,360] Map Units)

Age	Unit Name (Symbol)	Features and Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
QUATERNARY	Delta Gravel (Qhd) Drift of Late Wisconsin Age (Qd) Outwash Gravel (Qdo) Rubble of Jumbo Dome (Qdr) Morainal Deposits (Qdm) Lacustrine Silt and Clay (Qdl)	Qhd is gravel contained in tongue- to lobe-shaped layered deposits. Qd is composed of blanket deposits of heterogeneous pebble, gravel, cobble, sand, silt, and clay interlayers. Some large (2-m) boulders present locally. Sorting and bedding depends on degree of reworking by fluvial processes. Eskers present locally. Loess silt often blankets these units up to 1 m thick. Other units (Qdo, Qdr, Qdm, Qdl) range from scattered rubble, to layered heterogeneous lobes, to well-bedded silts and clays from lake deposits.	Low.	Eskers and diamictons may form steep slopes that would be unstable for infrastructure.	Scarps and slopes associated with these units are prone to mass movement and slope failures.	Modern remains possible; pollen, plant macrofossils, insects.	May contain campsites, settlements, and other artifacts.	Gravel, sand, and silt.	None.	Units support vegetation that prefers strong drainage.	Avoid areas of very large rubble for recreational opportunity development.	Clay-rich diamictons and eskers showcase glacial landforms.
	Glacial Erratics (Qde) Alluvium and Pediment Gravel (Qba) Outwash Gravel (Qbo) Glacial Erratics (Qbe) Colluvium (Qpc) Terrace and Pediment Gravel (Qtp) Dune Sand (Qsd) Rubble, Possibly Rock Glacier Origin (Qrb)	Units contain large, scant boulders dropped from drifting glacial ice, as well as angular slope deposits in lobe- to fan shapes. Gravels and sands are better sorted than heterogeneous rubble and boulders, possibly associated with an ice-cored rock glacier.	Low.	Colluvial units are often found at the base of unstable slopes. Surfaces are usually steep and irregular with large spaces between blocks.	Units are susceptible to frost creep, erosion, and mass wasting. Open work rubble surfaces are dangerous.	Modern remains possible; pollen, plant macrofossils, insects.	None documented.	Gravel, sand, and boulders.	None.	None documented.	Avoid areas of very large rubble for recreational opportunity development.	Glacial erratics are an interesting interpretive topic, and contain clues to glacial provenance.
QUATERNARY - TERTIARY	Hornblende Dacite of Jumbo Dome (QThj)	Unit contains hornblende-rich basaltic igneous intrusives.	High.	Suitable for most forms of infrastructure unless highly fractured and/or altered.	Unit associated with sheet-like exfoliation, which may lead to rockfall hazards.	None.	None documented.	Hornblende phenocrysts.	None.	Unit may alter to form calcium-, aluminum-, and magnesium-rich soils.	Unit is fine for most forms of recreation; may attract climbers.	Isotopically-determined date on this unit of 2.72 ± 0.25 million years.
TERTIARY AND OLDER	Thinly Covered Bedrock (Ttcb)	Undifferentiated bedrock of various ages overlain by thin layers of unspecified surficial deposits.	Variable, depending on properties of underlying bedrock and surficial deposits.			Unknown. Resources from bedrock units may be present.				Variable, depending on properties of underlying bedrock and surficial deposits.		Indicates where bedrock is located near the surface.

Map Unit Properties Table: Denali National Park and Preserve (Large Scale [1:63,360] Map Units)

Age	Unit Name (Symbol)	Features and Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
TERTIARY	Nenana Gravel (Tn) Coal Bearing Group, Undivided (Tcu) Lignite Creek Formation (Tlc) Gravel and Sand (Ts) Grubstake and Lignite Creek Formations, Undivided (Tgl)	Tn contains poorly consolidated layers of buff to reddish-brown pebble and boulder conglomerate with coarse sandstone, mudflow, and claystone interbeds. Some lignite present locally. Tlc contains crossbedded arkosic sandstone with pebble conglomeratic lenses and clay interbeds. Unit is mostly buff colored. Ts contains orangish- to buff colored sandy pebble, cobble, and gravel conglomeratic beds. Unit is typically well sorted and massively bedded with horizontal layering and crossbeds. Sands occur as broad lenses with total thickness ranges from 6 to 45 m. Tgl contains greenish-gray siltstone and claystone with some interbedded pebbly arkosic sandstone and silt and clay layers.	Moderately low.	Ts is exposed as cliffs in stream-incised areas, and may prove unstable on slopes. Units are permeable, but may be utilizable for waste water treatment. Avoid poorly consolidated areas for infrastructure.	Ts is prone to gullying and cliff formation, which may pose rockfall and mass movement hazards.	Pollen.	Chert pebbles may have provided tool material.	Lignite, sand, gravel, clay, and subbituminous coal.	None.	Units support wide range of habitats.	Units are fine for most recreation unless highly weathered, rendering them unstable for trail base.	Chert, argillite, graywacke, volcanic rocks, Felsic to intermediate intrusive rocks, and Cantwell conglomeratic clasts indicate source area.
	Suntrana Formation (Tsn) Sanctuary Formation (Tsc) Healy Creek Formation (Thc) Coal Bearing Sandstone (Tcs)	Tsn contains crossbedded pebbly sandstone, poorly consolidated and interbedded with silty claystone and some coal lenses. Appears white on weathered surfaces. Tsc is poorly consolidated and weathers brown- to gray with abundant banded (varved?) shale. Thc contains clay-rich quartz and micaceous sandstone, some quartz-chert conglomerate, claystone, and coal interbeds. Unit is poorly consolidated. Tcs is pebbly sandstone and conglomerate with lignite, silty claystone, and carbonaceous claystone interbeds.	Moderately low to low where consolidation is poor.	Most units are poorly consolidated, which may render them unsuitable for heavy infrastructure.	Tsc is extremely susceptible to landslides and slope creep.	Potential plant fossils associated with coal beds.	Chert pebbles may have provided tool material.	Subbituminous coal lenses, conglomerate, sand, gravel, clay, and silt. Lignite 3 to 6 m thick are correlative with nearby Dunkle coal mine deposits.	Not enough carbonate present.	Units support wide range of habitats.	Poor consolidation of units may render them too unstable for heavy recreational use.	Contains Oligocene coal units.
	Rhyolite (Tr) Latite (Tl) Andesite Plugs, Hypabyssal Intrusions, Plug-Domes and Associated Flows and Breccia (Ti) Vent Breccia (Tvb) Biotite Granite (Tg) Basaltic Composition Dikes (Tb) Diabase Sills (Tdb)	Tr contains assorted felsic volcanic extrusive layers in widespread deposits. Units contain intermediate composition (Si content) igneous intrusive, as well as basaltic composition flows and intrusive dikes. Volcanic breccia indicating forceful eruptions present locally. Tg is fine-grained in dikes to coarser-grained irregular intrusions with alkali feldspar, biotite, ilmenite, and quartz. Unit is commonly intruded by basaltic or lamprophyric dikes. Tb are narrow (0.3- to 2-m) basaltic dikes cutting older granites with abundant phenocrysts. Tdb are intrusive into the Cantwell Formation (Tc)	Moderately high to high.	Units may contain radon emitting materials. Avoid for basements and foundations.	Blockfall and exfoliation of large sheets of rock associated with several units.	None.	Phenocrysts may have provided trade material.	Geodes; Tb contains clinopyroxene phenocrysts. Tourmaline, zinnwaldite, topaz, zircon, apatite, and fluorite.	None.	Unit may weather to enriched soils.	Phenocrysts and other sharp fragments associated with unweathered portions of these units may prove undesirable for trail base.	Tg is dated by argon isotopes at 46 to 55 million years (composition indicates felsic magma source). Tb is dated by argon isotopes at 52 million years, with magnetic signature indicating early Tertiary paleolatitude?
	Cantwell Formation (Tc) Cantwell Formation, Tuff Bed (Tct)	Unit contains buff- to light gray sandstone and pebble conglomerate with some darker layers, siltstone, and shale interbedded. Several volcanic tuff layers are present locally (Tct). Unit is moderately- to well consolidated and provides bedrock source material for colluvial deposits.	Moderate.	Units are fine for most infrastructure unless highly heterogeneous and/or fractured	Unit is associated with slope processes, including landslides.	None documented in large scale source maps. See Kcs on small scale data table, below.	Abundant chert pebbles may have provided tool material.	Building stone.	Not enough carbonate present.	Units support slope preferring vegetation.	Units are fine for most recreation.	Source rocks well documented from volcanic rocks to the south.

Map Unit Properties Table: Denali National Park and Preserve (Large Scale [1:63,360] Map Units)

Age	Unit Name (Symbol)	Features and Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
MESOZOIC	Granite (MZgr)	Undifferentiated granite found in irregular bodies with no clear crosscutting relationships to place in an age bracket. Granite contains quartz, feldspar, biotite, and hornblende with various accessory minerals.	High.	Unit may alter into materials known to cause radon problems.	Sheet-like exfoliation of unit may pose blockfall hazard.	None.	Phenocrysts may have been used for trade materials.	Accessory minerals and phenocrysts. Building material.	None.	None documented.	Fine for most recreation unless highly fractured and/or altered.	Unit is widespread.
CRETACEOUS	Hornblende Biotite Granite (Kg) Intermediate Composition Plutonic Rocks (Km) Rhyolite Intrusive into the Cantwell Formation (Kr) Diabase Intrusive into the Cantwell Formation and Basalt Intrusive into the Birch Creek Schist (Kdb) Andesite Intrusive into the Cantwell Formation (Ka)	Kg is present in narrow dikes and small stocks with granitic porphyry textures of monzogranite to syenogranite compositions. Grain size ranges from less than 2 mm to 2 cm. Unit grades into Km , which contains northeast/southwest-oriented stocks and dikes with textures ranging from fine-grained porphyritic to medium grained. Unit is generally quartz poor with monzodiorite, monzonite, and diorite compositions, and alters to chlorite, carbonate, sericite, albite, epidote, and pyrite. Intrusions into Tc indicate a younger age for the unit than previously thought, additional investigation is needed to resolve. Compositions range from felsic to intermediate to mafic (basaltic), and most are present as narrow dikes.	Moderately high to high depending on degree of weathering.	Units may contain radon producing materials. Avoid for basements; avoid heavily fractured areas.	Units are susceptible to rockfall hazards on slopes.	None.	Pyrite may have been used for starting fires.	Hornblende phenocrysts, zircon, apatite, ilmenite, muscovite, and rutile. Hydrothermal alteration minerals include chlorite, calcite, and white mica.	None.	Units may give rise to fertile soils.	Suitable for most recreation; may be attractive to climbers.	Kg has minimum age of 61.4 to 71 million years (determined by argon isotopes) origin is likely volcanic island arc-related. Km is dated at 67 to 71 million years. Associated with gold-related Kuskokwim Mountains magmatic belt event.
CRETACEOUS - DEVONIAN	Undifferentiated Sedimentary Rocks of Devonian to Cretaceous Age (KDu)	Heterogeneous mix of sedimentary rocks, including sandstone, claystone, shale, limestone, conglomerate, and tuff (volcanic ash fall).	Moderate; variable among units.	Units are fine for most infrastructure unless highly heterogeneous and/or fractured.	Heterogeneous layers of these units may be prone to mass wasting; shrink-and-swell clays may be present in volcanic ash units.	Various fossils, none specified.	None documented.	Building stone.	Possible in limestone-rich units.	Units support wide range of habitats.	Units are fine for most recreation.	Records eras of sedimentary deposition settings.
CRETACEOUS - JURASSIC	Calcareous Sandstone and Argillite, with Coquinoid Limestone (KJs) Argillite and Sandstone (KJas) Sandstone and Argillite (KJsa) Conglomerate (KJc)	Units are rich in gray, poorly sorted sub-angular to sub-rounded lithic sandstones mixed with argillite layers and coquinoids locally. Some of the sandstone is calcareous (e.g., KJs). In outcrop, units weather to an orangish-brown color. Bedding is predominantly thin, parallel, and sheet-like. Some isoclinal folding is present, leaving cleavage in the argillaceous lithologies. KJas contains dark gray and black argillite and sandstone. Unit weathers orangish-brown with prominent cleavage parallel to bedding. Some white quartz veining is present locally. Argillite accounts for a smaller percentage in the lower units (e.g., KJsa), with massive sandstone beds becoming more prominent. Lowermost unit (KJc) is conglomerate with sandstone matrix. Clasts are mostly quartz, chert, and argillite.	Moderate to moderately high, depending on degree of weathering.	Prominent cleavage, thin bedding, heterogeneous composition, and minor dissolution in these units may render them weak on slopes for heavy infrastructure.	Slaty layers are prone to slides, and conglomeratic blocks may pose rockfall hazard if underlain by weaker claystone units.	Coquinoid beds with <i>Buchia sublaevis</i> shells in 2-m-thick layer (Lower Cretaceous age).	Chert nodules may have provided tool material; pyrite may have been used to start fires.	KJas contains white quartz veins, sulfides, crystal filled vugs, and fractures; KJsa contains rare tripolitic chert and white mica. Building stones.	Minor carbonate dissolution is possible in coquinoid layers.	Units support wide range of habitats.	Units are fine for most recreation.	Unit records orogenic provenance of sediments in a shallow (<125 m) shelf environment.

Map Unit Properties Table: Denali National Park and Preserve (Large Scale [1:63,360] Map Units)

Age	Unit Name (Symbol)	Features and Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
JURASSIC	Argillite, Cherty Argillite, and Minor Cherty Tuff and Basaltic Tuff (Jac) Calcareous Sandstone, Sandy Limestone, and Argillite (Js)	Jac is mostly deformed and slightly metamorphosed argillite with some minor chert-rich layers and tuff layers. Chert is light gray to black. Sheared areas are phyllitic. Js is thin to medium bedded, calcareous sandstone, with minor interbeds of limestone and argillite.	Moderate.	Suitable for most infrastructure unless highly altered and/or fractured. Avoid phosphatic areas.	Altered tuff layers may contain shrink-and-swell clays.	Radiolaria, ammonites, other Jurassic fossils.	Chert nodules may have provided tool material.	Phyllite. Phosphates in Js.	Not enough carbonate present.	Unit may produce soils rich in phosphorus.	Unit is fine for most light recreation. Avoid finely bedded areas for stability.	Radiolarians restrict Jac to Late Jurassic age. Ammonites in Js are Early Jurassic age (Sinemurian).
TRIASSIC	Redbed Sandstone and Conglomerate (TRrb) Brown Sandstone and Argillite (TRs) Limestone (TRul)	TRrb contains calcareous sandstone, siltstone, and conglomerate layers that weather to reddish-maroon colors. Hematite lends color to this unit; clasts are mostly well-rounded quartz, basalt, and volcanics. TRs contains thick-bedded yellowish-brown sandstone, argillite, and minor calcareous sandstone and limestone. TRul contains thick-bedded limestone, with shale, siltstone, and mudstone interbeds. Lithologies include 300 m of mudstone, wackestone, packstone, and rudstone.	Moderate to moderately high.	Units may contain shrink-and-swell clays; heterogeneous layering may be unstable on slopes. Fine for most infrastructure unless highly fractured.	If massive layers are underlain by weaker layers locally, rockfall hazard may result.	TRs contains Heterastridium (4-cm diameter), snails, and bivalves. TRul contains scleractinian thicket reefs, brachiopods, corals, bivalves, and gastropods.	Caves possible in this unit; may have been settlement areas.	Building stones.	Karst features such as dissolution holes and caves are possible for lower unit.	Units weather to calcium- and iron-rich soils locally.	Fine for most recreation; if caves are present, may attract speleologists.	Fossils record Late Triassic age; near-shore environment grading offshore in lower units.
	Basalt, Basaltic Tuff, and Limestone (TRlb) Basaltic Dikes and Sills (TRb) Intrusive Greenstone (TRgn) Red Colored Tuff, Andesite, Basalt, Graywacke, Conglomerate (TRvs) Limestone (TRll)	TRlb is a massive stack of basalt and basaltic tuff layers, 50 to 100 m thick, with fine-grained gray limestone interbeds. Basalt is tholeiitic and strongly magnetic. Alteration minerals include chlorite, albite, carbonate, and hematite. Unit is commonly intruded by fine-grained equigranular to porphyritic basaltic dikes and sills (TRb). TRgn is an intrusive greenstone unit. TRvs contains a red-weathered stack of volcanic and volcanics, including lithic tuff, conglomerate, graywacke, tuffaceous siltstone, and mudstone with some basalt and dacite flows. Calcareous cements and clasts are present. Lithic fragments include quartz, felsic volcanics, and plutonic and metamorphic rocks. Unit was exposed to contact metamorphism near Cretaceous plutons. TRll is thin-bedded, light gray to brown on weathered surfaces composed of carbonaceous mudstone and packstone present in fault-bounded slivers.	Moderate.	Calcareous layers and cements may dissolve, rendering unit unstable in some areas; units may contain shrink-and-swell clays. Avoid for heavy infrastructure.	If carbonate layers and cements are dissolved, rockfall hazards exist; heavily altered layers may fail when water saturated on slopes.	Limestone layers include colonial scleractinian corals, megalodontid bivalves, and brachiopod Spondylospira lewesensis; TRll contains at least 13 species of ammonites.	None documented.	Chlorite, albite, clinopyroxene, and hornblende. Tin-type mineralization; gold, silver, beryllium, uranium, rubidium, boron, cobalt, and nickel resources.	Dissolution possible for limestone interbeds.	Units weather to produce iron-, magnesium-, and aluminum-rich soils.	Fine for most recreation unless heavily altered and/or dissolved locally; may interfere with compass readings (TRb is strongly magnetic).	Basalts contain magnetite recording paleolatitudes of 5 degrees, possibly of hotspot origin correlative with Nikolai Greenstone flood basalts? TRb has argon isotope date of 179 million years. TRvs correlates with Tangle subterranean. TRll contains fossils found in northern Washington.
PERMIAN	Limestone (Pl)	Unit is approximately 100 m thick with a succession of limestone. Lower beds are medium- to thick-bedded limestone (packstone to grainstone). Upper unit contains light gray silicified mudstone, packstone, and grainstone. Unit weathers to orange-yellow-brown color.	Moderate.	Unit may be friable and too permeable for wastewater treatment infrastructure.	Dissolution may pose sinkhole hazard; susceptible to rockfall if highly dissolved on slopes.	Horridonid brachiopods, bryozoans, pelmatozoan debris, solitary rugose corals, ostracodes, and trilobites.	Caves possible in this unit; may have been settlement areas.	None documented.	Karst features such as dissolution holes and caves are possible.	Unit weathers to basic calcium- and magnesium-rich soils.	Avoid heavily dissolved areas for recreation; any caves may attract speleologists.	Unit contains megafauna of “Arctic Permian” type.

Map Unit Properties Table: Denali National Park and Preserve (Large Scale [1:63,360] Map Units)

Age	Unit Name (Symbol)	Features and Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
PERMIAN - PENNSYLVANIAN	Permian Mudstone and Graywacke (PPNs)	Unit is at least 200 m thick with thinly bedded mudstone and graywacke. Lower beds are gray-green graywacke turbidites that weather to yellow-brown, composed of lithic sandstones with thin intervals of mudstones. Thin-bedded, medium to dark gray and yellow-brown mudstone and siltstone comprise upper beds.	Moderate.	Unit is relatively weak and unstable for most heavy infrastructure, especially if exposed on slope.	Unit is prone to mass wasting hazards, including slumping, slope creep, and possibly debris flows if unvegetated slopes are water saturated.	Trace fossils of ichnogenera <i>Chondrites</i> and <i>Scalarituba</i> ; some brachiopods, bivalves, and bryozoans.	None documented.	None documented.	Not enough carbonate present.	None documented.	Unstable on slopes. Avoid for most use.	Unit records transition from Pennsylvanian to Permian in the area with conformable contacts.
PALEOZOIC	Tuff (PZt) Argillite and Tuff (PZst) Chert, Argillite, and Graywacke (PZs) Rhyolite Schist (PZr) Greenstone and Metamorphosed Basic to Intermediate Agglomerate (PZga) Marble (PZm) Schist of Sedimentary Origin (PZss)	PZt contains a mélange of dark gray-green andesitic to rhyodacitic composition ash and crystal tuff, subordinate argillite, and cherty argillite, siltstone, and graywacke, and minor volcanic flows. Bedding ranges from fine and flinty to more massive and crystalline. PZst contains similar assemblages with more siltstone and chert. PZs contains bedded chert, carbonaceous, fissile black argillite with yellow- and orange-weathering salts, basalt, cherty tuff, green volcanoclastic rock, siltstone, graywacke, and white quartz-pebble and black argillite conglomerate. PZr is metamorphosed silica-rich volcanic deposits. Units become less volcanic to more sedimentary, including metamorphosed limestone (marble) (PZm) and other mixed, slightly metamorphosed schists (PZss).	Moderately high.	Heterogeneous nature of units may render them unstable on slopes. Some units may contain radon-emitting materials and shrink-and-swell clays if altered.	Unit is susceptible to slope processes, including blockfall, landslides, slumping, and slope creep for finer grained units.	Radiolaria.	Massive chert deposits may have provided tool material.	Volcanic breccia, red argillite, and marble. Building material.	Some dissolution possible in limestone interbeds.	None documented.	Units fine for most recreation unless heavily altered.	Records basinwide volcanism and coincident sedimentary deposition as part of “West Fork terrane.”
MISSISSIPPIAN	Moose Creek Member (Mtmg)	Unit is composed of slightly metamorphosed basaltic to intermediate composition volcanic rocks. Green metavolcanic schists dominate this unit.	Moderately high.	Unit is fine for most infrastructure unless heavily altered and/or fractured.	Rockfall hazard when unit is exposed on slope.	None.	None documented.	Chlorite, epidote, and green micas.	None.	Unit weathers to iron- and magnesium-rich soils.	Units fine for most recreation unless heavily altered.	Unit records Mississippian-age volcanism.
DEVONIAN	Andesitic Tuff and Flows (Dv) Red and Brown Chert (Dc) Limestone (DI)	Dv contains green-weathering, pyroxene andesite tuff and flows, with compositions ranging from island-arc tholeiitic basalt (locally pillowed) to dacitic tuff. The lower contact is interlayered with massive red and brown ferruginous and manganiferous radiolarian chert (Dc). Slight greenschist metamorphism is prevalent with alteration minerals, including chlorite, albite, carbonate, and epidote. DI contains medium- to thick-bedded, medium to dark gray lime mudstone to wackestone that is locally fossiliferous.	Moderately high depending on degree of weathering.	Unit may contain radon-emitting materials and shrink-and-swell clays if altered.	Shrink-and-swell clays may buckle trail bases and may be slippery when water saturated; rockfall hazard associated with resistant units underlain by weaker units.	Radiolaria; DI contains rugose and tabulate corals, brachiopods, conodonts, and assorted megafossils.	Massive chert deposits may have provided tool material.	Manganese- and iron-rich cherts.	Karst features such as dissolution holes and caves are possibly found in DI .	Unit weathers to iron- and magnesium-rich soils.	Units fine for most recreation unless heavily altered.	Dv correlates with Dc and UNKsp as part of an island arc complex; units record Devonian island arc and basin depositional setting.

Map Unit Properties Table: Denali National Park and Preserve (Large Scale [1:63,360] Map Units)

Age	Unit Name (Symbol)	Features and Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
PRECAMBRIAN OR PALEOZOIC	Keevy Peak Formation (PZPCkp) Birch Creek Black Carbonaceous Schist (PZPCbcb) Schist and Slate (PZPCss) Gneiss and Schist (PZPCgs) Schist, Quartzite, and Marble (PZPCsqm) Stretched Conglomerate (PZPCscg) Schist (PZPCsch) Birch Creek Schist (PZPCbc)	PZPCkp contains a heterogeneous mixture of quartz-sericite schist, quartzite, and purple and green schist interlayered with slate arkosic gritlike schist, marble, and scant limy schist. PZPCbcb contains a similar assemblage with local layers of green chloritic and epidotic schist and impure marble. PZPCss contains fine-grained yellow, pale-green, and maroon schist and slate with some quartz-sericite schist and meta-chert. PZPCgs contains microcline and quartz gneiss with interlayered lenses of black carbonaceous schist. Lower units contain brown to gray cataclastic and crystalloblastic textures of interlayered quartz-sericite schist (PZPCsqm, PZPCbc, PZPCsch), chloritic to epidotic schist with some calcite (PZPCbc), stretched conglomerate (PZPCscg), quartzite, and marble present locally.	Moderately high.	Altered schist layers, as well as brittlely deformed layers, render these units locally unstable for heavy infrastructure.	Unit is susceptible to slope processes, including blockfall, landslides, slumping, and slope creep for altered, deformed, and fine-grained units.	None documented; deformation and metamorphism likely obscures any preexisting fossil remains.	Pyrite may have been used for fire starting.	Disseminated pyrite, marble, chlorite, and epidote.	Not enough carbonate or marble present.	Quartzite may form spires attractive to goats and birds.	Altered and deformed areas of units should be avoided for most forms of recreation.	PZPCss and PCPZgs may correlate with Nilkoka Group or Totatlanika Schist; units record deformation and metamorphic conditions of terrane formation.
UNKNOWN	Serpentine Gabbro and Silica-Carbonate Rocks (UNKsp) Aeromagnetic High, Probably Serpentine and Gabbro (UNKsp*)	Units are heterogeneous stacks of serpentinized, chromite-bearing dunite to altered, layered gabbro. Gabbro ranges from fine- to coarse-grained leucocratic to melanocratic altered clinopyroxene–plagioclase-bearing. Unit is altered locally to silica-carbonate magnetic rock.	Moderately high.	Avoid areas of heavy alteration and pervasive fracturing for infrastructure.	Highly altered nature of unit renders it weak on slope; rockfall hazards.	None.	None documented.	Clinzoisite, albite, sericite, chlorite, and epidote; magnetite(?).	None.	Unit contains abundant iron and magnesium for basic soil compositions.	Strong magnetic signature may interfere with compass readings.	Possibly part of feeder dike complex for Dv island arc complex.

Map Unit Properties Table: Denali National Park and Preserve (Small Scale [1:250,000] Map Units)

Age	Map Unit (Symbol)	Unit Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
QUATERNARY	Surficial Deposits, Undifferentiated (Qs)	Unit includes unconsolidated and interlayered silt, sand, and fluvial gravel, as well as glacial deposits—including moraines, till, and outwash. Unit also includes slope colluvium in fan-shaped lobes at the bases of slopes. Some terrace deposits and floodplain deposits present along rivers.	Very low.	High permeability of these units likely not suitable for wastewater treatment facilities.	Units are associated with heavy erosion and mass wasting, including frost creep along moderate slopes.	Modern remains possible; pollen, plant macrofossils, insects.	May contain campsites, settlements and other artifacts, and historic mining areas.	Gravel, sand, silt, placer gold deposits, and other heavy metal sediments.	None.	Units support riparian zones and valley floor vegetation and forests.	Units are fine for most light recreation. Avoid riparian areas and large rubble zones.	Units record surficial landscape evolution throughout the area.
TERTIARY	Kenai Group, Undivided (Tk) Sterling Formation (Tsf) Hornblende Dacite (Thd) Nenana Gravel (Tn) Sedimentary Rocks, Undivided (Tsu) Tsadaka, West Foreland, and Wishbone Formations, Undivided (Ttw)	Tk contains pebble and cobble conglomerate interlayered with medium-grained sandstone, some clay layers, and coal seams. Tsf contains massively bedded, coarse conglomerate that appears orange-tan to light gray in outcrop. Thd contains intrusive igneous hornblende dacite (intermediate silica content) of Jumbo Dome. Tn is more than 1,300 m thick containing conglomerate and sandstone with mudstone, claystone, and lignite interbeds. Unit is well sorted, but poorly consolidated, and appears yellowish to brownish in outcrop. Tsu and Ttw contain mixtures of dark gray shales, yellowish sandstones, multi-colored siltstone, pebble conglomerate, and some claystone and lignite beds locally. Ttw contains cobble to boulder conglomerate with other sedimentary lenses and some tuffaceous beds.	Tn and other sedimentary units are moderately low to moderate; igneous unit is high.	Heterogeneous nature of units renders them less stable on slopes. Avoid highly fractured areas for infrastructure.	Unit is prone to rockfall, especially in boulder conglomerates underlain by poorly consolidated units.	None documented.	None documented.	Plutonic conglomerate clasts in Ttw. Coal seams up to 5 m thick in Tk, lignite.	Not enough carbonate present.	Units support wide range of habitats.	Units are fine for most recreation. Avoid slope toes and highly fractured zones.	Thd has an isotopically-determined age of 2.79 ± 0.25 million years. Tk records estuarine and fluvial depositional environments during the Pliocene to Oligocene.
	Volcanic Rocks, Undivided (Tvu) Tyonek Formation (Tty) Coal-Bearing Rocks (Tcb) Hypabyssal Felsic and Intermediate Intrusive Rocks (Thf) Hypabyssal Mafic Intrusive Rocks (Thm) Andesite and Basalt (Tvb) Granitic and Volcanic Rocks, Undivided (Tiv) Granodiorite to Tonalite (Toem)	Tvu consists of interlayered volcanic flows of compositions ranging from basalt to rhyolite, some pyroclastic rocks, and intrusive rocks present locally. Tty is a mixture of sandstone, siltstone, shale, and claystone with various carbonaceous cements. Locally divided into sandstone member and conglomerate member. Tcb contains cyclic sequences of siltstone, claystone, mudstone, shale, sandstone, subbituminous coal and lignite, quartz, and pebble conglomerates. Thf comprises small stocks, sills, and dikes of rhyolitic to dacitic intrusive rocks. Thm is widely exposed, containing small stocks and irregularly shaped bodies of diorite porphyry, diabase, basalt, and lamprophyre. Tvb contains andesitic breccia and tuff layers with some local basalts and pyroclastic layers. Tiv contains portions of a granitic batholith and erosional remnants of rhyolitic flows. Toem contains biotite, biotite-hornblende, and hornblende granodiorite, quartz diorite, and quartz monzodiorite.	Moderate to high for intrusive igneous bodies; lower for poorly consolidated tuffs and breccias, heterogeneous units, and sedimentary units.	Units are very heterogeneous and may be slightly unstable on slopes; may also contain radon-emitting materials. Avoid for basements and foundations. Shrink-and-swell clays may be present, which could undermine roads and foundations.	Unit is prone to rockfall, particularly in areas where resistant igneous intrusive rocks are underlain by friable, weathered sedimentary rocks.	Scant shell fragments, bioturbation, plant fragments(?).	Chert pebbles in Tcb may have provided tool material.	Geodes; vesicles in Tvb may contain secondary minerals. Subbituminous coal and lignite.	Dissolution of carbonate cements may lead to unit instability.	Units support wide range of habitats and may weather to nutrient-rich soils.	Units are fine for most forms of recreation.	Records deposition environment in Tertiary basins.

Map Unit Properties Table: Denali National Park and Preserve (Small Scale [1:250,000] Map Units)

Age	Map Unit (Symbol)	Unit Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
TERTIARY	Fluviatile Sedimentary Rocks and Subordinate Volcanic Rocks (Tfv) Arkose Ridge Formation (Tar) Granite and Granodiorite (Tegr) Chickaloon Formation (Tch) Basalt (Tb) Volcanic Rocks of the Cantwell Formation (Tcv) Granitic Rocks (Tpgr) Granodiorite and Other Intermediate Plutonic Rocks (Thgd)	Tfv contains interlayered sequences of conglomerate, sandstone, siltstone, and mudstone with a few basalt-andesite flows. Tar is as much as 700 m thick of fluvially deposited feldspathic and biotitic sandstone, conglomerate, siltstone, and shale layers in coarsening upward sequences. Tegr contains biotite-hornblende granite and granodiorite present in irregular bodies. Some altered biotite and hornblende tonalite present locally. Tch contains a heterogeneous mixture more than 1,500 m thick of carbonaceous mudstone, siltstone, conglomeratic sandstone, and conglomerate with interbedded bituminous coal beds. Tb contains vesicular olivine basalt that appears brownish-black in outcrop. Tcv contains deformed sequences of andesite, altered basalt, rhyolite and interlayered dacite flows, felsic pyroclastic rocks, and minor sandstone and mudstone. Some calcareous rocks present locally. Tpgr is widespread unit of biotite-muscovite granite and quartz monzonite present generally in plutonic bodies. Thgd is biotite and biotite-hornblende granodiorite, with some smaller quartz diorite, monzonite, and monzodiorite plutons present locally.	Moderate to moderately high for sedimentary rocks; high for intrusive igneous rocks.	Heterogeneous nature of sedimentary layers may be unstable on slopes. Units may contain radon-emitting materials. Avoid for basements and foundations; fine for most infrastructure unless highly fractured.	Heterogeneous layers of these units may be prone to mass wasting and landslides on steep slopes; shrink-and-swell clays may be present in volcanic ash units.	Carbonized plant fragments (may correlate with Cantwell Formation).	Crystallized vesicles may have provided trade material.	Geodes; vesicles may contain secondary mineral crystallization, pillow lavas, and muscovite. Bituminous coal; may contain copper-bearing mineral ores.	Minor dissolution of carbonate cements.	Units may degrade into relatively magnesium-, iron-, and calcium-rich soils.	Units are fine for most recreation unless unaltered crystalline or glassy areas persist in basalt flows, rendering them unsafe for foot traffic.	Tar records deposition on alluvial fans overlain by braided stream deposits throughout the Tertiary and Cretaceous transition. Tegr dated at 45, 44, 48, and 52.8 million years. Tcv dated at 49 million years using potassium-argon isotopes of tuffs. Tpgr has dates between 52 and 65 million years. Thgd dated at 58 to 66 million years.
CRETACEOUS-TERTIARY	Rhyolite and Related Rocks (TKvr) Andesite and Related Rocks (TKvi) Intrusive Rocks (TKi) Granitic Rocks (TKg) Granodiorite, Tonalite, and Monzonite Dikes, and Stocks (TKgd) Gabbro and Leucogabbro (TKgb) Gneissose Granitic Rocks (TKgg)	TKvr consists of light gray to pink rhyolitic volcanic rocks with minor dacite, including flows, tuff, welded(?) tuff, and volcanic breccias. TKvi is predominantly andesitic, but has some minor dacite and basalt flows, tuffs, and breccias present locally. TKi contains intrusive rocks ranging from granite to diorite in composition, mostly intermediate silica contents. TKg is a widespread unit containing biotite and biotite-hornblende granite and lesser amounts of Granodiorite, quartz monzonite, and alkali granite locally. TKgd contains monzonite bodies and other intermediate silica composition dikes, stocks, and irregular intrusions. TKgb is present in small plutons of hypidiomorphic granular textured gabbro and leucogabbro. TKgg contains gneissose granodiorite, quartz diorite, and local granite in small plutonic bodies, structural grain strikes northeast-southwest and dips vertically to 80° northwest.	Moderately high to high.	Units may contain radon-emitting materials. Avoid for basements and foundations; fine for most infrastructure unless highly fractured.	Units are associated with blockfall and landslides if present on an undercut slope. Bulbous plutons may be subject to exfoliation (sloughing) of large blocks of rock.	None.	None documented.	Migmatite and mylonite layers. Building material.	None.	Portions of these units follow valley floors and support low forests.	Units are suitable for most recreation unless highly fractured and/or altered, rendering them unstable for trails and other uses.	TKvr age ranges from 70 to 50 million years. TKgd has dates from 59 to 75 million years from potassium-argon isotopes. TKgg has dates from uranium-lead isotopes in zircon and sphene, as well as dates from potassium-argon isotopes in mica, from 70 to 29 million years; youngest ages may relate to unroofing of Denali fault system.

Map Unit Properties Table: Denali National Park and Preserve (Small Scale [1:250,000] Map Units)

Age	Map Unit (Symbol)	Unit Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
PALEOCENE(?) - CRETACEOUS	Cantwell Formation, Sedimentary Rocks Subunit (Kcs) Matanuska Formation (Km) Wilber Creek Flysch (Kwcf) Nelchina Limestone (Knl)	Kcs contains a 4,000-m-thick interlayered sequence of polymictic conglomerate, sandstone, arkosic sandstone, siltstone, argillite, and shale, and a few thin coal beds. Km contains shallow marine shales with calcareous concretions. Unit appears well-indurated, dark gray, and thinly bedded. Some volcanic-lithic siltstone, sandstone, graywacke, and subordinate conglomerate present locally. Kwcf is a mixture of massively bedded siltstone, shale, sandstone, and conglomerate of clasts derived from quartzite, limestone, mafic and felsic igneous rocks, greenstone, diorite, sandstone, siltstone, phyllite, chert, rare grit, shale rip-ups, and very rare carbonatite. Units appear dark gray to greenish in outcrop and are well-sorted. Knl is shallow-water calcareous sandstone, siltstone, claystone, and thick-bedded to massive clastic limestone. Some conglomerate beds present locally.	Moderate.	Units are fine for most infrastructure unless highly heterogeneous and/or fractured; units may contain radon-emitting materials and/or shrink-and-swell clays from altered volcanic ash.	Unit is associated with slope processes, including landslides.	Paleocene(?) plant fossils; <i>Paragastroplices flexicostatus</i> of Albian age. Regionally, concretions in Km can be highly fossiliferous. Kcs contains dinosaur foot prints and other plants and traces.	None documented.	Calcareous concretions; carbonatite. Thin coal beds.	Knl may contain karst features such as dissolution holes and caves; some areas may be too clastic for extensive dissolution.	Unit may weather into calcium-, iron-, magnesium-, and aluminum-rich soils.	Units are suitable for most recreation unless highly fractured and/or altered	Kcs contains tuffs with potassium-argon isotope dates from Cretaceous. Km records shallow marine deposition environment during the Cretaceous between volcanic arcs.
CRETACEOUS	Mélanges of the Alaska Ridge (Kmar) Volcanic Rocks (Kvl) Granitic Rocks (Kg) Mafic and Ultramafic Rocks (Kmum)	Kmar contains four major suites: (1) cherty tuff, chert, argillite, and volcanoclastic sandstone; (2) dark gray to black argillite, slate, shale, graywacke, and subordinate chert, chert-pebble conglomerate, and polymict conglomerate; (3) limestone; and (4) mixed ultramafic rocks. Kvl includes dacite, andesite, basalt and assorted flows, tuffs, and dikes and sills locally; alteration of most units is widespread. Kg contains granitic to dioritic dikes, sills, and plutons. Kmum contains mixtures of gabbro, diorite, serpentinite, and mafic volcanoclastics. Most layers are altered and deformed to varying degrees present in lenses and dikes.	Moderate to high for granitic rocks.	Units may contain radon-emitting materials and/or shrink-and-swell clays from altered volcanics; heterogeneous layering may be unstable for foundations.	Heavy alteration associated with units render them likely to fail on slopes and if water saturated.	<i>Buchia</i> bivalves of Late Cretaceous to Jurassic age; Radiolaria and conodonts fragments.	Chert may have provided early tool material.	Tourmaline in Kg . Abrasives and building materials.	Kmar suite 3 is subject to karst dissolution.	None documented.	Highly altered units are unstable base.	Kmar records a nearshore environment caught up in a narrow belt during orogenesis. Kg has dates from 120 to 65 million years.
CRETACEOUS AND JURASSIC	Argillite, Chert, Sandstone, and Limestone (KJs) Kahlitna Flysch Sequence (KJf) Flysch Sequence 1 (KJfk) Flysch Sequence 2 (KJfn) Conglomerate, Sandstone, Siltstone, Shale, and Volcanic Rocks (KJcg)	KJs contains dark gray to greenish argillite interlayered with chert, sandstone, and thin shelly limestone layers. KJf , KJfk , and KJfn contain intensely deformed and metamorphosed turbidites. Units include dark gray to black argillite, graywacke, pebble conglomerate, black chert, and dark gray impure limestone beds. Units are tightly folded and faulted. KJcg is interbedded pebble and cobble conglomerate with sandstone, siltstone, shale, and volcanic flows and dikes (andesite to latite).	Moderate.	Units are fine for most infrastructure unless highly fractured, altered, and/or heterogeneous.	Intense deformation of unit renders it susceptible to fail if slope intersects predominant structural fabric.	Ammonites; bivalves <i>Inoceramus</i> , <i>Buchia sublaevis</i> , <i>B. keyserlingi</i> in coquina, <i>Inoceramus hobetsensis</i> ; broadleaf plant fossils.	None documented.	Feldspar porphyry in KJcg .	Some dissolution of limestone in KJs .	None documented.	Units can be too friable for climbing.	Argon isotopes of hornblende clast dates at 94.6 ± 1.5 to 101.0 ± 0.7 million years.

Map Unit Properties Table: Denali National Park and Preserve (Small Scale [1:250,000] Map Units)

Age	Map Unit (Symbol)	Unit Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
JURASSIC	Trondhjemite (Jtr) Naknek Formation (Jn) Chinitna Formation, Tuxedni Group, and Coeval Sedimentary Rocks (Jct) Mafic and Ultramafic Rocks (Jmu) Alaska-Aleutian and Chitina Valley Batholiths, Undifferentiated (Ji)	Jtr is foliated, altered, and deformed trondhjemite in northeast-trending plutons. Jn includes more than 1,400 m of thin- to thick-bedded gray siltstone, shale, sandstone, and conglomerate. Jct is a mixture of shallow marine clastic rocks ranging from conglomerate to siltstone to shale. All beds contain some volcanic elements, likely from nearby island arcs. Jmu contains dunite, harzburgitic dunite, wehrlite, websterite, alkali gabbro (monzogabbro), and clinopyroxenite in discontinuous belts, dikes, and sills. Ji contains widespread batholithic rocks of granite, granodiorite, quartz diorite, and tonalite as part of the Alaska-Aleutian Range batholith.	Moderate to high for granitic batholith units.	Heterogeneous nature of sedimentary layers may be unstable on slopes. Units may contain radon-emitting materials. Avoid for basements and foundations.	Sheet-like exfoliation weathering of plutonic units render them susceptible to large blockfall.	Megafossils; <i>Buchia</i> bivalves, ammonites.	None documented.	Olivine, phlogopite, and clinopyroxene. Abrasives and building materials.	Not enough carbonate present.	None documented.	Fine for most recreation unless highly fractured and/or altered.	Jtr potassium-argon isotope ages range from 129 to 149 million years. Jn constrains the age of uplift of the Alaska-Aleutian Range batholiths.
JURASSIC OR TRIASSIC?	Limestone and Marble (JTRlm) Talkeetna Formation (JTRtk)	JTRlm is fine- to medium-grained, gray metamorphosed limestone in discontinuous lenses as much as 30 m thick. Limestone is massive to poorly bedded. JTRtk contains greenstone and tuff layers. Greenstone is altered equivalent of basaltic lava, agglomerate, breccias, and tuff. Some interlayered sandstone and shale present locally.	Moderate.	Calcareous layers and cements may dissolve, rendering unit unstable in some areas. Tuff units may contain shrink-and-swell clays; avoid for heavy infrastructure.	If carbonate layers and cements are dissolved, rockfall hazards exist; heavily altered layers may fail when water saturated on slopes.	Shallow marine fossils.	Caves possible in this unit; may have been settlement areas.	Marble.	JTRlm is subject to karst dissolution.	Units may weather to produce calcium-, iron-, and magnesium-rich soils.	Fine for most recreation unless heavily altered and/or deformed.	JTRtk is extrusive product of an early Jurassic island arc setting.
JURASSIC - TRIASSIC	Red and Brown Sedimentary Rocks and Basalt (JTRsu) Crystal Tuff, Argillite, Chert, Graywacke, and Limestone (JTRct) Tatina River Volcanic and Equivalent Units (JTRtv)	JTRsu contains a redbed sequence of sandstone, siltstone, argillite, and conglomerates more than 2,000 m thick. Limestone and basalt interbeds present locally. JTRct is a deep marine sequence of tuff, minor chert, argillite, greywacke, and limestone that is heavily faulted and tightly folded. JTRtv consists of three distinct members: (1) upper green medium-grained, concretion-rich volcanoclastic sandstone, with gray, phosphatic shale, and minor tan chert-pebble conglomerate interbeds; (2) lower volcanic rocks, including greenish-gray pillow basalts interlayered with lenses of mudstone, shale, and siltstone gabbro bodies; and (3) brown silty shale and light gray, green, and black chert with some coarse volcanic sandstone and conglomerate.	Moderate.	Heterogeneous nature of units may render them unstable on slopes. Some units may contain shrink-and-swell clays from altered volcanics.	Unit is susceptible to slope processes, including blockfall, landslides, slumping, and slope creep for finer-grained units.	Radiolaria, other Late Triassic and Early Jurassic fossils, bivalve <i>Monotis subcircularis</i> , ammonites.	None documented.	Red chert pebbles. Phosphates and copper ore.	Dissolution is possible within limestone interbeds; may render units unstable.	None documented.	Fine for most recreation unless highly fractured; clay-rich layers may pose hazards to hikers if water saturated.	Records Jurassic – Triassic transition in marine and continental deposition settings.
JURASSIC? AND/OR PERMIAN	Ultramafic Rocks (JPsu)	Unit is a mixture of serpentized (altered) peridotite, harzburgite, and dunite. Unit is highly folded and contains some low-grade metasedimentary rocks and chert layers locally.	Moderate due to high degree of alteration.	Avoid areas of heavy alteration and pervasive fracturing for infrastructure.	Highly altered nature of unit renders it weak on slope; rockfall hazards.	None.	Chert layers may have provided tool material.	Olivine, chlorite, and actinolite. Abrasives.	None.	Unit may weather to produce iron- and magnesium-rich soils.	Unit may interfere with compass readings.	Unit contains oceanic crust caught up during orogenesis.

Map Unit Properties Table: Denali National Park and Preserve (Small Scale [1:250,000] Map Units)

Age	Map Unit (Symbol)	Unit Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
JURASSIC AND PENNSYLVANIAN	Uranatina Metaplutonic Complex (JPNaur)	The complex contains a greenschist facies (locally amphibolite facies) metamorphosed belt of schistose quartz diorite, quartz monzonite, granodiorite, hornblende diorite, amphibolite, and orthogneiss.	Moderately high.	Suitable for most infrastructure. Avoid schistose units and altered areas for heavy infrastructure.	Pervasive foliation and sheet-like exfoliation weathering of metaplutonic units render them susceptible to large blockfall.	None.	None documented.	Building materials.	None.	None documented.	Fine for most recreation unless heavily altered and/or deformed.	Jurassic potassium-argon isotope dates; uranium-lead isotope ages of 309 ± 11 to 153 ± 4 million years.
JURASSIC - CAMBRIAN	Mystic and Dillinger Stratigraphic Sequences, Undivided (JCmd)	Unit is a folded, structurally complex assemblage of siliciclastic turbidites, cherty pelagite, pillow basalt, wildflysch, chert, shale, reefoid limestone, and conglomerate and sandstone representing a mixture of depositional environments.	Moderate.	Heterogeneous nature of units renders them less stable on slopes. Avoid highly fractured areas for infrastructure.	Unit is prone to rockfall, especially in massive beds underlain by deformed units.	Permian- to Silurian-age marine fossils, conodonts.	Chert may have been useful for early tool making.	Basalt alteration minerals and amygdules. Basalts may contain copper-bearing minerals.	Some dissolution is possible in limestone layers.	None documented.	Deformed nature of unit renders it unstable on slopes.	Unit records deposition in deep marine, slope and shelf, and nonmarine settings over millions of years.
TRIASSIC	Calcareous Sedimentary Rocks (TRcs) Conglomerate and Volcanic Sandstone (TRcg) Nikolai Greenstone and Related Rocks (TRn) Gabbro, Diabase, and Metagabbro (TRgb) Metavolcanic and Associated Metasedimentary Rocks (TRnm) Limestone and Basalt Sequence (TRlb) Red Beds (TRr)	TRcs contains thinly bedded, gray calcareous shale, argillite, sandstone, siltstone, and sandy to silty and argillaceous limestone. Unit is intensely deformed. Cherty limestone is present locally, as well as associated dikes, sills, and plugs of altered diabase and gabbro. TRcg is only present locally, and contains cobble to boulder conglomerate with clasts of green volcanic rocks and red radiolarian chert overlain by massive volcanic sandstone. TRn contains greenschist facies metamorphosed subaerial and submarine basalt flows with minor volcanoclastic layers. Unit appears dark gray-green to brown and is more than 2,000 m thick. TRgb is predominantly altered hornblende-clinopyroxene gabbros with quartz gabbro and quartz diorite present in lenses. TRnm contains metabasalt, slate, and other metamorphosed sedimentary and igneous rocks. TRlb is interlayered limestone and amygdaloidal basalt flows. The limestone is massive and gray with local marbleized areas. TRr contains redbeds of sandstone, argillite, siltstone, and conglomerate with clasts of gabbro, serpentinite, and limestone (fossils).	Moderate to moderately high for metamorphic rocks.	Units are highly heterogeneous which may render the rock column relatively unstable. Units may contain radon emitting materials as well as shrink-and-swell clays (altered volcanics). Avoid highly fractured and/or altered areas for infrastructure.	If present on slopes, rockfall, blockfall hazards exist for massive units, whereas slope creep, slumping, and debris flow potential exists for fine-grained, deformed, and/or altered units.	conodonts <i>Negondolella polynathiformis</i> and <i>Epigondolella primitia</i> , bivalve <i>Monotis subcircularis</i> , radiolaria, <i>Heterastridium</i> sp.	None documented.	Red radiolarian chert nodules and mineralized amygdules in basalt flows. Slate, building materials, and marble.	Some dissolution possible in limestone-marble layers.	Units weather into nutrient-rich soils. Dissolution in limestone layers may provide burrow habitat.	Units are suitable for most recreation unless heavily altered, deformed, and/or undercut on slopes.	TRcs records turbidite environment during mid-Triassic. TRn is a diagnostic unit of the Wrangellia terrane. TRlb suggests shallow water deposition in an ocean island shield volcano environment.
TRIASSIC - PENNSYLVANIAN	Flysch-like Sedimentary Rocks (TRPNas)	Unit is pervasively deformed and folded with dark massive to thin-bedded marine Flysch-like layers, including conglomerate, sandstone, siltstone, argillite, limestone, and chert. Unit is several hundred meters thick. Unit is fault-bounded.	Moderate.	Deformed nature of unit may weaken its ability to support heavy infrastructure.	Fractured rocks are susceptible to blockfall and landslides.	Triassic Radiolaria, conodonts, Pennsylvanian brachiopods.	Chert may have been useful for early tool making.	Building material.	Some dissolution is possible in limestone layers.	None documented.	Avoid heavily deformed areas for recreation development.	Fault-bounded unit records structural evolution of the area.

Map Unit Properties Table: Denali National Park and Preserve (Small Scale [1:250,000] Map Units)

Age	Map Unit (Symbol)	Unit Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
TRIASSIC - DEVONIAN	Volcanic and Sedimentary Rocks (TRDv)	Unit contains interlayered greenish-gray to black chert, tuffs, volcanic conglomerate, with some maroon-colored volcanic mudstone, basaltic breccia, and graywacke-shale beds with lenses of limestone, and ammonite-bearing limestone present locally. Unit was metamorphosed to greenschist facies.	Moderate.	Unit may contain shrink-and-swell clays as well as radon-emitting material. Avoid for basements and foundations.	Mudstone-rich units are prone to failure on slopes, especially if water saturated.	Permian brachiopods, ammonites, conodonts. Devonian and Carboniferous fossils in the chert.	None documented.	Attractive maroon mudstone.	Some dissolution possible in limestone layers.	None documented.	Avoid particularly shaly areas for recreation development.	Unit is lower portion of Chulitna sequence.
SILURIAN - TRIASSIC	Limestone Blocks (TRSI)	Unit contains lenses and lozenge-shaped blocks of medium-bedded to massive gray limestone. Limestone is fine- to medium-grained and fossiliferous.	Moderately high.	Unit may be friable and too permeable for wastewater treatment infrastructure.	Dissolution may pose sinkhole hazard; susceptible to rockfall if highly dissolved on slopes.	Megafossils, conodonts.	Caves possible in this unit; may have been settlement areas.	None documented.	May contain karst features such as dissolution holes and caves.	If caves are present, unit may provide bat and other cave dweller habitat.	Any caves will attract caving interest.	Unit records marine basin conditions during a longstanding depositional setting.
MESOZOIC	Phyllite, Pelitic Schist, Calc-Schist, and Amphibolite of the MacLaren Metamorphic Belt (MZpca) Schist and Amphibolite (MZsa)	MZpca is a metamorphosed (amphibolite facies, then retrograde greenschist facies) belt of mainly phyllite, quartz-mica schist, calc-schist, amphibolite, and subordinate marble and meta-andesite, derived from mixed deposits of siltstone, graywacke, marl, andesite, and gabbro. Unit was folded extensively at least twice during orogenesis. MZsa contains fine- to medium-grained schist and hornblende amphibolites, with some irregular bodies of calc-silicate schist and quartzite and local gneissose granitic to gabbroic metaigneous rocks.	High; may be lower with increased degree of weathering.	Altered schist layers, as well as brittlely-ductilely deformed layers, render these units locally unstable for heavy infrastructure.	Unit is susceptible to slope processes, including blockfall, landslides, slumping, and slope creep for altered, deformed, and fine-grained units.	None documented; deformation and metamorphism likely obscures any preexisting fossil remains.	Garnets may have been used for early trade.	Garnets; abrasives.	Not enough carbonate present.	None documented.	Avoid schistose layers for recreation development.	MZpca has potassium-argon isotope dates from mica 48.0 to 30.6 million years and amphibole ages of 69.6 million years. Unit contains Meteor Peak fault. MZsa has potassium-argon isotope dates ranging from 65.9 to 31.9 million years with biotite ages youngest.
MESOZOIC OR PALEOZOIC	Intrusive and Volcanic Rocks, Undivided (MZPZi)	Units composition ranges spatially from granodiorite of Rainbow Mountain, diorite, and gabbro and quartz diorite, with dark gray to greenish dikes of basalt and granodiorite and some volcanics interlayered locally.	High; lower for less consolidated volcanic units.	Altered volcanics may contain shrink-and-swell clays, which could undermine road and building foundations. Avoid fractured areas.	Sheet-like exfoliation of plutonic areas poses blockfall hazard.	None.	None documented.	Building material.	None.	Unit may weather to produce calcium- and aluminum-rich soils.	Suitable for most recreation; may attract climbers. Avoid fractured areas.	Units are widespread and provide unique dating opportunity through isotope dating and cross cutting relationships.

Map Unit Properties Table: Denali National Park and Preserve (Small Scale [1:250,000] Map Units)

Age	Map Unit (Symbol)	Unit Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
MESOZOIC?, PALEOZOIC?, AND (OR) LATE PROTEROZOIC?	Ultramafic and Mafic Rocks, Undivided (MZZum)	Unit is widespread containing a heterogeneous mix of serpentinite, peridotite, dunite, gabbro, diorite, metabasite, and minor talc schist and roddingite. Textures of the different rock types range from intrusive to metamorphic. Devonian-age dikes present north of the Denali fault system.	Moderately high.	Avoid areas of heavy alteration such as serpentinite and talc schist and pervasive fracturing for infrastructure.	Highly schistose and altered nature of unit renders it weak on slope; rockfall hazards.	None.	None documented.	Talc, olivine, and roddingite; abrasives.	None.	Unit may weather to produce iron- and magnesium-rich soils.	Unit unstable on slopes. Avoid for trail development.	Argon isotopes from hornblende dates from 465 to 536 ± 2.6 million years.
PERMIAN	Eagle Creek Formation (Pe)	Unit contains as much as 1,000 m of alternating sequences of marine dark gray, thin-bedded argillite, and fine- to medium-grained limestone beds. Minor lenses of gray siltstone, bioclastic limestone, calcareous siltstone, radiolarian chert, shale, limestone, and pyritic sandstone are present locally.	Moderate.	Dissolved and fractured limestones may render unit too permeable for wastewater treatment.	Limestone dissolution may render units friable with increased fracture diameter, rendering unit weak on slopes.	Corals, bryozoans, brachiopods, echinoids, crinoids, fusulinids, and algal(?) fragments; Radiolaria.	Pyrite may have been used to start early fires.	Pyrite-bearing sandstone.	Limestones may contain karst features such as dissolution holes.	Small holes and cavities may provide burrow habitat.	Avoid areas of dissolution and fracture.	Unit records environmental conditions during the Permian; may contain extinction information.
PERMIAN - PENNSYLVANIAN	Station Creek and Slana Spur Formations, and Equivalent Rocks (PPNasc) Strelna Metamorphic Complex (PPNast) Marble (PPNaskm)	PPNasc contains a lower volcanic member of submarine andesite and basalt flows with breccias and pillows, and an upper volcanoclastic member of graywacke and volcanilutite grading upwards. PPNast is composed of folded, faulted, and metamorphosed metasedimentary and metavolcanic rocks, including quartzofeldspathic and quartz-mica schist with lenses of marble, metachert, and orthogneiss. Volcanic layers are now greenstones. PPNaskm occurs as 30-m-thick lenses of white, medium- to coarse-grained marble.	Moderately high.	Heterogeneous and deformed nature of units may render them unstable on slopes. Some units may contain radon-emitting materials and shrink-and-swell clays if altered.	Units are susceptible to slope processes, including blockfall, landslides, slumping, and slope creep for finer-grained units.	Late Paleozoic bryozoans?	None documented.	Migmatite and greenstones; marble.	Marble may be susceptible to minor dissolution.	None documented.	Units are fine for most recreation unless highly fractured, altered, and/or deformed.	Unit records seafloor volcanism and conditions from Permian and Pennsylvanian.
PERMIAN - DEVONIAN	Sheep Creek Formation and Correlative Siliciclastic Units (PDsc)	Unit is heterogeneous mixture of sandstone, conglomerate, siltstone, and argillite with local beds of chert and limestone. Unit contains clasts of Devonian limestone.	Moderate.	Units are fine for most infrastructure unless highly heterogeneous and/or fractured.	Heterogeneous layers of these units may be prone to mass wasting.	Conodonts, fusulinids, brachiopods, corals, plant remains.	None documented.	Building material.	Some dissolution is possible in limestone layers	None documented.	Units are fine for most recreation.	Unit records deposition spanning from Permian to Devonian.
PENNSYLVANIAN	Tetelna Volcanics (PNat)	Unit appears greenish in outcrop composed of volcanic flows, mud, and debris flows, with lapilli-tuff interbedded with volcanoclastic rocks. Composition is mostly andesitic, and the entire unit was metamorphosed at lower greenschist facies.	Moderately high.	Unit may contain radon-emitting materials and shrink-and-swell clays if altered.	Sedimentary rocks may weather out differentially, posing a rockfall hazard.	None.	None documented.	Lapilli.	None.	May weather to produce aluminum- and calcium-rich soils.	Units are fine for most recreation.	Unit records widespread Pennsylvanian volcanism and later metamorphic-deformation conditions.

Map Unit Properties Table: Denali National Park and Preserve (Small Scale [1:250,000] Map Units)

Age	Map Unit (Symbol)	Unit Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
MISSISSIPPIAN - DEVONIAN	Totatlanika Schist (MDt)	Unit is a mylonite assemblage of gritty semischist with intense deformation and low-grade metamorphic minerals and textures. Some chloritic quartzo-feldspathic schist and augen gneiss, phyllitic schist and semischist, phyllite, metavolcanic rocks, quartzite, marble, and greenstone are present locally with original textures often evident.	Moderate.	Altered schists and phyllite layers render these units locally unstable for heavy infrastructure.	Unit is susceptible to blockfall, landslides, slumping, and slope creep for altered, deformed, and fine-grained units.	None documented; deformation and metamorphism likely obscures any preexisting fossil remains.	Augen may have been valuable trade material.	Clear to bluish gray quartz augen (“eyes”). Marble and abrasives.	Some minor dissolution possible in marble.	None documented.	Avoid schistose areas for recreation development.	Uranium-lead isotopes in zircon provide an age of 375 million years; dates extrusive protolith.
MISSISSIPPIAN? – DEVONIAN?	Keevy Peak Formation (PZk)	Unit is a heterogeneous mixture of phyllite, meta-argillite, quartzite, metachert(?), and lesser amounts of calcareous phyllite, marble, and mafic and felsic metavolcanic rocks. Unit is multiply deformed and metamorphosed at greenschist facies.	Moderate.	Altered schist layers, as well as brittlely deformed layers, render these units locally unstable for heavy infrastructure.	Unit is susceptible to blockfall, landslides, slumping, and slope creep for altered, deformed, and fine-grained units.	None documented; deformation and metamorphism likely obscures any preexisting fossil remains.	None documented.	Phyllite; marble.	Some minor dissolution possible in marble.	None documented.	Altered and deformed areas of units should be avoided for most forms of recreation.	Uranium-leab isotopes in zircon dates extrusion age at 346 ± 1 million years.
DEVONIAN	Serpentinite, Basalt, Chert, and Gabbro (Dsb) Phyllite, Slate, Siliceous Siltstone, and Argillite (Dps) Yanert Fork Sequence and Correlative Rocks (Dy)	Dsb contains sheared and altered serpentinite, chert, and pillow basalt forming lenticular blocks. Dps contains gray to silvery phyllite, slate, siliceous siltstone, and argillite with some thin limestone and calcareous siltstone interbeds. Dy contains metasedimentary and metavolcanic rocks of varying grade and fabric, including argillite, slate, phyllite, phyllonite, semischist, impure quartzite, schistose stretched-pebble conglomerate, banded metachert, felsic metatuff, and metabasalt. Some original pillows and layering present locally.	Moderate.	Sheared and altered areas should be avoided for heavy infrastructure; heterogeneous nature of units render them locally unstable.	Large blocks of massive to intact rock surrounded by sheared and/or altered rock pose blockfall hazards. Schistose cleavage may cause large slides locally, depending on orientation with slope.	Radiolaria.	None documented.	Blocky building material and copper-bearing minerals.	Some dissolution possible in limestone layers.	Limestone dissolution holes may provide burrow habitat. Units may weather into nutrient-rich soils.	Units are fine for most recreation unless highly fractured.	Dsb was part of a dismembered ophiolite assemblage. Dy records a volcanic-bearing turbiditic marine sequence on a continental slope.
DEVONIAN AND SILURIAN	Unnamed Limestone (DSmdl) Whirlwind Creek Formation and Unnamed Correlative Units (DSwc) Limestone (DSI) Tolovana Limestone (DSt)	DSmdl contains light gray to brown limestone, some meta-limestone locally. DSwc is shallow marine limestone and dolostone with minor fossils. DSI contains gray, massive recrystallized limestone with sandy, silicified limestone and dolostone interbeds. Some brecciated and micritic limestones present locally. DSt contains alternating green and maroon limey mudstone grading upward into yellowish-brown silty and shaly lime mudstone and wackestone, grading into gray peloid and ooid-rich lime packstone and grainstone and scant dolostones present locally.	Moderate, depending on degree of alteration and dissolution of carbonate layers.	Unit may be friable and too permeable for wastewater treatment infrastructure.	Dissolution may pose sinkhole hazard; susceptible to rockfall if highly dissolved on slopes.	Graptolites, <i>dasclydacean</i> algae, corals, stromatoporoids, brachiopods, conodonts, Ostracodes.	Caves possible in this unit; may have been settlement areas.	Ooids; limestone building material.	Karst features such as dissolution holes and caves are possible for DSwc and DSt .	If caves are present, they may provide habitat for bats and other cave dwellers.	Caves if present may attract speleologists.	Unit records Silurian deep-water conditions and Late Devonian shallow-water depositional environment.
DEVONIAN - CAMBRIAN	Dillinger Sequence, Undivided (DCd)	Unit contains mixtures of shales, siltstones, cherts, and silty limestones as slope or basinal deposits	Moderate; lower for very hetero- geneous areas.	Heterogeneous nature of unit may render it unstable on slopes.	If carbonate cements are dissolved, integrity of unit may be compromised; possible slope processes hazards.	Conodonts, graptolites, corals.	Chert layers may have provided tool material.	None documented.	Some dissolution possible in limestone layers.	Unit supports wide variety of habitats.	Unit is fine for most recreation.	Unit contains turbidites from an ancient marine slope environment.

Map Unit Properties Table: Denali National Park and Preserve (Small Scale [1:250,000] Map Units)

Age	Map Unit (Symbol)	Unit Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
SILURIAN - CAMBRIAN	Post River Sandstone, Lyman Hills Formation, and Correlative Units (SCpl)	Unit contains sequences of shale, siltstone and chert, grading into silty limestone and shale with cross lamination textures.	Moderate.	Suitable for most infrastructure unless highly fractured.	If massive units are underlain by shale units, a hazardous rockfall situation can exist.	Cambrian conodonts, Ordovician and Silurian graptolites.	Chert layers may have provided tool material.	None documented.	Minor dissolution possible in limestone layers.	None documented.	Unit is fine for most recreation.	Unit records basinal – slope depositional setting with turbidity currents.
ORDOVICIAN	Chert (Oc)	Unit contains dark gray to black chert layered with siliceous slate, argillite, and some greenstone, impure limestone, and dolostone.	Moderately high.	Suitable for most infrastructure unless highly fractured.	Slate layers may undermine unit stability on a slope causing a blockfall hazard. If carbonate cements are dissolved, unit may be prone to failure.	Graptolites.	Chert layers may have provided tool material.	Slate.	Minor dissolution possible in limestone layers.	None documented.	Avoid slate-rich layers due to sharp surfaces.	Unit records Ordovician depositional setting in open basin.
CAMBRIAN? AND PROTEROZOIC	Wickersham Grit, Undivided (CZw) Argillaceous Upper Unit (CZwa)	CZw is a poorly sorted sequence of quartzite, feldspathic quartzite, grit, calcareous siltstone, and fine-grained sandstone, and some scant dark limestone and chert layers. Unit is brittlely deformed and recrystallized with imbricate structures. CZwa is argillaceous with tan-weathering calcareous siltstone and sandstone, with some quartzite and chert present locally. Some dark limestone thin beds interlayered locally.	Moderate; deformed areas are more susceptible to erosion.	Pervasive deformation compromises stability of unit for heavy infrastructure. Avoid fractured areas for wastewater treatment facilities.	Pervasive fracturing incites a likely blockfall hazard.	None documented; deformation and metamorphism likely obscures any preexisting fossil remains.	None documented.	Cataclasite. Quartzite building materials.	Minor dissolution possible in limestone layers.	None documented.	Avoid unit for most recreation due to high fracture density.	Unit records deposition in nearshore environments as well as nature of deformation during orogenesis and uplift.
PALEOZOIC	Spruce Creek Sequence and Correlative Rocks (PZsc) Volcanic and Sedimentary Rocks (PZvs) Calcareous and Phyllitic Rocks (PZkcp)	PZsc contains mafic to felsic metavolcanic rocks, some minor greenstone (altered basalt) sills. Mafic layers are dark, massive; and schistose with metabasalt, andesitic, and amphibolitic layers with minor quartz to mafic schist, pelitic schist, phyllite, metasilstone, metatuff, and minor chert and marble. Felsic layers contain blastoporphyritic metarhyolite and felsic schist. PZvs contains dark, greenish-gray mafic volcanic rocks and black phyllite, chert pebble conglomerates, tuffs, and graywackes. Basalt pillows, breccias, and flows present in volcanic rocks. PZkcp contains greenschist facies calc-phyllite, marble, and phyllite.	Moderate.	Units contain heterogeneous assemblages that may weaken rock columns on slopes; altered volcanic units may contain shrink-and-swell clays.	Mass wasting, blockfall, and landslides are a potential hazard associated with these heterogeneous units.	None.	Phenocrysts may have attracted trade interest and historic mining activity.	Quartz, orthoclase, and plagioclase phenocrysts; stibiconite; kermesite; stibnite; pyrite; boulangerite; cervantite; and arsenopyrite. Marble, building materials, copper-bearing minerals, polymetallic vein deposits, tactite, and stratiform occurrences; support placer gold deposits.	Some minor dissolution is possible in marble.	Units may weather to produce nutrient-rich soils.	Units are fine for most recreation unless altered and/or heavily fractured.	Uranium-lead isotopes provide dates between 364 and 375 million years.

Map Unit Properties Table: Denali National Park and Preserve (Small Scale [1:250,000] Map Units)

Age	Map Unit (Symbol)	Unit Description	Erosion Resistance	Suitability for Infrastructure	Hazards	Paleontological Resources	Cultural Resources	Mineral Occurrence	Karst Issues	Habitat	Recreation	Geologic Significance
PALEOZOIC AND(OR) PROTEROZOIC?	Quartz and Pelitic Schist of the Yukon-Tanana Upland (PZZYqs) Schist and Amphibolite (PZZYsa)	PZZYqs contains multiply metamorphosed quartzite, schistose quartzite, and quartz-mica schist with minor gritty quartzite, chlorite schist, calc-silicate schist, marble, magnetite-biotite schist, amphibolite, and greenstone. PZZYsa contains pelitic schist containing subordinate quartzite, quartz-schist, calc-silicate rocks and calc-schist, marble, amphibolite, graphitic schist, and augen gneiss interlayers. Unit is largely fault-bounded. Unit contains local augen gneiss.	Moderately high.	Avoid schistose layers for heavy infrastructure.	Schistose nature of units renders them friable, and—when combined with layers of more resistant, hard rock—a dangerous rockfall situation occurs.	None.	Augen may have provided valuable trade materials; historic mining activity.	Augen gneiss. Marble, magnetite, graphite, and polymetallic mineral deposits.	Some minor dissolution possible in marble.	Some high calcium soils could be derived from these units.	Avoid friable schistose layers for heavy recreation.	Uranium-lead isotopes in zircon provide dates of 365 million years for protolith, potassium-argon isotopes providing dates ranging from 105 to 120 million years for metamorphic event timing.
PALEOZOIC OR PRECAMBRIAN	Pelitic and Quartzose Schist of the Alaska Range (PZZaqs) Pelitic and Quarzitic Schist (PZZrqs)	PZZaqs contains pelitic and quartzose schist with local calc-schist and feldspathic layers at greenschist to amphibolite facies metamorphic grade. PZZrqs is composed mostly of muscovite- and quartz-rich schist, and lesser calc-schist, quartzofeldspathic schist, gneiss, amphibolite, and quartzite, with scant marble interbeds. Metamorphic history includes greenschist, blueschist, granulite, and amphibolite facies conditions overprinting each other.	Moderate.	Avoid schistose layers for heavy infrastructure.	Schistose nature of units renders them friable and prone to rockfall and failure on a slope.	None.	None documented.	High temperature and pressure metamorphic minerals. Thin marbles and abrasives.	Some minor dissolution possible in thin marble interbeds.	Quartzite may form spires attractive to goats and birds.	Avoid friable schistose layers for heavy recreation.	PZZrqs records conditions during extreme pressure and temperature of orogenesis.
PROTEROZOIC	Gritty Lower Unit (Zwg) Metamorphic Basement Rocks of the Nixon Fork Sequence, Undivided (ZYnm)	Zwg contains poorly sorted quartzite and gritty quartzite with conspicuous granule conglomerate (contains white to blue quartz augen in a cherty, quartzofeldspathic-wacke matrix). ZYnm is metamorphosed (greenschist facies) quartz and pelitic schists with some heterogeneous pods, lenses, and layers of calc-schist, quartzofeldspathic schist, marble, schistose felsic metavolcanic rocks, greenstone, and gneissic and schistose plutonic rocks.	High to moderately high for schistose layers.	Avoid highly fractured areas for infrastructure; fine for most light infrastructure.	Blockfall and landslides are a potential hazard associated with these heterogeneous units.	None documented; deformation and metamorphism likely obscures any preexisting fossil remains.	Augen may have provided valuable trade materials.	Single-crystal milky white to blue quartz augen. Marble, abrasives and attractive building stones.	Some minor dissolution possible in marble.	None documented.	Units are fine for most recreation. Avoid micaceous areas for climbing.	Unit records beach deposits? Potassium-argon isotope ages date metamorphism at 296 to 291 million years; Uranium-lead isotopes in zircon provide ages of 1,250 ± 50 million years.
UNKNOWN	Bedrock of Unknown Type or Age (UNKbu) Ultramafic and Associated Rocks (UNKmlu)	UNKmlu contains serpentinized (altered) ultramafic rocks, as well as altered basalts, tuffs, and recrystallized chert. Unit appears green to maroon in a small sliver of outcrop. UNKbu contains a mixed rubble of metasiltstone and chert, as well as unknown bedrock areas.	Moderate.	Avoid areas of heavy alteration and/or a high concentration of fractures and joints.	Units are altered and poorly consolidated in areas, rendering them susceptible to rockfall and mass wasting.	Fossils possible, but none were specified.	None documented.	Secondary mineralization in basalt vesicles. Attractive green and red stones.	Unknown.	Unit supports development of iron- and magnesium-rich soils.	Units are fine for most recreation, although high iron content may interfere with compass reading.	Unit presents future research possibilities to determine precise timing and provenance.

Geologic History

This section describes the rocks and unconsolidated deposits that appear on the digital geologic map of Denali National Park and Preserve, the environment in which those units were deposited, and the timing of geologic events that created the present landscape.

Ongoing research continues to yield new interpretations of the complex geologic history of Alaska and the Denali National Park and Preserve area.

Precambrian (prior to 542 million years ago)

The geologic story of Denali National Park and Preserve begins in Precambrian time (fig. 21). The oldest rocks in the park are the metamorphosed and deformed remnants of near shore beach, shallow to deep marine, and volcanic deposits. These rocks were deposited in an ancient ocean basin off the coast of the North American craton. Approximately 1 billion years ago, during the Proterozoic Eon, quartz-rich sands and muds were being shed from the ancient craton and deposited in a shallow sea. During this time, North America was part of a large supercontinent known as Rodinia (Condie and Sloan 1998).

Paleozoic Era (542 to 251 million years ago)

During the early Paleozoic, volcanic activity began in the marine basin, possibly accompanying the breakup of Rodinia, spreading basaltic lava and erupting volcanic ash to mix with the terrigenous sediments. In shallow sea areas, during the Cambrian, Silurian, and Devonian periods, marine animals (including corals) thrived, creating layers of limy muds mixed with the quartz-rich sands and volcanic layers in the park (Gilbert 1979).

Following deposition, these early deposits experienced several episodes of folding deformation and metamorphism under extreme temperature and pressure conditions. Metamorphism is bracketed between the middle Paleozoic age of the youngest protoliths and the widespread mid-Cretaceous granitic intrusions (Dusel-Bacon et al. 1993). Deformation under conditions that produce greenschist and amphibolite rocks resulted in northwest-trending folds. A second episode at lower greenschist-facies conditions overprinted these folds with northeast trending folds during the Cretaceous (Dusel-Bacon et al. 1993). During greenschist to amphibolite-facies conditions, quartz sand deposits were hardened into quartzites; mixed sand, mud, and volcanic layers were converted to micaceous schists, gneisses, and phyllites; and limy mud deposits were metamorphosed into marble.

This assemblage of rocks forms much of the widespread Yukon-Tanana terrane, which creates the basement complex of much of interior Alaska (Gilbert 1979; Moore et al. 1994). It is separated from younger terranes toward the south within the park by the Hines Creek fault, the northern branch of the Denali fault system. Timing of deformation for the rocks north of the Hines Creek fault

is poorly constrained between fossil ages from the middle Paleozoic protoliths and undeformed Mesozoic age granitic igneous intrusions (Dusel-Bacon et al. 1993).

As early as the Middle Silurian, intermittent periods of tectonic collision and subduction of oceanic crust and occasional island arcs were occurring west of the North American continent, slowly adding pieces to the continental land mass. At least two subduction zones were present during the Mississippian, Pennsylvanian, and Permian periods. There is no evidence of significant mountain building (orogenesis) during this time in the Denali area. In the western conterminous United States, however, the Antler terrane was accreted during the Mississippian Antler orogeny (Condie and Sloan 1998).

Mesozoic Era (251 to 65.5 million years ago)

The Antler orogeny started a long sequence of accretions along the western edge of the continent as the plates (Kula-Farallon-Pacific and North American plates) collided and slid past each other. The accretion of the metamorphosed and deformed ocean basin and volcanic flows of the Yukon-Tanana terrane onto the North American craton in the Denali area was possibly completed by about 225 million years ago in the early Triassic or prior to the Jurassic (Moore et al. 1994). Several large terranes, including Sonomia (Middle Triassic) and Stikinia (Late Triassic), were added to western Canada.

South of the Hines Creek fault, part of other tectonic terranes (Pingston, Windy, Dillinger, and McKinley terranes) are belts of metamorphic rocks containing gray slate, marble, metamorphosed intrusive mafic rocks (gabbros), and greenstones (altered basalts) (Gilbert 1979; Lillie 2005). These rocks are part of a complex suture zone disrupted by strike-slip faulting along the Denali fault system (described below) and thrust or normal bounding faults (Coney et al. 1980). These rocks were deposited in the latter half of the Paleozoic and beginning of the Mesozoic Era as muds, sand, and fossil debris in a deep ocean basin at tropical latitudes. Marine environments dominated the area for hundreds of millions of years. Some early minor deformation created fissures, allowing basaltic magma to intrude as plutons and extrude as flows (Gilbert 1979).

As the Peninsular and Wrangellia segments of the composite Talkeetna superterrane (elsewhere called the southern Alaska or Wrangellia composite terrane) were being dragged toward the southern boundary of the continent, the sediments trapped between the approaching terrane and the continent were squeezed,

metamorphosed, and deformed to form a northeastward tapering wedge of rocks (Farewell terrane?) called flysch. This accompanied an eastward-increasing metamorphic grade sequence during the Late Mesozoic (Dusel-Bacon et al. 1993). The Upper Jurassic-Cretaceous Kahiltna assemblage records a basin that persisted between the terranes and now contains flysch (Ridgway et al. 2002). Some portions of the flysch wedge are exposed in mountains of the eastern portion of the park, such as Mt. Pendelton and Scott Peak. Masses of accreted island arcs, forming portions of the Pingston and McKinley terranes, were also caught between southern Alaska and the Talkeetna superterrane about 200 million years ago. The culmination of this event involved the emplacement of structurally interleaved tectonic fragments and low- to medium-grade metamorphism, creating the Maclaren metamorphic belt.

In the Jurassic, the block of rocks that eventually became the Brooks Range in northern Alaska collided with North America, producing high-pressure metamorphism, deformation, and thrusting along the southern margin of the continental terranes (Moore et al. 1994). In the Early Jurassic, northward thrusting off the coast of central Alaska caused terranes to “telescope” together. Large nappes and thrusts were forced onto the southern and western edges of the older accreted terranes (Condie and Sloan 1998). The Nevadan orogeny of the western conterminous United States also occurred during this time. Extreme crustal thickening along the plate margins led to partial melting at depth and the emplacement of Jurassic age igneous batholiths (granite, granodiorite, quartz diorite, and tonalite) of the Alaska-Aleutian Range (Wilson et al. 1998). Between the Denali fault system and the Border Ranges to the south, a medium-grade metamorphic event occurred across most of the southern Peninsular and Wrangellia terranes prior to final accretion onto the continent. This metamorphism accompanied local intrusions of tonalitic to granodioritic plutons of Early to Middle Jurassic age in the Peninsular terrane and of Late Jurassic age in the Wrangellia terrane (Dusel-Bacon et al. 1993). Jurassic age plutons are present in the Talkeetna Mountains (Miller 1994).

Metamorphism within the Chugach accretionary wedge, south of the park, was associated with north-directed underthrusting beneath the Peninsular and Wrangellia terranes beginning in the Early to Middle Jurassic. This event created metabasalts, metachert, and other metasedimentary rocks under greenschist to high-pressure blueschist metamorphic facies conditions (Dusel-Bacon et al. 1993).

Starting in the Jurassic, strong orogenic activity began the formation of the Alaska Range. This was in response to subduction of the Kula Plate beneath southern Alaska. Accompanying subduction was terrane accretion, thrust faulting, crustal thickening, isostatic adjustment, and plutonism (Dumoulin et al. 1998). As the subducting Kula Plate (described below) descended farther into the crust beneath southern Alaska, dewatering of the oceanic crust as well as increasing heat caused local melting. This melting created magma plumes that rose to form

relatively buoyant granitic plutons that now compose the core of the Alaska Range. Some of the oldest plutons are dated at 149 million years ago, whereas the younger plutons are 29 million years old (Wilson et al. 1998).

Intense deformation, plutonism, metamorphism, and volcanism continued in the Denali area in response to subduction and subsequent terrane accretion during the Cretaceous (fig. 22) (Condie and Sloan 1998). During most of the accretion events, rising crustal temperatures led to massive gold vein formation (Goldfarb et al. 1997). Movement along the Denali fault might have started during this time. Accretion of the Talkeetna superterrane—transported from the south along the margin of western Canada by dextral strike-slip movement—was complete by 110 to 85 million years ago during the Cretaceous (Hickman et al. 1990; Miller et al. 2002).

Prior to regional uplift, much of the Denali area was composed of continental lowlands, similar to the areas north of the Alaska Range today. Streams flowed predominantly south toward the coast, depositing alluvial sand, gravel, and silt. During the early stages of uplift, streams began diverting to northerly courses (fig. 23). A pull-apart asymmetrical basin graben developed in the area between the Hines Creek and southern fault strands of the Denali fault system. This so-called Cantwell Basin filled with as much as 4,000 m (13,000 ft) of southward thickening sediments (Hickman et al. 1990).

Cenozoic Era (the past 65.5 million years)

Intense regional volcanism peaked around 56 and 38 million years ago, leading to vast igneous intrusions and deposits of red, yellow, and brown basalts, rhyolites, andesites, and other volcanic rocks (Teklanika Formation). Some of these extrusive rocks are present at Polychrome Pass (older event) and Mt. Galen (younger event) (Cole 2004). Contemporaneous granitic igneous intrusions formed Mt. Eielson and Mt. Foraker (Gilbert 1979). Hydrothermal activity increased as well, leading to the development of gold-bearing quartz vein systems (fig. 24) (Goldfarb et al. 1997). Extensive volcanic flows covered the stream sediments that ultimately became the shale, sandstone, coal beds, and pebble conglomerate of the Tertiary lower Cantwell Formation (Wahrhaftig 1970; Gilbert 1979; Hickman et al. 1990).

During the Jurassic and Cretaceous, three major plates existed west of the North American continent separated by mid ocean spreading ridges: the Kula, Farallon, and Pacific plates. Most remnants of the Kula and Farallon plates were lost to subduction beneath northeastern Asia and western North America, respectively. The Juan de Fuca and Cocos plates in the eastern Pacific basin are remnants of the otherwise subducted Farallon plate. Subduction of the Farallon plate drove the Laramide orogeny spanning from Mexico to Canada from the Late Cretaceous into the Tertiary.

By the mid-Tertiary, the Pacific plate dominated the basin. Dated volcanic layers (53.9 ± 1.6 million years ago based on argon isotopes [$^{40}\text{Ar}/^{39}\text{Ar}$]) in the upper

Cantwell Formation are cut by younger high angle, northwest-trending, right-lateral shear zones, suggesting that the once collisional boundary between Alaska and the Pacific plate was changing to a more translational ("side-swiping") boundary by this time (Cole et al. 1996).

Once the Farallon spreading ridge (East Pacific Rise) was consumed, the margin between the Pacific and western North American plates became more transform in nature, around 30 million years ago (creating the San Andreas fault), and the Pacific plate continued to travel northwestward, subducting beneath Alaska, and dragging terranes with it (Condie and Sloan 1998).

From the Cretaceous to early Tertiary, accompanying the accretion of the Chugach accretionary prism to the south, the southern edge of the Talkeetna superterrane was exposed to further deformation and overprinting of a lower-grade metamorphic event (Dusel-Bacon et al. 1993). The Chugach split into a sequence of accretionary prisms and mélangé complexes that underthrust each other sequentially from south to north, each accompanied by pervasive deformation and metamorphism throughout the Cretaceous into the early Tertiary. The accretionary wedge in its entirety was part of southern Alaska by about 67 million years ago, followed by the accretion of the Prince William accretionary prism at approximately 50 million years ago (Lillie 2005). Between these large terranes, forearc basins, narrow volcanic arcs, and small-scale marine basins received alluvial, volcanic, and shallow marine sediments that record uplift to the north and subsequent erosion of vast amounts of sediments, as well as thrust faulting to the south associated with continued accretion in the Talkeetna area (Trop et al. 2003).

As accretion of the Yakutat Terrane that started ~20 million years ago persists today, low-pressure, amphibolite facies metamorphism and deformation is occurring on the southern coast of Alaska (Dusel-Bacon et al. 1993; Lillie 2005). The Pacific plate continues to move northwestward at a rate of approximately 5 cm (2 in.) per year. This active margin has led to extensive faulting throughout the region over the past 100 million years. Some of this deformation is accommodated along the right-lateral Denali fault system running parallel to the Alaska Range. Fresh fault scarps and offset stream gravels attest to recent movement along the fault (Wahrhaftig 1970), as well as the magnitude 7.9 Denali fault earthquake in 2002 (Eberhart-Phillips et al. 2003). Maximum local uplift rates in Denali are estimated at ~1 km/million years (0.6 mi/million years) beginning in the late Tertiary as a result of changes in motion of the Pacific plate relative to North America. This, in conjunction with the geometry (bend) along the Denali fault, caused local crustal blocks to be rapidly forced upward (Fitzgerald et al. 1993).

Block faulting during the mid-Tertiary created downfaulted inland basins, which collected vast deposits of sediments, including the coal-bearing Suntrana Group (30 to 5 million years ago) (Wahrhaftig 1970; Gilbert 1979). The Tertiary basin coal-bearing rocks record terrestrial cyclic sequences of siltstone, claystone, mudstone, shale, pebbly sandstone, subbituminous coal and lignite, and minor amounts of pebble conglomerate (Wilson et al. 1998). These basins supported vast forests and swampy bog environments (Gilbert 1979). The coal-bearing rocks were covered by sand and gravel shed from erosion of further uplift of the Alaska Range during the Pliocene Epoch (5.3 to 2.6 million years ago). These thick sediments comprise the poorly consolidated Nenana Gravel (Gilbert 1979; Wilson et al. 1998).

Erosion kept pace with exhumation of the Alaska Range until the beginning of Pliocene time. Around 6 million years ago, the Alaska Range began to uplift rapidly, causing most major streams of the central portions of the range to flow north (Gilbert 1979). During the Pliocene–Pleistocene, several global ice age events also occurred. Glacial ice covered vast extents of the Alaska Range, carving U-shaped valleys. Interglacial period climates supported the growth of open boreal woodland, dense alder shrub vegetation, open taiga, graminoid tundra, and birch shrub tundra through at least the past 50,000 years. Holocene coniferous forests only became established after 6,500 years before present (Elias et al. 1996). This vegetation is recorded in recent peat, swamp, bog, and other organic deposits in the Quaternary unconsolidated sediments in the foothills north of the Alaska Range in the park.

Between four and seven major glacial advances altered the landscape at Denali between 2 million and 10,000 years ago (Gilbert 1979; Brease 2004), starting with the Teklanika glacial advance, which began possibly in the late Pliocene to early Pleistocene. Other advances, including the Browne, Bear Creek, Dry-Lignite Creek, Healy-McLeod Creek, and Riley Creek, are also locally recorded in the glacial deposits at Denali. The Carlo Creek Readvance began in the early Holocene, 8,000 years ago (Brease 2004). End units from this advance form lakes that trap fine-grained sands and other lacustrine deposits today (Wahrhaftig 1952). The past glacial advances left ice erosional and glacial depositional features in most of the river valleys within the park. U-shaped valleys are present in the Savage, Toklat, Thorofare, Sanctuary, Teklanika, and McKinley Rivers' drainages. Large glacial erratic deposits are located throughout lower areas of the park (Brease 2004). Glacial processes continue today as smaller glaciers cover more than 1 million acres within Denali National Park and Preserve. Table 1 summarizes the geologic history in the Denali area of Alaska.

Eon	Era	Period	Epoch	Ma	Life Forms	North American Events			
Phanerozoic	Cenozoic	Quaternary	Holocene	0.01	Age of Mammals	Modern humans	Cascade volcanoes (W)		
			Pleistocene			Extinction of large mammals and birds	Worldwide glaciation		
		Tertiary	Neogene	Pliocene		2.6	Large carnivores	Sierra Nevada Mountains (W)	
				Miocene		5.3	Whales and apes	Linking of North and South America	
			Paleogene	Oligocene		23.0		Basin-and-Range extension (W)	
				Eocene		33.9			
				Paleocene		55.8	Early primates	Laramide Orogeny ends (W)	
				65.5					
		Mesozoic	Cretaceous				Age of Dinosaurs	Mass extinction	Laramide Orogeny (W)
						145.5		Placental mammals	Sevier Orogeny (W)
	Jurassic			199.6	Early flowering plants	Nevadan Orogeny (W)			
	Triassic				First mammals	Elko Orogeny (W)			
						Mass extinction	Breakup of Pangaea begins		
						Flying reptiles	Sonoma Orogeny (W)		
						First dinosaurs			
	Paleozoic	Permian			Age of Amphibians	Mass extinction	Supercontinent Pangaea intact		
						Coal-forming forests diminish	Ouachita Orogeny (S)		
							Alleghanian (Appalachian) Orogeny (E)		
							Ancestral Rocky Mountains (W)		
		Pennsylvanian		299	Coal-forming swamps				
					Sharks abundant				
		Mississippian		318.1	Variety of insects				
					First amphibians				
		Devonian		359.2	First reptiles	Antler Orogeny (W)			
					Mass extinction				
		Silurian		416	First forests (evergreens)	Acadian Orogeny (E-NE)			
			443.7						
Ordovician			First land plants						
			Mass extinction						
			First primitive fish	Taconic Orogeny (E-NE)					
			Trilobite maximum						
Cambrian		488.3	Rise of corals						
			Early shelled organisms	Avalonian Orogeny (NE)					
				Extensive oceans cover most of proto-North America (Laurentia)					
					542				
Proterozoic	Precambrian			2500	First multicelled organisms	Supercontinent rifted apart			
					Formation of early supercontinent				
					Grenville Orogeny (E)				
					First iron deposits				
Archean						Abundant carbonate rocks			
Hadean					Early bacteria and algae				
				≈4000		Oldest known Earth rocks (≈3.96 billion years ago)			
					Origin of life?	Oldest moon rocks (4–4.6 billion years ago)			

Figure 21: Geologic timescale. Included are major life history and tectonic events occurring on the North American continent with an emphasis on events affecting Alaska. Red lines indicate major unconformities between eras. Isotopic ages shown are in Ma. Compass directions in parentheses indicate the regional location of individual geologic events. Adapted from the U.S. Geological Survey, <http://pubs.usgs.gov/fs/2007/3015/>, with additional information from the International Commission on Stratigraphy, <http://www.stratigraphy.org/view.php?id=25>.




Figure 22: Three-dimensional model of an active margin setting with successive terrane accretion, uplift, volcanism, and subduction. Graphic by Trista L. Thornberry-Ehrlich (Colorado State University), adapted from Lillie (2005) fig. 11.8.




Figure 23: Reversal of stream direction from southward to northward accompanying uplift of the Alaska Range in the early Cenozoic. Graphic is by Trista L. Thornberry-Ehrlich (Colorado State University), adapted from Gilbert (1979) fig. 13.




Figure 24: Map of Alaska during the Paleocene and Eocene epochs showing generalized locations of major terranes, faults, as well as significant gold districts (black circles). Graphic by Trista L. Thornberry-Ehrlich (Colorado State University), adapted from Goldfarb et al. (1997) fig. 7.

Table 1: Summary of geologic events in the Denali area. Ma = million years ago; ybp = years before present. Colors (also used on the Map Unit Properties Table) according to the Commission for the Geological Map of the World. Table modeled after Brease (2004) table 34.1.

Eon	Era	Period		Epoch	Geologic Events	
PHANEROZOIC	CENOZOIC	QUATERNARY		HOLOCENE	Cantwell ash fall with volcanism from southerly source (~3,700 ybp) and multiple glaciation events.	
				PLEISTOCENE	Multiple glacial advances starting ~2 Ma to 8,000 ybp; intense glacial erosion of rapidly uplifting Alaska Range.	
		TERTIARY	NEOGENE	PLIOCENE	Deposition of Nenana Gravel shed from uplifting Alaska Range; units include sandstone, conglomerates, claystone, and lignite. Uplift and exhumation of Mt. McKinley. Movement along Denali fault system likely resumes.	
				MIOCENE	Deformation and thrust faulting cause a surge in uplift of the Alaska Range; beginning of Yakutat terrane accretion. Regional subsidence due to crustal thickening and continued sedimentation and coal deposits of the Tanana foreland.	
			PALEOGENE	OLIGOCENE	Deposition in subsidence basins north and south of the Alaska Range; several phases of igneous intrusion and volcanism at 38 Ma.	
				EOCENE	Cantwell volcanism and McKinley intrusive sequence from 41 to 57 Ma results in flows, breccias, and tuffs, as well as granodiorite intrusion with some sulfide mineralization; Prince William wedge accretes.	
				PALEOCENE	Emergent Alaska Range continues to shed sediments into foreland pull-apart basin, leading to piles of sandstone, siltstone, shale, tuff layers, coal, and conglomerates of the Cantwell Formation; strike-slip movement along Denali fault. Continued intrusion of granites in the Alaska Range, including the McKinley and Ruth plutons.	
		MESOZOIC	CRETACEOUS		Multiple phases of igneous intrusive activity (granites and granodiorites), volcanism, and orogenesis as the Chugach wedge accretes to the continent; continued flysch deposition in shallow basins; pervasive deformation and metamorphism at 115 to 106 Ma, 74 Ma, and 65 to 60 Ma; uplift of Alaska Range continues. Final closer of ocean between Talkeetna Superterrane, and previously accreted terranes to the north.	
			JURASSIC		Orogenic activity increases as Talkeetna superterrane is accreting, pushing miniterranes within the intervening basin toward the continent; intense deformation and metamorphism; continued deposition of Mesozoic flysch in segmented, forearc, and backarc basins	
	TRIASSIC		Final accretion of Yukon-Tanana terrane; abundant submarine basalt flows form Nikolai Greenstones; continued deposition of redbed sandstones, conglomerates, tuffs, argillites, and limestones; Pingston, McKinley, and Chulitna terranes are pushed toward the margin of North America.			
	PALEOZOIC		PERMIAN		Deposition of alternating limestone and argillite beds of the Eagle Creek Formation, as well as massive marine limestone, mudstone, and greywacke.	
			PENNSYLVANIAN		Widespread volcanism forms andesitic Tetelna Volcanics, later metamorphosed to greenschist; continued marine deposition of cherts, pillow basalts, shales, fossiliferous limestones, sandstones, and argillites.	
			MISSISSIPPIAN		Moose Creek Member basaltic to intermediate volcanism, later metamorphosed to greenschists (Totatlanika Schist).	
			DEVONIAN		Andesitic tuff from island arc volcanism, red and brown chert deposits, and shallow marine basin limestone.	
		SILURIAN		Marine deposition and coral growth; intermittent volcanic activity. Continued deposition of turbidites, sandstones, argillites, dolomitic limestones, cherts, volcanic flows and ash falls, shales, and conglomerates; intermittent igneous activity, including mafic dike intrusions; multiple phases of deformation and metamorphism change sediments to quartzites, phyllites, slates, marbles, gneisses, meta-volcaniclastic schists, and greenstones.		
	ORDOVICIAN					
	CAMBRIAN					
PRECAMBRIAN				Shallow marine basins covered large area south of ancient North American continent; resulted in deposition of quartz-rich sediments interlayered with volcanic flows and ash and limestone. Formations include Keavy Peak Formation and Birch Creek Schist.		

Glossary

This glossary contains brief definitions of technical geologic terms used in this report. Not all geologic terms used are referenced. For more detailed definitions or to find terms not listed here please visit: <http://geomaps.wr.usgs.gov/parks/misc/glossary.html>. Definitions are based on those in the American Geological Institute Glossary of Geology (fifth edition; 2005).

accretionary prism. A wedge-shaped body of rock consisting of rocks scraped off of subducting oceanic crust at a subduction zone. Rocks consist of trench-fill turbidites, seamounts and surrounding reefs. Accretionary prisms form in the same manner as a pile of snow in front of a snowplow.

active margin. A tectonically active margin where lithospheric plates converge, diverge or slide past one another (also see “passive margin”). Typically associated with earthquakes and, in the case of convergent and divergent boundaries, volcanism.

alluvial fan. A fan-shaped deposit of sediment that accumulates where a hydraulically confined stream flows to a hydraulically unconfined area. Commonly out of a mountain front into an area such as a valley or plain.

alluvium. Stream-deposited sediment.

alpine glacier. A glacier occurring in a mountainous region on the sides of a mountain.

altiplanation. Soliflucation and related mass movements that tend to produce flat or terrace-like surfaces, especially at high elevation and latitudes where periglacial processes predominate.

andesite. A dark-colored, fine-grained extrusive rock that, when porphyritic, contains phenocrysts composed primarily of zoned sodic plagioclase and one or more of the mafic minerals with a groundmass composed generally of the same minerals as the phenocrysts.

angular unconformity. An unconformity where the strata above and below are oriented differently; generally caused by structural deformation and erosion prior to deposition of the upper bed.

anticline. A convex-upward (“A” shaped) fold. Older rocks are found in the center.

anticlinorium. A large, regional feature with an overall shape of an anticline. Composed of many smaller folds.

aquifer. A rock or sedimentary unit that is sufficiently porous that it has a capacity to hold water, sufficiently permeable to allow water to move through it, and currently saturated to some level.

arête. A rocky sharp-edged ridge or spur, commonly present above the snowline in rugged mountains sculptured by glaciers, and resulting from the continued backward growth of the walls of adjoining cirques.

argillite. A compact rock, derived from mudstone or shale, more highly cemented than either of those rocks. It lacks the fissility (easily split) of shale or the cleavage of slate. It is regarded as a product of low-temperature metamorphism.

ash (volcanic). Fine pyroclastic material ejected from a volcano (also see “tuff”).

asthenosphere. Earth’s relatively weak layer or shell below the lithosphere.

axis (fold). A straight line approximation of the trend of a fold, that divides the two limbs of the fold.

base flow. Stream flow supported by groundwater flow from adjacent rock, sediment, or soil.

base level. The lowest level to which a stream can erode its channel. The ultimate base level for the land surface is sea level, but temporary base levels may exist.

basement. The undifferentiated rocks, commonly igneous and metamorphic, that underlie the rocks exposed at the surface.

basin (sedimentary). Any depression, from continental to local scales, into which sediments are deposited.

basin (structural). A doubly-plunging syncline in which rocks dip inward from all sides (also see “dome”).

batholith. A massive, discordant pluton, greater than 100 km², (40 mi²) often formed from multiple intrusions.

bed. The smallest sedimentary strata unit, commonly ranging in thickness from one centimeter to a meter or two and distinguishable from beds above and below.

bedding. Depositional layering or stratification of sediments.

bedrock geology. The geology of underlying solid rock as it would appear with the sediment, soil, and vegetative cover stripped away.

bergschrand. The crevasse occurring at the head of an alpine glacier, which separates the moving snow and ice of the glacier from the relatively immobile snow and ice adhering to the headwall of a cirque.

block (fault). A crustal unit bounded by faults, either completely or in part.

blueschist. A schistose metamorphic rock with a blue color due to the presence of sodic amphibole, e.g. the minerals glaucophane or crosstie, and commonly mottled bluish-gray lawsonite. Often associated with high pressure, relatively low temperature metamorphic conditions.

braided stream. A stream, clogged with sediment that forms multiple channels that divide and rejoin.

breccia. A coarse-grained, generally unsorted sedimentary rock consisting of cemented angular clasts greater than 2 mm (0.08 in).

brittle. Describes a rock that fractures before sustaining deformation.

calcareous. Describes rock or sediment that contains calcium carbonate (CaCO₃).

carbonaceous. Describes a rock or sediment with considerable carbon, especially organics, hydrocarbons, or coal.

cataclasite. A fine-grained rock formed by pervasive fracturing, milling, crushing, and grinding by brittle deformation, typically under conditions of elevated pressure.

cementation. Chemical precipitation of material into pores between grains that bind the grains into rock.

chemical sediment. A sediment precipitated directly from solution (also called nonclastic).

chemical weathering. Chemical breakdown of minerals at Earth's surface via reaction with water, air, or dissolved substances; commonly results in a change in chemical composition more stable in the current environment.

chert. A extremely hard sedimentary rock with conchoidal (smooth curved surface) fracturing. It consists chiefly of interlocking crystals of quartz (syn: flint).

cirque. A deep, steep-walled, half-bowl-like recess or hollow located high on the side of a mountain and commonly at the head of a glacial valley. Produced by the erosive activity of a mountain glacier. It often contains a small round lake.

clastic. Describes rock or sediment made of fragments of pre-existing rocks.

clay. Can be used to refer to clay minerals or as a sedimentary fragment size classification (less than 1/256 mm [0.00015 in]).

cleavage. The tendency of a rock to split along parallel, closely spaced planar surfaces. It is independent of bedding and is produced by deformation or metamorphism.

concordant. Strata with contacts parallel to the attitude of adjacent strata.

congeliturbate. Mass of soil or other unconsolidated material moved or disturbed by frost action.

conglomerate. A coarse-grained, generally unsorted, sedimentary rock consisting of cemented rounded clasts larger than 2 mm (0.08 in).

continental crust. The crustal rocks rich in silica and alumina that underlie the continents; ranging in thickness from 35 km (22 mi) to 60 km (37 mi) under mountain ranges.

continental shield. A continental block of Earth's crust that has remained relatively stable over a long period of time and has undergone only gentle warping compared to the intense deformation of bordering crust

convergent boundary. A plate boundary where two tectonic plates are colliding.

cordillera. A Spanish term for an extensive mountain range that is used in North America to refer to all of the western mountain ranges of the continent.

craton. The relatively old and geologically stable interior of a continent (also see "continental shield").

crevasse. A deep fissure or crack in a glacier, caused by stresses resulting from differential movement over an uneven surface. Crevasses may be as much as 100 m (330 ft) deep.

cross section. A graphical interpretation of geology, structure, and/or stratigraphy in the third (vertical) dimension based on mapped and measured geological extents and attitudes depicted in a vertically oriented plane.

cross-bedding. Uniform to highly varied sets of inclined sedimentary beds deposited by wind or water that indicate distinctive flow conditions (e.g., direction and depth).

crust. Earth's outermost compositional shell, 10 to 40 km (6 to 25 mi) thick, consisting predominantly of relatively low-density silicate minerals (also see "oceanic crust" and "continental crust").

crystalline. Describes a regular, orderly, repeating geometric structural arrangement of atoms.

dacite. A fine-grained extrusive rock with the same general composition as andesite, but having a less calcic plagioclase and more quartz.

debris cone. A cone or mound of ice or snow on a glacier, covered with a veneer of debris thick enough to protect the underlying material from ablation.

debris flow. A rapid and often sudden flow or slide of rock and soil material involving a wide range of types and sizes.

deformation. A general term for the process of faulting, folding, and shearing of rocks as a result of various Earth forces such as compression (pushing together) and extension (pulling apart).

dike. A tabular, discordant igneous intrusion.

dip. The angle between a bed or other geologic surface and horizontal.

dip-slip fault. A fault with measurable offset where the relative movement is parallel to the dip of the fault.

disconformity. An unconformity at which the bedding of the strata above and below are parallel.

discordant. Having contacts that cut across or are set an angle to the orientation of adjacent rocks.

divergent boundary. An active boundary where tectonic plates are moving apart (e.g., a spreading ridge or continental rift zone).

drainage basin. The total area from which a stream system receives or drains precipitation runoff.

drift. All rock material (clay, silt, sand, gravel, boulders) transported by a glacier and deposited directly by or from the ice, or by running water emanating from a glacier.

ductile. Describes a rock that is able to sustain deformation before fracturing.

eolian. Formed, eroded, or deposited by or related to the action of the wind.

extrusive. Of or pertaining to the eruption of igneous material onto the surface of Earth.

facies (metamorphic). The pressure-temperature regime that results in a particular, distinctive metamorphic mineralogy (i.e., a suite of index minerals).

facies (sedimentary). The depositional or environmental conditions reflected in the sedimentary structures, textures, mineralogy, fossils, etc. of a sedimentary rock.

fan delta. An alluvial fan that builds into a standing body of water. This landform differs from a delta in that a fan delta is next to a highland and typically forms at an active margin.

fault. A break in rock along which relative movement occurs between the two sides.

fault gouge. Soft, uncemented, pulverized, clay-like material found along some faults formed by friction as the fault moves.

felsic. Describes an igneous rock having abundant light-colored minerals such as quartz, feldspars, or muscovite. Compare to “mafic.”

flysch. A marine sedimentary facies characterized by a thick sequence of poorly fossiliferous, thinly bedded, graded marls and sandy and calcareous shales and muds, rhythmically interbedded with conglomerates (rare), coarse sandstones, and graywackes. Typically deposited in deep ocean basins near convergent plate boundaries and rising mountains.

footwall. The mass of rock beneath a fault surface (also see “hanging wall”).

formation. Fundamental rock-stratigraphic unit that is mappable, lithologically distinct from adjoining strata, and has definable upper and lower contacts.

fracture. Irregular breakage of a mineral. Any break in a rock (e.g., crack, joint, fault).

frost wedging. The breakup of rock due to the expansion of water freezing in fractures.

gelifluction. Progressive lateral flow of earth material under periglacial conditions.

geology. The study of Earth, including its origin, history, physical processes, components, and morphology.

graben. A down-dropped structural block bounded by steeply dipping, normal faults (also see “horst”).

granodiorite. A group of plutonic rocks containing quartz, plagioclase, and potassium feldspar with biotite, hornblende, or, more rarely, pyroxene, as the mafic components.

graywacke. A term commonly used in the field for a dark gray to dark green, very hard, dense sandstone of any composition but with a chlorite-rich matrix; these rocks have undergone deep burial.

greenschist. A schistose metamorphic rock, whose green color is due to the presence of the minerals chlorite, epidote, or actinolite, corresponds with metamorphism at temperatures in the 300–500°C (570–930°F) range.

hanging valley. A tributary glacial valley whose mouth is high above the floor of the main valley, which was eroded by the main body of the glacier.

hanging wall. The mass of rock above a fault surface (also see “footwall”).

horn. A high pyramidal peak with steep sides formed by the intersection walls of three or more cirques.

horst. Areas of relative “up” between grabens, representing the geologic surface left behind as grabens drop. The best example is the Basin-and-Range province of Nevada. The basins are grabens and the ranges are weathered horsts. Grabens become a locus for sedimentary deposition (also see “graben”).

igneous. Refers to a rock or mineral that originated from molten material; one of the three main classes of rocks—igneous, metamorphic, and sedimentary.

intrusion. A body of igneous rock that invades (pushes into) older rock. The invading rock may be a plastic solid or magma.

island arc. A line or arc of volcanic islands formed over and parallel to a subduction zone.

isostasy. The process by which the crust “floats” at an elevation compatible with the density and thickness of the crustal rocks relative to underlying mantle.

isostatic adjustment. The shift of the lithosphere to maintain equilibrium between units of varying mass and density; excess mass above is balanced by a deficit of density below, and vice versa.

joint. A semi-planar break in rock without relative movement of rocks on either side of the fracture surface.

laccolith. A mushroom- or arcuate-shaped pluton that has intruded sedimentary strata and domed up the overlying sedimentary layers. Common on the Colorado Plateau.

lacustrine. Pertaining to, produced by, or inhabiting a lake or lakes.

lamination. Very thin, parallel layers.

landslide. Any process or landform resulting from rapid, gravity-driven mass movement.

lava. Still-molten or solidified magma that has been extruded onto Earth’s surface through a volcano or fissure.

lignite. A brownish-black coal that is intermediate in coalification between peat and subbituminous coal.

limbs. Either side of a structural fold.

lithification. The conversion of sediment into solid rock.

lithology. The physical description or classification of a rock or rock unit based on characters such as its color, mineral composition, and grain size.

lithosphere. The relatively rigid outmost shell of Earth’s structure, 50 to 100 km (31 to 62 mi) thick, that encompasses the crust and uppermost mantle.

loess. Windblown silt-sized sediment, generally of glacial origin.

mafic. Describes dark-colored rock, magma, or minerals rich in magnesium and iron. Compare to “felsic.”

magma. Molten rock beneath Earth’s surface capable of intrusion and extrusion.

mantle. The zone of Earth’s interior between crust and core.

matrix. The fine grained material between coarse (larger) grains in igneous rocks or poorly sorted clastic sediments or rocks. Also refers to rock or sediment in which a fossil is embedded.

meander. Sinuous lateral curve or bend in a stream channel.

mechanical weathering. The physical breakup of rocks without change in composition.

mélange. A mappable body of jumbled rock that includes fragments and blocks of all sizes, both exotic and native, embedded in a fragmented and generally sheared matrix.

member. A lithostratigraphic unit with definable contacts; a member subdivides a formation.

meta-. A prefix used with the name of a sedimentary or igneous rock, indicating that the rock has been metamorphosed.

metamorphic. Describes the process of metamorphism or its results. One of the three main classes of rocks—igneous, metamorphic, and sedimentary.

metamorphism. Literally, a change in form. Metamorphism occurs in rocks through mineral alteration, genesis, and/or recrystallization from increased heat and pressure.

- mid-ocean ridge.** The continuous, generally submarine, seismic, median mountain range that marks the divergent tectonic margin(s) in Earth's oceans.
- migmatite.** Literally, "mixed rock" with both igneous and metamorphic characteristics due to partial melting during metamorphism.
- mineral.** A naturally occurring, inorganic crystalline solid with a definite chemical composition or compositional range.
- monocline.** A one-limbed flexure in strata, which are usually flat-lying except in the flexure itself.
- monzonite.** A group of plutonic rocks containing approximately equal amounts of alkali feldspar and plagioclase, little or no quartz, and commonly augite as the main mafic mineral. Intrusive equivalent of latite.
- moraine.** A mound, ridge, or other distinct accumulation of unsorted, unstratified glacial drift, predominantly till, deposited chiefly by direct action of glacial ice.
- mole track.** A small, geologically short-lived ridge, formed by the humping up and cracking of the ground where movement along a large strike-slip fault occurred in heavily alluviated terrain.
- moulin.** A roughly cylindrical, nearly vertical hole or shaft in the ice of a glacier, scoured out by swirling meltwater as it pours down from the surface.
- mylonite.** A fine-grained, foliated rock typically found in localized zones of ductile deformation, often formed at great depths under high temperature and pressure.
- nappe.** A sheetlike, allochthonous (manufactured elsewhere) rock unit that has moved in a predominantly horizontal surface. The mechanism may be thrust faulting, recumbent folding, or gravity sliding.
- nonconformity.** An erosional surface preserved in strata in which crystalline igneous or metamorphic rocks underlie sedimentary rocks.
- normal fault.** A dip-slip fault in which the hanging wall moves down relative to the footwall.
- obduction.** The process by which the crust is thickened by thrust faulting at a convergent margin.
- oceanic crust.** Earth's crust formed at spreading ridges that underlies the ocean basins. Oceanic crust is 6 to 7 km (3 to 4 mi) thick and generally of basaltic composition.
- orogeny.** A mountain-building event.
- outcrop.** Any part of a rock mass or formation that is exposed or "crops out" at Earth's surface.
- outwash.** Glacial sediment transported and deposited by meltwater streams.
- overbank deposits.** Alluvium deposited outside a stream channel during flooding.
- overburden.** Non-economic, often unconsolidated, rock and sediment overlying an ore, fuel, or sedimentary deposit.
- paleogeography.** The study, description, and reconstruction of the physical landscape from past geologic periods.
- paleontology.** The study of the life and chronology of Earth's geologic past based on the fossil record.
- Pangaea.** A theoretical, single supercontinent that existed during the Permian and Triassic periods (also see Laurasia and Gondwana).
- parent (rock).** The original rock from which a metamorphic rock was formed. Can also refer to the rock from which a soil was formed.
- passive margin.** A margin where no plate-scale tectonism is taking place; plates are not converging, diverging, or sliding past one another. An example is the east coast of North America. (also see "active margin").
- pebble.** Generally, small rounded rock particles from 4 to 64 mm (0.16 to 2.52 in) in diameter.
- periglacial.** Describes processes, climates, and features at the margin of former or existing glaciers or icesheets influenced by the cold temperatures of the ice.
- permeability.** A measure of the relative ease with which fluids move through the pore spaces of rocks or sediments.
- phenocryst.** A coarse crystal in a porphyritic igneous rock.
- placer.** A surficial mineral deposit formed by mechanical concentration of mineral particles from weathered debris. The mineral concentrated is usually a heavy mineral such as gold, cassiterite, or rutile.
- plate tectonics.** The concept that the lithosphere is broken up into a series of rigid plates that move over Earth's surface above a more fluid asthenosphere.
- pluton.** A body of intrusive igneous rock that crystallized at some depth beneath Earth's surface.
- plutonic.** Describes igneous rock intruded and crystallized at some depth beneath Earth's surface.
- porosity.** The proportion of void space (e.g., pores or voids) in a volume of rock or sediment deposit.
- porphyry.** An igneous rock consisting of abundant coarse crystals in a fine-grained matrix.
- potassium feldspar.** An alkali feldspar rich in potassium (e.g., orthoclase, microcline, sanidine, adularia).
- Principal of Original Horizontality.** The concept that sediments are originally deposited in horizontal layers and that deviations from the horizontal indicate post-depositional deformation.
- Principle of Superposition.** The concept that sediments are deposited in layers, one atop another, i.e., the rocks on the bottom are oldest with the overlying rocks progressively younger toward the top.
- progradation.** The seaward building of land area due to sedimentary deposition.
- protolith.** The parent or unweathered and/or unmetamorphosed rock from which regolith or metamorphosed rock is formed.
- provenance.** A place of origin. The area from which the constituent materials of a sedimentary rock were derived.
- radioactivity.** The spontaneous decay or breakdown of unstable atomic nuclei.
- radiometric age.** An age in years determined from radioisotopes and their decay products.
- red beds.** Sedimentary strata composed largely of sandstone, siltstone, and shale that are predominantly red due to the presence of ferric oxide (hematite) coating individual grains.
- regolith.** General term for the layer of rock debris, organic matter, and soil that commonly forms the land surface and overlies most bedrock.
- regression.** A long-term seaward retreat of the shoreline or relative fall of sea level.

relative dating. Determining the chronological placement of rocks, events, or fossils with respect to the geologic time scale and without reference to their numerical age.

reverse fault. A contractional high-angle (greater than 45°) dip-slip fault in which the hanging wall moves up relative to the footwall (also see “thrust fault”).

rhyolite. A group of igneous rocks, typically porphyritic and commonly exhibiting flow texture, with phenocrysts of quartz and alkali feldspar in a glassy to cryptocrystalline groundmass. The fine-grained extrusive equivalent of granite.

rift valley. A depression formed by grabens along the crest of an oceanic spreading ridge or in a continental rift zone.

rock. A solid, cohesive aggregate of one or more minerals.

roundness. The relative amount of curvature of the “corners” of a sediment grain, especially with respect to the maximum radius of curvature of the particle.

sand. A clastic particle smaller than a granule and larger than a silt grain, having a diameter in the range of 1/16 mm (0.0025 in) to 2 mm (0.08 in).

sandstone. Clastic sedimentary rock of predominantly sand-sized grains.

scarp. A steep cliff or topographic step resulting from displacement on a fault, or by mass movement, or erosion. Also called an “escarpment.”

seafloor spreading. The process in which tectonic plates diverge and new lithosphere is created at oceanic ridges.

seamount. An elevated portion of the sea floor, 1,000 m (3,300 ft) or higher, either flat-topped or peaked.

sediment. An eroded and deposited, unconsolidated accumulation of rock and mineral fragments.

sedimentary rock. A consolidated and lithified rock consisting of clastic and/or chemical sediment(s). One of the three main classes of rocks—igneous, metamorphic, and sedimentary.

sequence. An informal rock-stratigraphic unit that is traceable over large areas and defined by a major sea level transgression-regression sediment package.

serpentinite. A rock consisting almost wholly of serpentine-group minerals, e.g. antigorite and chrysotile, commonly derived from the alteration of peridotite. Accessory chlorite, talc, and magnetite may be present.

shale. A clastic sedimentary rock made of clay-sized particles that exhibit parallel splitting properties.

sill. A tabular, igneous intrusion that is concordant with the surrounding rock.

silt. Clastic sedimentary material intermediate in size between fine-grained sand and coarse clay (1/256 to 1/16 mm [0.00015 to 0.002 in]).

siltstone. A variable-lithified sedimentary rock with silt-sized grains.

slickenside. A smoothly polished and often striated surface representing deformation of a fault plane.

slope. The inclined surface of any geomorphic feature or measurement thereof. Synonymous with gradient.

slump. A generally large, coherent mass movement with a concave-up failure surface and subsequent backward rotation relative to the slope.

soil. Surface accumulation of weathered rock and organic matter capable of supporting plant growth and often overlying the parent material from which it formed.

soliflucation. The slow downslope movement of waterlogged soil, normally at 0.5–5.0 cm/year (0.2–2 in/year), especially the flow occurring at high elevations in regions underlain by frozen ground that acts as a downward barrier to water percolation, initiated by frost action and augmented by meltwater resulting from alternate freezing and thawing of snow and ground ice.

spring. A site where water issues from the surface due to the intersection of the water table with the ground surface.

strata. Tabular or sheet-like masses or distinct layers of rock.

stratigraphy. The geologic study of the origin, occurrence, distribution, classification, correlation, and age of rock layers, especially sedimentary rocks.

stream. Any body of water moving under gravity flow in a clearly confined channel.

strike. The compass direction of the line of intersection of an inclined surface with a horizontal plane.

strike-slip fault. A fault with measurable offset where the relative movement is parallel to the strike of the fault. Said to be “sinistral” (left-lateral) if relative motion of the block opposite the observer appears to be to the left. “Dextral” (right-lateral) describes relative motion to the right.

subduction zone. A convergent plate boundary where oceanic lithosphere descends beneath a continental or oceanic plate and is carried down into the mantle.

subbituminous. A black coal, intermediate in rank between lignite and bituminous coal. It is distinguished from lignite by higher carbon and lower moisture content.

subsidence. The gradual sinking or depression of part of Earth’s surface.

suture. The linear zone where two continental landmasses become joined due to obduction.

system (stratigraphy). The group of rocks formed during a period of geologic time.

tectonic. Relating to large-scale movement and deformation of Earth’s crust.

terraces (stream). Step-like benches surrounding the present floodplain of a stream due to dissection of previous flood plain(s), stream bed(s), and/or valley floor(s).

terrane. A large region or group of rocks with similar geology, age, or structural style.

terrestrial. Relating to land, Earth, or its inhabitants.

terrigenous. Derived from the land or a continent.

theory. A hypothesis that has been rigorously tested against further observations or experiments to become a generally-accepted tenet of science.

thermokarst topography. An irregular land surface containing cave-in lakes, bogs, caverns, pits, and other small depressions formed in a permafrost region by the melting of ground ice.

thrust fault. A contractional dip-slip fault with a shallowly dipping fault surface (less than 45°) where

the hanging wall moves up and over relative to the footwall.

till. Unstratified drift, deposited directly by a glacier without reworking by meltwater, and consisting of a mixture of clay, silt, sand, gravel, and boulders ranging widely in size and shape.

tonalite. A plutonic rock with the composition of diorite, but with an appreciable amount of quartz (between 5 and 20 percent of the light-colored constituents).

topography. The general morphology of Earth's surface, including relief and locations of natural and anthropogenic features.

trace (fault). The exposed intersection of a fault with Earth's surface.

trace fossils. Sedimentary structures, such as tracks, trails, burrows, coprolites (dung), etc., that preserve evidence of organisms' life activities, rather than the organisms themselves.

transgression. Landward migration of the sea as a result of a relative rise in sea level.

travertine. A limestone deposit or crust, often banded, formed from precipitation of calcium carbonate from

saturated waters, especially near hot springs and in caves.

trend. The direction or azimuth of elongation or a linear geological feature.

tuff. Generally fine-grained, igneous rock formed of consolidated volcanic ash.

unconformity. An erosional or non-depositional surface bounded on one or both sides by sedimentary strata that marks a period of missing time.

uplift. A structurally high area in the crust, produced by movement that raises the rocks.

vent. An opening at Earth's surface where volcanic materials emerge.

volcanic. Related to volcanoes. Igneous rock crystallized at or near the Earth's surface (e.g., lava).

water table. The upper surface of the saturated zone; the zone of rock in an aquifer saturated with water.

weathering. The set of physical, chemical, and biological processes by which rock is broken down.

xenolith. A foreign rock entrained in magma as an inclusion.

Literature Cited

This section lists references cited in this report. A more complete geologic bibliography is available from the National Park Service Geologic Resources Division.

- Adamczak, S., Jr., R. D. Abbott, and F. J. Wuttig. 2004. "Earthquake Hazards: Geotechnical Design Concerns. A Review Based on the November 3, 2002 Alaska Earthquake." In *It's a Cool World*, eds. Daniel W. Smith, David C. Sego, and Charles A. Lenzion. Proceedings of the International Symposium on Cold Regions Engineering 12: unpaginated.
- Adams, P. C. 1999. "Fluvial Geomorphic Processes and Succession on the Tanana River Floodplain." In *Arctic Science Conference Program and Abstracts* 50: 161.
- Adema, G. W., P. Brease, and A. Bucki. 2003. "Glacier Monitoring in Denali National Park and Preserve." In *Geological Society of America Abstracts With Programs* 35 (6): 132.
- Alaska Department of Environmental Conservation (ADEC). 2006. *Contaminated Sites Program – Database Results*. Division of Spill Prevention and Response, Contaminated Sites Program <http://www.dec.state.ak.us/SPAR/csp/> (accessed August 30, 2006).
- Aochi, H., O. Scotti, and C. Berge-Thierry. 2005. "Dynamic Transfer of Rupture Across Differently Oriented Segments in a Complex 3-D Fault System." In *Geophysical Research Letters* 32 (21): 4 pp.
- Axford, Y. L. and A. Werner. 1997. "A Lacustrine Record of Late-Glacial/Holocene Environmental Change: Pork Chop Pond, Denali National Park, AK." In *Geological Society of America Abstracts With Programs* 29 (6): 35, 1997.
- Braile, L. W. 2009. Seismic monitoring. In *Geological Monitoring*, eds. R. Young and L. Norby, 229-244, Boulder, CO: Geological Society of America.
- Brease, P. 2004. "Denali National Park and Preserve: South Central Alaska." In *Geology of National Parks*, eds. A. G. Harris, E. Tuttle, and S. D. Tuttle: 477-504.
- Brease, P., L. Stromquist, A. Fiorillo, and S. T. Hasiotis. 2009. "Cretaceous Dinosaurs in Denali—A Newly Discovered Resource Requires a New Management Plan." In *Geological Society of America Abstracts With Programs* 41 (7): 150.
- Bucki, A. K. and V.B. Valentine. 2003. "Thickness Changes of the Muldrow Glacier Near the Traleika Confluence, Denali National Park, Alaska." In *Arctic Science Conference Program and Abstracts* 54: 199.
- Bundtzen, T. K. 1981. Geology and Mineral Deposits of the Kantishna Hills, Mt. McKinley Quadrangle, Alaska. MS Thesis. Anchorage, AK: University of Alaska.
- Burghardt, J. E. 1997. Mineral Report: Validity Examination of the Comstock #1 Lode, Comstock #2 Lode, Comstock #5 Lode, and Comstock #6 Lode Mining Claims. Unpublished report. Denver, CO: Geologic Resources Division, National Park Service.
- Capps, S. R. 1933. *The Eastern Portion of Mount McKinley National Park [Alaska]*. Bulletin B 0836-D: 219-300. Reston, VA: U. S. Geological Survey.
- Carver, G., G. Plafker, M. Metz, L. Cluff, B. Slemmons, E. Johnson, J. Roddick, and S. Sorensen. 2004. "Surface Rupture on the Denali Fault Interpreted From Tree Damage During the 1912 Delta River Mw 7.2-7.4 Earthquake: Implications for the 2002 Denali Fault Earthquake Slip Distribution." In *Bulletin of the Seismological Society of America* 94 (6B): S58-S71.
- Cassidy, J. F. and G. C. Rogers. 2004. "The M (sub w) 7.9 Denali Fault Earthquake of 3 November 2002: Felt Reports and Unusual Effects across Western Canada." In *Bulletin of the Seismological Society of America* 94 (6): Part B 53-57.
- Clague, J. J. 1979. "The Denali Fault System in Southwest Yukon Territory: A Geologic Hazard?" In *Paper - Geological Survey of Canada* (79-1A): 169-178.
- Clark, M. H. and M. S. Duffy. 2003. *Soil survey of Denali National Park area, Alaska*. Palmer, AK: Natural Resource Conservation Service. <http://soildatamart.nrcs.usda.gov/Manuscripts/AK651/0/DenaliPark.pdf>. Accessed 10 September 2010.
- Clautice, K. H. and R. J. Newberry. 1999. "Geology and Geophysics of the Healy A-6 Quadrangle, Chulitna Mining District." In *Arctic Science Conference Program and Abstracts* 50: 207-208.
- Clautice, K. H., R. J. Newberry, R. B. Blodgett, T. K. Bundtzen, B. G. Gage, E. E. Harris, S. A. Liss, M. L. Miller, R. R. Reifensstuhl, and D. S. Pinney. 2001. Bedrock Geologic Map of the Chulitna Region, South-Central Alaska. RI2001-1A, 1C, 1D. State of Alaska Department of Natural Resources: Division of Geological and Geophysical Surveys.
- Cole, R. B. 2004. "Dynamics of Tertiary Volcanism in Denali National Park, Alaska." In *Geological Society of America Abstracts With Programs* 36 (5): 54-55.

- Cole, R. B., K. D. Ridgway, P. W. Layer, and J. Drake. 1996. "Volcanic History, Geochronology, and Deformation of the Upper Cantwell Formation, Denali National Park, Alaska: Early Eocene Transition Between Terrane Accretion and Strike-Slip Tectonics." In *Geological Society of America Abstracts With Programs* 28 (7): 313.
- Cole, R. B., W. M. Brewer, and B. P. Hooks. 2002. "Melange and Late Paleozoic-Triassic(?) Chert, Limestone, and Pillow Basalt Along a Terrane Suture Zone South of the Denali Fault, Reindeer Hills, Alaska." In *Geological Society of America Abstracts With Programs* 34 (6): 44.
- Condie, K. C. and R.E. Sloan. 1998. *Origin and Evolution of the Earth, Principles of Historical Geology*. Dubuque, IA: Prentice-Hall, Inc.
- Coney, P. J., N. J. Silberling, and D.L. Jones. 1980. "Accretionary Tectonic Styles in the Alaska Range." In *Eos, Transactions, American Geophysical Union* 61 (46): 1114.
- Csejtey, B., Jr., D. P. Cox, R. C. Evarts, G. D. Stricker, and H. L. Foster. 1982. "The Cenozoic Denali Fault System and the Cretaceous Accretionary Development of Southern Alaska." In *Journal of Geophysical Research*. B 87 (5): 3741-3754.
- Csejtey, B., Jr., A. B. Ford, C. T. Wrucke, J. T. Dutro, Jr., A. G. Harris, and P. F. Brease. 1995. "Geological Correlations across the Denali Fault in South-Central Alaska: Implications for Cenozoic Fault Displacement." In *Geological Society of America Abstracts With Programs* 27 (5): 12-13.
- Denali National Park and Preserve. 2009. *Denali National Park and Preserve Resource Stewardship Strategy 2008-2027 Summary*. <http://www.nps.gov/denali/naturescience/rss.htm>. Denali Park, AK: Denali National Park and Preserve. Accessed December 2009.
- Densmore, R. V. 2005. "Succession on Subalpine Placer Mine Spoil: Effects of Revegetation with *Alnus Viridis*, Alaska, U.S.A." In *Arctic, Antarctic, and Alpine Research* 37 (3): 297-303.
- Dumoulin, J. A., D. C. Bradley, and A. G. Harris. 1998. *Sedimentology, Conodonts, Structure, and Regional Correlation of Silurian and Devonian Metasedimentary Rocks in Denali National Park, Alaska*. U. S. Geological Survey Professional Paper P 1595: 71-98.
- Dusel-Bacon, C., B. Csejtey, Jr., H. L. Foster, E. O. Doyle, W. J. Nokleberg, and G. Plafker. 1993. *Distribution, Facies, Ages, and Proposed Tectonic Associations of Regionally Metamorphosed Rocks in East and South-Central Alaska*. U. S. Geological Survey Professional Paper P 1497-C: C1-C72.
- Dyson, J. L. 1966. *Glaciers and Glaciation in Glacier National Park*. Glacier Natural History Association, in cooperation with the National Park Service.
- Eberhart-Phillips, D., P. J. Haeussler, J. T. Freymueller, A. D. Frankel, C. M. Rubin, P. Crow, N. A. Ratchkovski, G. Anderson, G. A. Carver, A. J. Crone, T. E. Dawson, H. Fletcher, R. Hansen, E. L. Harp, R. A. Harris, D. P. Hill, S. Hreinsdóttir, R. W. Jibson, L. M. Jones, R. Kayen, D. K. Keefer, C. F. Larsen, S. C. Moran, S. F. Personius, G. Plafker, B. Sherrod, K. Sieh, N. Sitar, and W. K. Wallace. 2003. "The 2002 Denali Fault Earthquake, Alaska: A Large Magnitude, Slip-Partitioned Event." In *Science* 300 (5622): 1113-1118.
- Edwards, P., M. Tranel, P. Brease, and P. Sousanes. 2000. "Stream and River Water Quality in Denali National Park and Preserve, Alaska." In *American Water Resources Association Technical Publication Series* TPS 00-1: 203-207.
- Elias, S. A., S. K. Short, and C. F. Waythomas. 1996. "Late Quaternary Environments, Denali National Park and Preserve, Alaska." In *Arctic* 49 (3): 292-305.
- Eppinger, R. G., P. H. Briggs, J. G. Crock, A. L. Meier, S. J. Sutley, and P. M. Theodorakos. 2002. *Environmental-Geochemical Study of the Slate Creek Antimony Deposit, Kantishna Hills, Denali National Park and Preserve, Alaska*. Professional Paper P 1662: 123-141. Reston, VA: U. S. Geological Survey.
- Fitzgerald, P. G., E. Stump, and T. F. Redfield. 1993. "Late Cenozoic Uplift of Denali and its Relation to Relative Plate Motion and Fault Morphology." In *Science* 259 (5094): 497-499.
- Ford, A. B., B. Csejtey, Jr., and C. T. Wrucke. 2003. "Denali Fault Geologic Relations at Gunsight Pass, Denali National Park, Alaska." In *Geological Society of America Abstracts With Programs* 35 (6): 561.
- Gangloff, R. A. 1999. "Arctic Dinosaurs and Their Cretaceous Record in Alaska." In *Arctic Science Conference Program and Abstracts* 50: 288-289.
- Garcia, R., F. Crespon, V. Ducic, and P. Lognonne. 2005. "Three-Dimensional Ionospheric Tomography of Post-Seismic Perturbations Produced by the Denali Earthquake from GPS Data." In *Geophysical Journal International* 163 (3): 1049-1064.
- Gilbert, W. G. 1979. *A Geologic Guide to Mount McKinley National Park*. Anchorage, AK: Alaska Natural History Association.
- Goldfarb, R. J., L. D. Miller, D. L. Leach, and L. W. Snee. 1997. "Gold Deposits in Metamorphic Rocks of Alaska." In *Economic Geology Monographs* 9: 151-190.

- Gough, L. P. and J. G. Crock. 1997. Distinguishing Between Natural Geologic and Anthropogenic Trace Element Sources, Denali National Park and Preserve. Professional Paper P 1574: 54-71. Reston, VA: U. S. Geological Survey.
- Greenwood, S. M. and G. Rowell. 1980. "Mount McKinley." In *America's Magnificent Mountains*, pref. Sterling B. Hendricks, 90-103, Washington, DC: National Geographic Society Special Publications Division.
- Haeussler, P. J. 2005. "What made Denali so Tall? Structural Geology of the High Peaks of the Alaska Range." In *Geological Society of America Abstracts With Programs* 37 (7): 79.
- Harp, E. L., R. W. Jibson, R. E. Kayen, D. K. Keffer, B. L. Sherrod, G. A. Carver, B. D. Collins, R. E. S. Moss, and R. N. Sitar. 2003. "Landslides and Liquefaction Triggered by the M 7.9 Denali Fault Earthquake of 3 November 2002." In *GSA Today* 13 (8): 4-10.
- Hickman, R. G., K. W. Sherwood, and C. Craddock. 1990. "Structural Evolution of the Early Tertiary Cantwell Basin, South-Central Alaska." In *Tectonics* 9(6): 1433-1449.
- IPCC, 2007, Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change: Geneva, Switzerland, Intergovernmental Panel on Climate Change, 104 p. (http://www.ipcc.ch/publications_and_data/ar4/syr/en/contents.html, accessed March 6, 2010.)
- Jones, D.L., N. J. Silberling, W. G. Gilbert, and P. J. Coney. 1980. "Age, Character, and Distribution of Accreted Terranes in the Central Alaska Range, South-Central Alaska." In *Eos, Transactions, American Geophysical Union* 61 (46): 1114.
- Karl, T. R., Melillo, J. M., and Peterson, T. C. 2009. *Global climate change impacts in the United States*. New York, NY: Cambridge University Press. <http://www.globalchange.gov/publications/reports/scientific-assessments/us-impacts>. Accessed 17 September 2010.
- Karpilo, R. D., Jr. 2009. Glacier monitoring techniques. In *Geological Monitoring*, eds. R. Young and L. Norby, 141-162, Boulder, CO: Geological Society of America.
- Karpilo, R. D., Jr., G. W. Adema, and B. F. Molina. 2004. "Documenting Glacier Change in Denali National Park and Preserve, Alaska." In *Geological Society of America Abstracts With Programs* 36 (5): 229.
- Lillie, R. J. 2005. *Parks and Plates: The Geology of our National Parks, Monuments, and Seashores*. New York: W.W. Norton and Company.
- Lowey, G. W. 1998. "A New Estimate of the Amount of Displacement on the Denali Fault System Based on the Occurrence of Carbonate Megaboulders in the Dezadeash Formation (Jura-Cretaceous), Yukon, and the Nutzotin Mountains Sequence (Jura-Cretaceous), Alaska." In *Bulletin of Canadian Petroleum Geology* 94: 4333-4359.
- Lord, M. L., D. Germanoski, and N. E. Allmendinger. Fluvial geomorphology: Monitoring stream systems in response to a changing environment. In *Geological Monitoring*, eds. R. Young and L. Norby, 69-103, Boulder, CO: Geological Society of America.
- Mann, D. H. 1999. "Development of Taiga Vegetation and Soils over Millennial Time Scales in Denali National Park." In *Arctic Science Conference Program and Abstracts* 50: 229.
- Matmon, A., D. Schwartz, H. Stenner, J. Lienkamper, T. Dawson, P. J. Haeussler, L. Staft, and R. C. Finkel. 2004. Determining Holocene and Late Pleistocene Slip Rates Along the Denali Fault Using Cosmogenic (Super 10) Be Analysis of Boulders on Displaced Moraines." In *Geological Society of America Abstracts With Programs* 36 (5): 137.
- Matmon, A., D. P. Schwartz, P. J. Haeussler, R. Finkel, J. J. Lienkamper, H. D. Stenner, and T. E. Dawson. 2006. "Denali Fault Slip Rates and Holocene-Late Pleistocene Kinematics of Central Alaska." In *Geology* 34: 645-644.
- McCauley, C. A., M. R. Lilly, D. M. Nyman, and D. M. White. 1999. "Measuring the Potential of Permafrost as an Environmental Barrier by a Diesel/Jet Fuel In-situ Infiltration Test in Bethel, Alaska." In *Arctic Science Conference Program and Abstracts* 50: 275-276.
- Meisling, K. E., M. C. Gardner, G. W. Cushing, and S. C. Bergman. 1987. "A Reconstruction of Southern Alaska at 75 Ma." In *Geological Society of America Abstracts With Programs* 19 (7): 769.
- Metz, P. A., C. J. Freeman, D. D. Adams, and J. C. Balla. 1989. "Late Proterozoic Precious Metal Bearing Sequences in Interior Alaska." In *Geological Society of America Abstracts With Programs* 21 (5): 116-117.
- Miller, M. L., D. C. Bradley, T. K. Bundtzen, and W. C. McClelland. 2002. "Late Cretaceous through Cenozoic Strike-Slip Tectonics of Southwestern Alaska." In *Journal of Geology* 110 (3): 247-270.
- Miller, T. P. 1994. "Pre-Cenozoic Plutonic Rocks in Mainland Alaska." In *The Geology of Alaska*, eds. G. Plafker and H. C. Berg, 535-554 Boulder, CO: Geological Society of America.
- Molina, B. F. and M. Sfraga. 1999. "Measuring and Monitoring Changes in Alaska's Glaciers with Ground, Aerial, and Space Photography: A History." In *Arctic Science Conference Program and Abstracts* 50: 78-79.

- Moore, T. T., A. Grantz, and S. M. Roeske. 1994. "Continent-Ocean Transition in Alaska: The Tectonic Assemblage of Eastern Denalia." In *Phanerozoic Evolution of North America Continental-Ocean Transition*, ed. R. C. Speed. Geological Society of America, DNAG Continental-Ocean Transect Volume: 399-441.
- National Park Service. 2006. Soil Survey Geographic (SSURGO) database for Denali National Park and Preserve, Alaska. Denver, CO: National Park Service Geologic Resources Division.
<http://science.nature.nps.gov/nrdata/datastore.cfm?ID=48854>. Accessed 10 September 2010.
- National Park Service. 2009. Glaciers/Glacial Features. Online feature. Denali Park, AK: Denali National Park and Preserve. <http://www.nps.gov/denali/naturescience/glaciers.htm>. Accessed 16 September 2010.
- Newkirk, S. R. 2005. "Tectonic Setting and Lithogeochemical Characteristics of Metavolcanics and Metabasite Sills of the Delta District, East-Central Alaska Range, Alaska." In *Geological Society of America Abstracts With Programs* 37 (2): 35.
- Nixon, G. T. 2003. "Use of Spinel in Mineral Exploration: The Enigmatic Giant Mascot Ni-Cu-PGE Deposit - Possible Ties to Wrangellia and Metallogenic Significance." In *Geological Fieldwork 2002: A Summary of Field Activities and Current Research*. Geological Fieldwork, Report: 2003-1: 115-128. Victoria, BC: British Columbia Ministry of Energy and Mines, Resource Development Division, Geological Survey Branch.
- Nokleberg, W. J., G. Plafker, and F. H. Wilson. 1994. "Geology of South-Central Alaska." In *The Geology of Alaska*, eds. George Plafker and Henry C. Berg., 311-366 Boulder, CO: Geological Society of America.
- Norris, F. 1998. "Gold Rush-Era Mining Sites in Alaska's National Parks." In *CRM (Washington, D.C.)* 21 (7): 30-31.
- Osterkamp, T. E. and M. T. Jorgenson. 2009. Permafrost conditions and processes. In *Geological Monitoring*, eds. R. Young and L. Norby, 205-227, Boulder, CO: Geological Society of America.
- Petrone, K. C., L. D. Hinzman, and R. D. Boone. 1999. "The Influence of Permafrost and Fire Disturbance on Nitrogen and Carbon Concentrations in Three Subarctic Streams." In *Arctic Science Conference Program and Abstracts* 50: 202.
- Pfeiffer, T. J., and S. L. Bednarz. 2000. "The Influence of Structural Geology on a Massive Landslide in Denali National Park, Alaska." In *Annual Meeting - Association of Engineering Geologists* 43 (4): 107.
- Plafker, G., W. J. Nokleberg, and J. S. Lull. 1989. "Bedrock Geology and Tectonic Evolution of the Wrangellia, Peninsular, and Chugach Terranes Along the Trans-Alaska Crustal Transect in the Chugach Mountains and Southern Copper River Basin, Alaska." In *Journal of Geophysical Research, B, Solid Earth and Planets* 94 (4): 4255-4295.
- Popovics, L. M., A. M. Milner, and C.L. Ping. 1999. "The Effect of Soil and Stream Water Quality on Primary and Secondary Productivity of Rock Creek, Denali National Park and Preserve, Alaska." In *Arctic Science Conference Program and Abstracts* 50: 194-195.
- Prindle, L. M. 1907. *The Bonnifield & Kantishna Regions, Alaska*. Bulletin 314-L. Reston, VA: U.S. Geological Survey.
- Ratchkovski, N. A., S. Wiemer, and R. A. Hansen. 2004. "Seismotectonics of the Central Denali Fault, Alaska, and the 2002 Denali Fault Earthquake Sequence." In *Bulletin of the Seismological Society of America* 94 (6): Part B 156-174.
- Redfield, T. F. and P. G. Fitzgerald. 1993. "Denali Fault System of Southern Alaska: An Interior Strike-Slip Structure Responding to Dextral and Sinistral Shear Coupling." In *Tectonics* 12 (5): 1195-1208.
- Redfield, T. F. and P. G. Fitzgerald. 2000. "Plate Kinematics, Escape Tectonics, and the Denali Fault System of Cenozoic South Central Alaska." In *Eos, Transactions, American Geophysical Union* 81 (48): Suppl. 1123.
- Redfield, T. F., D. W. Scholl, M. E. Beck, and P. G. Fitzgerald. 2005. "Escape Tectonics and Plate Kinematics in the South-Central Alaskan Orocline." In *Geological Society of America Abstracts With Programs* 37 (7): 79.
- Reed, B.L. and S. W. Nelson. 1977. *Geologic map of the Talkeetna Quadrangle, Alaska*. Miscellaneous Field Studies Map 870-A. Reston, VA: U.S. Geological Survey.
- Richter, D. H. and D. L. Jones. 1973. "Structure and Stratigraphy of Eastern Alaska Range, Alaska." In *Arctic Geology Memoir - American Association of Petroleum Geologists* 19: 408-420.
- Ridgway, K. D., J. M. Trop, W. J. Nokleberg, C. M. Davidson, and K. R. Eastham. 2002. "Mesozoic and Cenozoic Tectonics of the Eastern and Central Alaska Range: Progressive Basin Development and Deformation in a Suture Zone." In *Geological Society of America Bulletin* 114 (12): 1480-1504.
- Roeske, S. M., B. A. Hampton, K. D. Ridgway, and G. E. Gehrels. 2005. Cryptic Strike-Slip Faults and Dismembered Ophiolites: Potential Tracers of the Alaska Range Suture Zone." In *Geological Society of America Abstracts With Programs* 37 (4): 80.

- Roush, J., and P. Brease. 1997. Glacier Monitoring in Denali National Park and Preserve, Alaska: Integrating Field Study and Remote Sensing. Open-File Report 98-0031. Reston, VA: U.S. Geological Survey.
- Ruppert, N.A., K.D. Ridgway, J.T. Freymueller, R.S. Cross, and R.A. Hansen. 2008. Active tectonics of interior Alaska: Seismicity, GPS geodesy, and local geomorphology. In *Active Tectonics and Seismic Potential in Alaska*, eds Freymueller, J.T., et al., *Geophysical Monograph Series*, 179: 109-133.
- Salisbury and Dietz, Inc. 1984. *Kantishna Hills and Dunkle Mine Areas, Denali National Park and Preserve, Alaska*. 1983 Mineral Resource Studies, Contract S0134031. Washington, DC: U.S. Bureau of Mines.
- Santucci, V. L., J. P. Kenworthy and A. L. Mims. 2009. Monitoring in situ paleontological resources. In *Geological Monitoring*, eds. R. Young and L. Norby, 189-204, Boulder, CO: Geological Society of America.
- Schwartz, D. P., P. J. Haeussler, A. Matmon, T. E. Dawson, G. Seitz, H. Stenner, S. Bemis, E. Molhoek, B. Sherrod, A. J. Crone, S. Personius, P. A. Craw, and J. Beget. 2005. *Earthquake Geology of the Denali Fault System, Alaska*. Open-File Report 2005-1131: 25-29. Reston, VA: U. S. Geological Survey.
- Seraphim, R. H. 1975. "Denali: A Monmetamorphosed Stratiform Sulfide Deposit." In *Economic Geology and the Bulletin of the Society of Economic Geologists* 70 (5): 949-959.
- St. Amand, P. 1957. Geological and geophysical synthesis of the tectonics of portions of British Columbia, the Yukon Territory and Alaska. *Geological Society of America Bulletin* 68: 1343-1370.
- Stone, D. B.. 1980. "Timing of Accretion of Alaskan Tectonostratigraphic Terranes." In *Eos, Transactions, American Geophysical Union* 61 (46): 1114.
- Stone, D. B. 1984. "Alaska: Geology and Tectonics." In *Biennial Report - Geophysical Institute, University of Alaska* 1983-84: 1-19.
- Symons, D. T. A., M. J. Harris, J. E. Gabites, and C. J. R.Hart. 2000. "Eocene (51 Ma) End to Northward Translation of the Coast Plutonic Complex: Paleomagnetism and K-Ar Dating of the White Pass Dikes." In *Tectonophysics* 326 (1-2): 93-109.
- Thorson, R. M. 1986. "Late Cenozoic Glaciation of the Nenana Valley." In *Glaciation in Alaska-The Geologic Record*, eds. T. D. Hamilton, K. M. Reed, and R. M. Thorson: 99-122. Fairbanks, AK: Alaska Geological Society.
- Trop, J. M., K. D. Ridgway, and T. L. Spell. 2003. "Sedimentary Record of Transpressional Tectonics and Ridge Subduction in the Tertiary Matanuska Valley-Talkeetna Mountains Forearc Basin, Southern Alaska." In *Special Paper - Geological Society of America* 371: 89-118.
- Van Maanen, J. L. and G. L. Solin. 1988. Hydraulic and Channel Characteristics of Selected Streams in the Kantishna Hills Area, Denali National Park and Preserve, Alaska, 1982-84. U. S. Geological Survey Open-File Report 88-0325.
- Wahrhaftig, C. 1952. Geologic Map of Part of Healy C-4 Quadrangle, Alaska, Showing Pleistocene Deposits Along the Nenana River. U.S. Department of Interior: Geological Survey.
- Wahrhaftig, C. 1970. Geologic Map of the Healy D-4 Quadrangle, Alaska. Geologic Quadrangle Map GQ-806. Reston, VA: U.S. Geological Survey.
- Werner, A., R. M. Thorson, and N. W. Ten Brink. 1990. "Geology of the McKinley River Area, Denali National Park and Preserve, Alaska." In *Geological Society of America Abstracts With Programs* 22 (7): 176.
- West, A. W., M. Keskinen, and S. Swanson. 1995. "Petrology of the Mt. McKinley Pluton, Denali National Park, Alaska." In *Geological Society of America Abstracts With Programs* 27 (5): 83-84.
- White, J. C. 1997. "Deformation Processes Along the Denali Fault System." In *Lithoprobe Report* 56: 143-145.
- Wieczorek, G. F. and J. B. Snyder. 2009. Monitoring slope movements. In *Geological Monitoring*, eds. R. Young and L. Norby, 245-271, Boulder, CO: Geological Society of America.
- Wilson, F. H., J. H. Dover, D. C. Bradley, F. R. Weber, T. K. Bundtzen, and P. J. Hauessler (compilers). 1998. *Geologic Map of Central (Interior) Alaska*. Open-File Report 98-133. Reston, VA: U.S. Geological Survey.
- Zhang, T., R. G. Barry, and K. Knowles. 1999. "Characteristics of Permafrost and Ground Ice Distribution in the Northern Hemisphere." In *Arctic Science Conference Program and Abstracts* 50: 117-118.


Appendix A: Overview of Digital Geologic Data

The following page is an overview of the digital geologic data for Denali National Park and Preserve. For a poster-size PDF of this map and complete digital data, please see the included CD or visit the Geologic Resources Inventory publications web site (http://www.nature.nps.gov/geology/inventory/gre_publications.cfm).

The overview incorporates the small scale (1:250,000) data set that encompasses the entire park and preserve. Larger scale (1:63,360) data is available for Chulitna region and Healy quadrangle.





Overview of Digital Geologic Data for Denali National Park and Preserve





Overview of Digital Geologic Data for Denali National Park and Preserve



DENA - Geologic Units			
Water	Tegr - Granite and granodiorite	JTRsu - Chulitna sequence, red and brown sedimentary rocks and basalt	PDsc - Mystic sequence, Sheep Creek Formation and correlative siliciclastic units
Glacier	Tcv - Volcanic rocks of the Cantwell Formation	JTRct - Chulitna sequence, crystal tuff, argillite, chert, graywacke, and limestone	MDt - Totatlanika Schist
Qs - Surficial deposits, undifferentiated	Tpgr - Granitic rocks	JTRtv - Mystic sequence, Tatina River Volcanic and equivalent units	Dsb - Chulitna sequence, Serpentine, basalt, chert and gabbro
Tk - Kenai Group, undivided	TKvr - Rhyolite and related rocks	JCmd - Mystic and Dillinger stratigraphic sequences, undivided	D5mdl - Mystic and Dillinger sequence, unnamed limestone
Tsf - Kenai Group, Sterling Formation	TKvi - Andesite and related rocks	TRcs - Calcareous sedimentary rocks	D5wc - Nixon Fork sequence, Whirlwind Creek Formation and unnamed correlative units
Thd - Hornblende dacite	TKi - Intrusive rocks, undivided	TRcg - Conglomerate and volcanic sandstone	DSl - Limestone
Tn - Nenana Gravel	TKg - Granitic rocks	TRn - Nikolai Greenstone and related rocks	DCd - Dillinger sequence, undivided
Tsu - Sedimentary rocks, undivided	TKgd - Granodiorite, tonalite, and monzonite dikes, and stocks	TRnm - Metavolcanic and associated metasedimentary rocks	SCpl - Dillinger sequence, Post River Sandstone, Lyman Hills Formation, and correlative units
Tvu - Volcanic rocks, undivided	TKgb - Gabbro and leucogabbro	TRlb - Chulitna sequence, limestone and basalt sequence	Oc - Chert
Tty - Kenai Group, Tyonek Formation	TKgg - Gneissose granitic rocks	TRr - Chulitna sequence, red beds	CZw - Wicksham Grit, undivided
Tcb - Coal-bearing rocks	Kcs - Cantwell Formation, sedimentary rocks subunit	TRPNas - Flysch-like sedimentary rocks	PZk - Keevy Peak Formation
Thf - Hypabyssal felsic and intermediate intrusive rocks	Kmar - Melanges of the Alaska Range	TRDv - Chulitna sequence, volcanic and sedimentary rocks	PZsc - Spruce Creek sequence and correlative rocks
Thm - Hypabyssal mafic intrusive rocks	KJs - Argillite, chert, sandstone, and limestone	TRSI - Melanges of the Alaska Range, limestone blocks	PZvs - Volcanic and sedimentary rocks
Tvb - Andesite and basalt	KJf - Kahiltina flysch sequence, undivided	MZPZi - Intrusive and volcanic rocks, undivided	PZZaqs - Pelitic and quartzose schist of the Alaska Range
Tv - Granitic and volcanic rocks, undivided	KJfk - Kahiltina flysch sequence, flysch sequence 1	MZZum - Ultramafic and mafic rocks, undivided	UNKmlu - Melanges of the Alaska Range, ultramafic and associated rocks
Toem - Granodiorite to tonalite	KJfn - Kahiltina flysch sequence, flysch sequence 2	PPNasc - Skolai Group, Station Creek and Slana Spur Formations, and equivalent rocks	UNKbu - Bedrock of unknown type or age
Tfv - Fluvial sedimentary rocks and subordinate volcanic rocks	Jct - Chinitna Formation, Tuxedni Group, and coeval sedimentary rocks	PPNast - Skolai Group, Strelna metamorphic complex	
	Ji - Alaska-Aleutian Range and Chitina Valley batholiths, undifferentiated	PPNaskm - Skolai Group, marble	

Appendix B: Scoping Session Participants

The following is a list of participants from the GRI scoping session for Denali National Park and Preserve, held on February 24–26, 2004. The contact information and email addresses in this appendix may be outdated; please contact the Geologic Resources Division for current information. The scoping meeting summary was used as the foundation for this GRI report. The original scoping summary document is available on the GRI web site: http://www.nature.nps.gov/geology/inventory/gre_publications.cfm.

LAST NAME	FIRST NAME	AFFILIATION	TITLE	PHONE	E-MAIL
Adema	Guy	Denali National Park and Preserve	physical scientist	907-683-6356	guy_adema@nps.gov
Blodgett	Robert	U.S. Geological Survey	paleontologist	907-786-7416	rblodgett@usgs.gov
Brease	Phil	Denali National Park and Preserve	geologist		phil_brease@nps.gov
Claudice	Karen	Alaska Division of Geological and Geophysical Surveys	geologist	907-451-5023	karen@dnr.state.ak.us
Connors	Tim	NPS, Geologic Resources Division	geologist	303-969-2093	tim_connors@nps.gov
Dickison	George	NPS, Alaska Regional Office	GIS coordinator	907-644-3546	george_dickison@nps.gov
Fiorillo	Tony	Dallas Museum of Natural History	paleontologist	214-421-3466 x 234	tfiorillo@dmnhnet.org
Gamble	Bruce	U.S. Geological Survey	geologist	907-786-7479	bgamble@usgs.gov
Giffen	Bruce	NPS, Alaska Regional Office	geologist	907-644-3572	bruce_giffen@nps.gov
Halloran	Jim	NPS, Alaska Regional Office	geologist	907-644-3574	jim_halloran@nps.gov
Heise	Bruce	NPS, Geologic Resources Division	geologist	303-969-2017	bruce_heise@nps.gov
Kucinski	Russell	NPS, Alaska Regional Office	geologist	907-644-3571	russ_kucinski@nps.gov
Liebsher	Tom	Yukon-Charley Rivers National Preserve	chief, natural resources	907-455-0620	thomas_liebsher@nps.gov
Norby	Lisa	NPS, Geologic Resources Division	geologist	303-969-2318	lisa_norby@nps.gov
Rosenkrans	Danny	Wrangell-St. Elias National Park and Preserve	geologist	907-823-7240	danny_rosenkrans@nps.gov
Sharp	Devi	Wrangell-St. Elias National Park and Preserve	chief, natural resources	907-822-7212	devi_sharp@nps.gov
Spencer	Page	NPS, Alaska Regional Office	ecologist	907-644-3448	page_spencer@nps.gov
Stevens	Deanne	Alaska Division of Geological and Geophysical surveys	geologist	907-451-5014	deanne@dnr.state.ak.us
Stromquist	Linda	NPS, Alaska Regional Office	geologist	907-644-3576	linda_stromquist@nps.gov
Wesser	Sara	NPS, Central Alaska Network	network coordinator	907-644-3558	sara_wesser@nps.gov
Wilder	Doug	NPS, Central Alaska Network	data manager	907-455-0661	doug_wilder@nps.gov
Wilson	Ric	U.S. Geological Survey	geologist	907-786-7448	fwilson@usgs.gov

The Department of the Interior protects and manages the nation's natural resources and cultural heritage; provides scientific and other information about those resources; and honors its special responsibilities to American Indians, Alaska Natives, and affiliated Island Communities.

NPS 184/105592, September 2010

National Park Service
U.S. Department of the Interior



Natural Resource Program Center
1201 Oakridge Drive, Suite 150
Fort Collins, CO 80525

www.nature.nps.gov