





Common Native Trees



Tanoak

Lithocarpus densiflorus var. *densiflorus*

Leaves leathery, toothed.
Bark grayish brown.
Acorns hairy-capped and are important
Source of food for wildlife.



Red Alder

Alnus rubra


Grows in open and damp habitats.
Leaves coarsely toothed.
Bark white-gray.



California Laurel

(Pepperwood, Bay)
Umbellularia californica

Leaves shiny green, yellow in
fall, aromatic when rubbed in
hands.



Douglas-fir

Pseudotsuga menziesii var. *menziesii*


Found on ridges.
Bark brown and furrowed, brittle.
Often carries more
lichen than Coast Redwoods.



Big-leaf Maple

Acer macrophyllum

Grows along streams.
Bark covered with mosses.
Leaves yellow in fall.



Sitka Spruce

Picea sitchensis

Grows close to coast.
Needles grow on all sides of twig.
Bark gray, chip-like.



Coast Redwood

Sequoia sempervirens
(sun needles)

Fibrous reddish-brown bark, deep
furrows extend up tree. Cones
olive sized.



Coast Redwood

Sequoia sempervirens
(shade needles)

Scaly needles (left) from top of tree.
Flat needles (shown) from lower
reaches where light is
less prevalent.



Western Hemlock

(Octopus Tree)
Tsuga heterophylla

Often grows on
decaying logs and stumps.
Drooping branches with soft
needles.