

THE ECOLOGICAL ROLE OF ELK IN ROCKY MOUNTAIN NATIONAL PARK

FINAL REPORT

JUNE 2006



Ryan J. Monello¹, Therese L. Johnson¹, and R. G. Wright²

¹National Park Service, Rocky Mountain National Park, Estes Park, CO 80517

²Department of Fish and Wildlife Resources, University of Idaho, Moscow, ID 83844

TABLE OF CONTENTS

Introduction	1
Physical and Biological Characteristics of the Park	2
Human History of the Region	7
Summary	9
The Elk Population: Historical Information and Contemporary Research	10
Introduction – Elk Population Indices and Models.....	10
Ecological Carrying Capacity (K).....	10
Ecosystem Simulation Model (SAVANNA)	10
Statistical Differences	11
Pre-Settlement Conditions	11
Early Management (1915 – 1968).....	12
Natural Regulation	17
Science-Based Management	18
Elk Research (1968 – present)	19
Elk Distribution and Movements	19
Elk Population Dynamics.....	23
Elk Densities	26
Elk Survival	27
Chronic Wasting Disease.....	27
Summary	28
Focal Plant Communities	28
Aspen	28
Aspen Ecology	28
Aspen in RMNP	29
Willow	32
Willow Ecology	32
Willow in RMNP	32
Willow Distribution in RMNP.....	34
Willow Growth and Size in RMNP	35

TABLE OF CONTENTS

Willow Reproduction in RMNP	36
Bitterbrush and Sagebrush	37
Bitterbrush and Sagebrush Ecology	37
Bitterbrush and Sagebrush in RMNP.....	38
Herbaceous Communities on the Winter Range	38
Coniferous Forest Communities in RMNP.....	40
Alpine Tundra	41
Summary	41
Ungulate Interactions	43
Mule Deer	43
Competition with Elk.....	45
Chronic Wasting Disease.....	45
Bighorn Sheep.....	46
Competition with Elk.....	46
Moose.....	47
Summary	47
Effects of Elk on Biodiversity.....	48
Birds.....	48
Ptarmigan	48
Songbirds	49
Butterflies.....	50
Beaver	50
Beaver Ecology	50
Beaver in RMNP.....	51
Summary	52
Nutrient Cycles	52
Conclusions.....	53
Literature Cited	56

LIST OF FIGURES

Figure 1.	Land ownership adjacent to and near Rocky Mountain National Park.	3
Figure 2.	Elevation (in feet) and major vegetation covertypes of Rocky Mountain National Park.	4
Figure 3.	Elk ranges in Rocky Mountain National Park and vicinity.	5
Figure 4.	Approximate number of elk inhabiting the Estes Valley during winter from 1913 (initial year of reintroduction) To 2005.	13
Figure 5.	Approximate deer population size in the Estes Valley during winter from 1915 to 2003.	15
Figure 6.	Elk subpopulation boundaries during winter on the east side of Rocky Mountain National Park and vicinity.	21
Figure 7.	Elk migration routes and major stream drainages in the Rocky Mountain National Park area.	22
Figure 8.	Elk population dynamics (five-year moving average) in Rocky Mountain National Park and the Town of Estes Park.	24
Figure 9.	Woody shrub distribution on the elk winter range in Rocky Mountain National Park.	33
Figure 10.	Primary areas of use for moose, bighorn sheep, and mule deer in Rocky Mountain National Park.	44

APPENDIX

Appendix 1.	Common plant species found in Rocky Mountain National Park covertypes.	72
-------------	--	----

INTRODUCTION

Many national parks and equivalent protected areas in North America have experienced substantial increases in the numbers of several species of large grazing and browsing herbivores in the past several decades - numbers that are now considered by many managers and scientists to exceed levels that have existed at any time since the arrival of European colonists (Wright 1992). There is a great deal of scientific debate over the extent of this increase, in part because there is still considerable argument on how to obtain accurate counts of existing herbivore populations let alone how to ascertain the validity of historic counts (Buckland et al. 1993). The reasons for the increases in species such as Rocky Mountain elk (*Cervus elaphus*), white-tailed deer (*Odocoileus virginianus*), and to a lesser extent bison (*Bison bison*) are varied but largely come down to two causes; the almost complete elimination of a major herbivore predator the gray wolf (*Canis lupus*) and the prohibition of human harvest of wildlife in protected areas. Concerns about increased numbers of large herbivores have primarily evolved around the potential adverse effects that the high densities of these animals could be having on native vegetation, other animal species, and ecosystem properties in general. We refer to densities of animals perceived to be at numbers or densities that may be detrimental to the components of the natural system as overabundant, recognizing, as McShea et al. (1997) point out, that this concept is a value judgement having a clear meaning only when used in a specific context.

Caughley (1981) discussed four situations where the term overabundance could be applied to animal populations and all have been used in North America to support the argument that populations of certain herbivore species were overabundant. These situations are: 1) when animals threaten human life and livelihood; 2) when the animals depress the densities of favored species; 3) when the animals are considered to be too numerous for their own good; and 4) when their high numbers are deemed to cause ecosystem dysfunction. For many years, the National Park Service (NPS) culled large ungulates, primarily antelope (*Antilocapra americana*), bison, elk, and mule deer (*Odocoileus hemionus*) in several western parks and more recently white-tailed deer in an eastern area, largely because of concerns associated with reason number three (Robish and Wright 1995, Frost et al. 1997). In general, the typical stated concern was that animals would destroy their food supply and starve. In reality there was also a concern that visitors would see starving or dying animals and react negatively (Wright 1992). In recent years the scientists within and working with the NPS have begun to appreciate Caughley's fourth point with particular attention given to the decline of ecologically important vegetation communities such as aspen and willow in the case of elk use, and rare plants and breeding bird species in the case of white-tailed deer.

This report originated because of the long-term concerns about whether the population of elk in Rocky Mountain National Park (RMNP) is overabundant. We recognize that this is not the first time this issue has been raised at RMNP. In fact, concerns about elk numbers in the park have existed since the 1930s and have been addressed by many - albeit short term - management programs over the intervening years. However, this is the first time a comprehensive review of this issue has been conducted. The objectives of this report are to: 1) present the great wealth of information, both historical and contemporary, that has been

gathered on the elk population in the park and Estes Park area in the past 70+ years; 2) summarize the research that has been conducted on the effects of elk on plant communities and other animal populations in and around the park; and 3) determine if elk population dynamics and their effects deviate from that expected under natural conditions.

PHYSICAL AND BIOLOGICAL CHARACTERISTICS OF THE PARK

Rocky Mountain National Park is located in north-central Colorado and is surrounded by a mix of private and public land (Fig. 1). Established in 1915, it encompasses 1,076 km² (415 square miles) and is visited by over three million people annually. Elevations range from approximately 2,400 m (~8,000 feet) to over 4,300 m (14,255 feet), with approximately 1/3 of the park above timberline (3,350 m or 11,000 feet). Lower elevations typically consist of dense forests and open meadows, while upper elevations are characterized by alpine tundra and large expanses of rock (Fig. 2). A detailed account of the cover types shown in Figure 2 can be found in Appendix 1.

The climate on the eastern and western sides of RMNP varies greatly. Lower elevation areas on the east side of the park are generally warmer and drier than those on the west side of the park. Average summer (June through August) temperatures vary from 14° to 17° C (57° to 63° F) in the Town of Estes Park and from 11° to 14° C (52° to 57° F) in the Town of Grand Lake. Precipitation in the summer generally ranges from 4 to 6 cm (1.60 to 2.40 inches) of rain each month in both Estes Park and Grand Lake. However, there are noticeable precipitation differences in the winter; Grand Lake averages over 360 cm (143 inches) of snow each year while Estes Park averages 64 cm (25 inches) of snow each year.

The focus of this report is the elk winter range, which is where elk spend the majority of the year (Fig. 3). The **primary winter range** is located on the eastern side of RMNP below 3,000 m (~10,000 feet) in elevation. It is approximately 450 km² (~170 square miles) and encompasses the Estes Valley and portions of RMNP. The Estes Valley is not a formal name and is only used in this report to represent the larger valley (which covers portions of RMNP and Estes Park) from the Town of Estes Park (Figs. 1, 3). This report also uses the term **core winter range**, which differs from the primary winter range. Core winter range areas consist of Moraine Park, Beaver Meadows, and Horseshoe Park, where elk concentrate their foraging activities during winter (Fig. 3).

Moraine Park, Beaver Meadows, and Horseshoe Park consist of low lying valleys created by receding glaciers 10,000 year ago. These valleys lie at 2,400 m (~8,000 feet) in elevation and run east-west to form continuous elk habitat with the Town of Estes Park (Fig. 3). The terrain of Moraine Park, Beaver Meadows, and Horseshoe Park consists of large, open meadows and shrubland areas that are bordered by lodgepole pine (*Pinus contorta*) and Douglas-fir (*Pseudotsuga menziesii*) on north facing slopes and ponderosa pine (*Pinus ponderosa*) on south facing slopes. During winter, elk forage almost exclusively in the open meadows and adjacent slopes that consist of grassland, upland shrub, and riparian willow habitat.




Figure 1. Land ownership adjacent to and near Rocky Mountain National Park (RMNP).




Figure 2. Elevation (in feet) and major vegetation covertypes of Rocky Mountain National Park.




Figure 3. Elk ranges in Rocky Mountain National Park and vicinity.

The morphology and vegetation in and around the valley floors of Moraine Park, Beaver Meadows, and Horseshoe Park is very similar. Each has a branching river that runs through the valley floor and is bordered by riparian shrubs and mesic grasslands. Dry grasslands occur on the valley floors away from the rivers, where water tables are typically lower. On south facing slopes with <40% canopy cover of Ponderosa pine, upland shrubs tend to dominate. Scattered stands of aspen (*Populus tremuloides* Michx.) are found throughout the dry grassland and upland shrub areas. The valley floors and upland shrub areas make up almost 1,300 ha (3,200 acres) of winter range within the park. Of this, approximately 36% can be classified as riparian shrubs and mesic grasslands, 32% as dry or upland grasslands, and 11% as upland shrub areas.

Common riparian shrub species found in these areas include mountain willow (*Salix monticola*), flat-leaved willow (*S. planifolia*), Geyer willow (*S. Geyeriana*), mountain birch (*Betula occidentalis*), and alder (*Alnus tenuifolia*). Mesic grasslands and the understory of willow and birch communities consist of Canadian reed grass (*Calamagrostis canadensis*), sedges (*Carex* spp.), smooth brome (*Bromus inermis*), Kentucky bluegrass (*Poa pratensis*), timothy (*Phleum pratense*), cow parsnip (*Heracleum sphondylium*), and golden banner (*Thermopsis divaricarpa*). Dry grasslands are made up of a variety of grasses (*Agropyron* spp.) and sedges (*Carex* spp.), timber oatgrass (*Danthonia intermedia*), Parry's oatgrass (*Danthonia parryi*), and long-haired needlegrass (*Stipa comata*). Upland shrub areas consist of antelope bitterbrush (*Purshia tridentata*), big sagebrush (*Artemisia tridentata*), rabbitbrush (*Chrysothamnus viscidiflorus*), blue grama (*Bouteloua gracilis*), and mountain muhly (*Muhlenbergia montana*) (Stevens 1980a). Elk diets on the park winter range from fall to spring primarily consist of graminoids (58-76%), willows (8-16%), upland shrubs (0-13%), pine trees (0-12%), forbs (0-7%), and aspen (0-7%) (Singer et al. 2002; see also Hobbs et al. 1981).

The Town of Estes Park lies within the primary winter range and has similar plant species as found in the park, but has more extensive grasslands, two golf-courses that are fertilized and irrigated, and is slightly lower in elevation. Elk are able to freely use these winter range areas because they have habituated to human activities. Elk diets during winter have not been investigated in the town area.

The majority of elk (75-90%) spend summer and early fall on the western side of the Continental Divide (in the Kawuneeche Valley) or the higher elevations in RMNP (Figs. 2, 3). The Kawuneeche Valley is centered around the Colorado River and has extensive wet meadow areas surrounded by lodgepole pine. Vegetation in the Kawuneeche Valley is similar to Moraine Park and Horseshoe Park; however, winter use by elk is minimal in comparison to the east side, due to greater snowfall.

Alpine tundra provides the other principal summer range for elk. There are over 134 km² (50 square miles) of alpine tundra in the park and elk use ~39% of it from June until October (Figs. 2, 3). Tundra vegetation has been described by Kiener (1939), Griggs (1956), and Willard (1963), which Stevens (1980b) summarizes into the following types: fellfields covered by cushion plants (e.g., mosses); alpine turf and alpine marsh, both primarily

composed of sedges and wildflowers; snowbed, characterized by rushes, grasses, and wildflowers; and riparian willow areas. The distribution of these plant associations is primarily determined by physiography, snow accumulation, moisture availability, exposure, temperature, and substrate (Kiener 1939, Griggs 1956, Willard 1963, Stevens 1980*b*). A detailed list of plant species and communities found in the alpine tundra and other vegetation communities of RMNP can be found in Appendix 1.

HUMAN HISTORY OF THE REGION

Humans have played a role in the RMNP area for thousands of years by harvesting plants and animals, and more recently by development, fire suppression, haying, logging, market hunting, predator control, ranching, recreational activities, and water diversions. The effects of many of these past and current human activities are just now being fully realized by researchers and cannot be discounted when considering such issues as elk and vegetation interactions.

Little is known about the first humans to spend time in RMNP. Clovis spear points indicate prehistoric people were in the area immediately after the last glaciers receded about 10,000 years ago (Butler 1997). The Ute Indians have been the most consistent users of the park and probably have been in Colorado for at least 7,000 years. Although moved to reservations in Utah and southwestern Colorado in 1868, they continued to use the park on a sporadic basis until the late 1800s. Archeological evidence also indicates that the Apache were in the park and surrounding mountains by about 1500, and were known to be in the area in the early 1800s (Toll 1962, Brunswig 2001, Brunswig and Doerner 2001). The Arapaho arrived in the area in the late 1700s or early 1800s and numerous battles are recorded between them and the Apache and Ute for the plants and animal resources in the park (Toll 1962). Due to the harsh winter conditions of the mountains, hunters and gatherers from prehistoric and early historic times only used RMNP seasonally from late spring to early fall (Benedict 1992, Butler 1997).

The only known ethnographic account of Native American use of the park prior to European American settlement in 1860 is *Arapaho Names and Trails: A Report of a 1914 Pack Trip* (Toll 1962). The Arapaho indicate that the area was a “game bag” where bison, elk, and deer were hunted, but little information exists to indicate how consistent their use was or what their effect on animal populations might have been. Information on Ute or Apache use of the area is non-existent except for the archeological record. Unfortunately, the highly acidic and basic soils (depending on parent rock origin) in the area do not allow for the long-term preservation (>200 years) of animal bone in archeological sites through which quantification by species can be made (B. Butler personal communication).

Some of the earliest evidence of consistent seasonal use of the park are rock game drives and associated ambush pits. These drives and pits date from about 7,000 years ago until the Native Americans acquired horses in the late 1700s. Game drives were typically built above treeline and consisted of rock walls arranged as a funnel that allowed humans on foot to herd animals towards hunters hiding behind rock blinds or in ambush pits (Benedict 1996). It is unknown how effective game drives were, but it is clear that they were used almost

exclusively for elk, and that some of the largest game drive systems in the southern Rocky Mountains are located in the park (Benedict 1996). Benedict (1996) estimates that the largest complex in the park would have required about 50 people to operate, indicating that a substantial amount of elk may have been harvested from these drives. However, it is impossible to determine a numerical range or frequency of such harvests. There are at least six rock game drives in the park, and 51 have been recorded just south of the park in the Indian Peaks Wilderness (Cassells 1995).

High altitude game drives are only one source of obtaining meat, and it is worth noting that the atlatl and bow and arrow were used for thousands of years throughout the area, especially in the trees and meadows at lower elevations. Thus, although many animals might be acquired in a single game drive, the animals obtained via bow and arrow may have accounted for the majority of animals harvested throughout an entire season (B. Butler personal communication).

Bison were present in RMNP during the 18th and 19th centuries, but there is no evidence to indicate if they were present prior to these times. Toll (1962) notes that by the time the area was settled the Arapaho had moved north to hunt bison because none were left in the RMNP area. Fryxell (1928) documents numerous accounts of bison remains in and around the park, such as a large cache of bison skulls found along the current day Devil's Gulch Road. Fryxell (1928) also recounts Oliver Toll's 1913 trip with two Arapaho elders, during which the Arapaho described numerous bison hunting grounds they used in the mid-1800s. These areas included locations in Estes Park (e.g., near Mount Olympus), RMNP (e.g., Thatchtop Mountain), and Grand Lake. However, it is unknown how many bison inhabited the RMNP area or what their migration patterns were like.

The first permanent settlers in Estes Park was the Estes Family, who moved to the area in 1860. Initially, the Estes family and other settlers made a living by market hunting and ranching. These activities resulted in the extirpation of elk by 1880 (Sprague 1925, Buchholtz 1983) and likely resulted in large changes to current elk winter range areas. Gray wolves and grizzly bears (*Ursus arctos horribilis*), once commonly observed in the area, were noted to be gone by the early 1900s (Sprague 1925, Estes 1939, Stevens 1980a, Buchholtz 1983). All of the core winter range areas on the east side of the park (Moraine Park, Beaver Meadows, and Horseshoe Park) were intensively grazed by livestock, hayed, and developed by the late 1800s (Estes 1939, Buchholtz 1983). With the loss of elk, difficulties associated with ranching in the area, and eventual creation of RMNP in 1915, the primary source of income for the area turned from hunting and ranching to tourism related activities. Ranchers began operating lodges and opportunities such as climbing Longs Peak attracted visitors from around the world. Settlement of the area resulted in groups such as the Estes Park Protective and Improvement Association who, in conjunction with the U.S. Forest Service, reintroduced 28 elk to Estes Park in 1913-14.

Settlement on the west side of the park in the Kawuneeche Valley paralleled that on the east with unsuccessful mining (1879-1884) and ranching operations giving way to tourism and resorts by the time the park was established. Although the Kawuneeche Valley today

contains some relatively large riparian areas, it is unknown what conditions were like prior to European American settlement. The Grand Ditch (created in 1895) diverts 30% (10 m³/s or 360 cfs) of the total annual runoff from the upper Colorado River watershed and has likely contributed to a much drier meadow-like environment than that which historically occurred in the valley (Cooper et al. 2000). An oral history interview indicates that much of the area from Grand Lake north to the Holzwarth Never Summer Ranch was cleared of trees in the late 1800s to produce open areas to raise grass for cattle and horses. Areas to the west of the park, in particular Middle Park and North Park, seem to have been preferred by Native Americans for hunting over the Kawuneeche Valley due to the numerous large herds of bison, antelope, and elk found throughout these areas (B. Butler personal communication).

Settlement of the RMNP area appears to have also caused an increase in fire frequency during the 1800s, followed by a large decrease during the 20th century. Veblen et al. (2000), working in areas adjacent to the park, found fire frequency increased dramatically during the 19th century due to a dry climate and settlement activities such as mining. Since the early 1900s, however, fire suppression efforts in the area have been widespread, with varying levels of effectiveness (e.g., Veblen et al. 2000, Sherrif et al. 2001). Historic photos from RMNP and the surrounding area indicate lower elevation forests, consisting primarily of ponderosa pine (*Pinus ponderosa*), were much more sparse in the 1800s compared to current day conditions (Veblen and Lorenz 1991).

Today, tourism related activities remain the primary source of income in the towns of Estes Park and Grand Lake (U.S. Census Bureau 2001). Development has also increased dramatically during the last 20 years, and there are now over 10,000 residents in and around Estes Park. Current zoning indicates Estes Park is 45% developed (year 2000 estimate), and building projections indicate build-out will be reached in approximately 40 years (Hobbs et al. 2003). Likewise, considerable development can be found just outside the park's boundaries in the Grand Lake area.

Summary

There is little information to indicate what conditions might have been like when Native Americans inhabited the park or when settlers arrived. Elk were noted to be plentiful and wolves and grizzly bears were often seen in the area, but there is no specific information on animal abundance or vegetation conditions. It is clear, however, that humans have historically and will continue to play a large role in the RMNP ecosystem. Native Americans used the area seasonally and hunted elk extensively, but the size or consistency of their harvests are unknown. Settlement had a dramatic influence on the area by eliminating the major predators and elk from the RMNP area, and later reintroducing elk and developing homes and ranches on the east and west sides of the park.

THE ELK POPULATION: HISTORICAL INFORMATION AND CONTEMPORARY RESEARCH

Introduction – Elk Population Indices and Models

This report uses various indices and models to describe elk population dynamics and their effects on vegetation. This section provides definitions for terms and a summary of an ecosystem simulation model frequently referred to in this report.

Ecological Carrying Capacity (K) – Various definitions of carrying capacity exist in the scientific literature. This report uses the concept and definition of ecological carrying capacity (also referred to as K), defined as the level at which the elk population is limited by the forage base of a given area (Caughley 1976). The elk population is considered to be at K when there is a long-term stability of elk numbers and vegetation biomass on the winter range. Ecological carrying capacity can also be thought of to represent the largest number of elk that can be supported by the winter range and it does not consider the potential effect of predators that have been extirpated. Wolves may or may not limit elk at some level below K.

Estimates of K for winter range areas in RMNP and the Estes Valley have been made using several methods. These include 1) measuring annual forage biomass and energy requirements of herbivores to calculate the number of elk that can be supported by the winter range; 2) fitting a population model to elk survey data to determine if the elk numbers exhibit long-term stability (>10 years); and 3) using an ecosystem simulation model that incorporates plant and soil responses to herbivory, elk population dynamics, and weather to estimate the total number of elk that can be supported over a long-term time frame. Results from all of these methods are presented and discussed below.

Estimates of K vary annually and over longer time frames because the winter range forage base is largely dependent on annual precipitation. Elk numbers do not correspond exactly to such estimates because they are long lived animals and their population responses can be delayed and are also affected by winter severity, human harvest, and predation. Despite such variations, elk population dynamics and estimates of K are valuable tools for managers because they provide benchmarks to describe research results and management goals.

Ecosystem Simulation Model (SAVANNA) – Numerous results in this report refer to an ecosystem simulation model called SAVANNA (Coughenour 2002). This model was originally developed for studies of African grasslands and has been used in western U.S. and Canadian national parks as an ecosystem management tool where ungulates and their interactions with plants are a principal concern. The model is unique because it represents linkages between plant production and population processes, as well as between ungulate energy balance and their population processes. SAVANNA was parameterized for RMNP to provide an objective tool to evaluate historic, current, and future elk and vegetation conditions under different management scenarios. The following is a description of the model taken from Coughenour (2002):

SAVANNA is a spatially explicit, process-oriented model of grassland, shrubland, savanna, and forested ecosystems developed originally for studies in East Africa. The model has been applied to Elk Island National Park in Alberta, the Pryor Mountain Wild Horse Range, northern Australia, South Africa, and Tanzania. SAVANNA simulates processes at landscape through regional spatial scales over annual to decadal time scales. The model is composed of site water balance, plant biomass production, plant population dynamics, litter decomposition and nitrogen cycling, ungulate herbivory, ungulate spatial distribution, ungulate energy balance, and ungulate population dynamics submodels. A wolf predation and wolf population dynamics submodel were derived from a model used to assess wolf reintroduction into Yellowstone National Park.

The model was used to represent plant and soil responses to varying levels of herbivory, food limitation of the herbivore population, and predation. The model was based on a single vegetation map that included RMNP, Estes Park, and Arapahoe-Roosevelt National Forest. Simulated elk populations took into consideration the number of elk removed by hunter harvest. SAVANNA evaluated elk dynamics and vegetation conditions under a variety of ecological and management scenarios, most notably:

Undisturbed model simulations – includes wolves, undisturbed vegetation and water table levels (both estimated with historic photos), and weather data from 1775-1909.

Current model simulations – current vegetation and water table levels and weather values from 1949-1998. These simulations were done with and without elk management reductions and/or wolves to evaluate interactions between elk dynamics, vegetation conditions, and predators.

Future model simulations – current vegetation and water table levels and stochastic (random) weather patterns from 1949-1998. Future model simulations were done with and without elk management reductions, wolves, and/or fencing on the winter range to estimate the effects of various management options.

Statistical Differences – This report often refers to significant differences due to elk herbivory (e.g., vegetation production inside versus outside exclosures) or other factors. Unless otherwise noted, all differences reported in this report are statistically different using an alpha value of 0.05.

Pre-Settlement Conditions

It is clear that elk were numerous in the Estes Valley prior to settlement by European colonists, but there is no information on the magnitude of the population. The first permanent settlers in the Estes Valley were the Estes family, who arrived in 1860 and noted an abundance of elk and deer, but did not provide any population estimates. Estes (1939) wrote: “One fall and winter the writer killed one hundred head of elk, besides other game, such as mountain sheep, deer, and antelope. By this time [1863] we had made a trail to Denver, where we sold many dressed skins and many hindquarters of deer, elk and sheep”. Market

hunting in the Estes Valley did not end until the late 1870s when elk had been eliminated, or nearly so, from the area. Abner Sprague, one of the first settlers in Moraine Park, described the decline: “Our [elk] only lasted about three years. They came down from their high range just before Christmas, 1875, by the thousands and were met by hunters with repeating rifles and four horse teams; hauled to Denver for three or four cents per pound. In 1876 fewer came down; in [18]77 very few were seen on [the east] side of the divide. In 1878 I killed my last elk, and to get him had to go over Flat Top [Mountain]” (Sprague 1925). This was the last recorded kill of an elk during the 1800s, but it is impossible to determine the accuracy of Sprague’s statement that there were of “thousands of elk”. It is clear, however, when Estes’s first-hand accounts are also considered, that elk were indeed abundant but were extirpated (or nearly so) from the Estes Valley by 1880. No viable population existed again until elk were reintroduced in 1913-14.

SAVANNA simulated pre-settlement scenarios with and without wolves. When wolves were present, the total elk population size fluctuated between 1,500 and 3,500 animals. Fluctuations in elk numbers were due to a tri-trophic equilibrium between elk, plants, and wolves (Coughenour 2002). In time periods with more plant biomass there were more elk, but the population was still limited by wolves.

A pre-settlement scenario without wolves was also examined to determine the role of wolves in elk population dynamics. Without wolves, elk were found to increase to approximately 3,800 animals and then exhibit a long-term decline in numbers due to deteriorating forage conditions. With no wolves present the elk population size was only limited by available food resources. This can also be thought of as a bi-trophic equilibrium between elk and plants (versus the tri-trophic equilibrium above). Coughenour (2002) concluded that wolves limited the elk population size to 15-40% below their food limited carrying capacity. Elk in the presence of wolves also maintained a higher body condition than those without wolves.

Although it has been suggested that elk from the park migrated to the plains during winter (Clarke et al. 1994), there is no evidence for this assertion. Estes (1939), referring to animal migrations during the 1860s, stated “winter drove all the game down to the foothills, except the elk, they would remain in the park [referring to the Estes Valley] until summer, then they went up over the range or mountains”. Within RMNP, the comparison of historic game drives and current elk migration patterns suggests elk use the same routes today as they did historically (Benedict 1996, Larkins 1997).

Early Management (1915 – 1968)

Twenty-eight elk were reintroduced into the Estes Valley in 1913-14 by the U.S. Forest Service and Estes Valley Improvement Association. No hunting was allowed in or near the Estes Valley until 1939 and, along with the creation of RMNP in 1915, the population quickly grew from 30 animals in 1915 to approximately 350 animals by 1930 (Stevens 1980a; Fig. 4). Concern about the size of the population first arose in the early 1930s because elk started eating the bark of live aspen trees (McLaughlin 1931 *cited in* Guse 1966). Wright et al. (1933) concluded this was an indication that “the elk herd was reaching the limit




Figure 4. Approximate number of elk inhabiting the Estes Valley during winter from 1913 (initial year of reintroduction) to 2005. Estimates are derived from a number of sources, including Stevens 1980a, Lubow et al. 2002, and unpublished data (Rocky Mountain National Park). Estimates made prior to 1994 consist primarily of ground counts and their accuracy is unknown. Estimates since 1994 consist of aerial counts for the park, ground counts for the town, and are corrected with a sightability model (Lubow et al. 2002). Decreased elk numbers since 2002 are due in part to elk migrations out of Estes Park during winter.

of its food supply and that range abuse and starvation were in the offing”. Wright et al. (1933) suggested that a major cause of such effects was that the most important elk winter range areas were privately owned (e.g., Beaver Meadows). These areas were heavily grazed by cattle and horses during the summer, leaving little forage for elk in the winter. To remedy the situation, the park purchased over 12 km² (~5 square miles) of private land in core elk winter range areas (i.e., low lying areas of Moraine Park, Beaver Meadows, and Horseshoe Park) in 1932 (Ratcliff 1941a).¹ No other immediate actions were recommended in favor of studying the effect of these new areas on range conditions.

Subsequent studies through the 1930s and 1940s indicated grass cover and production had stabilized or was increasing, but shrubs and aspen on the elk winter range were still heavily browsed (Dixon 1939, Ratcliff 1941a, Packard 1947). Ratcliff (1941a) observed heavy use and mortality of mountain sagebrush (*A. tridentata* ssp. *vaseyana*) and willow (*Salix* spp.) throughout Mill Creek, Moraine Park, Beaver Meadows, and the Deer Mountain area. Aspen stands located on grassland areas of the elk winter range (e.g., Moraine Park) were not able to reproduce because all of the young trees were being eaten each year (Dixon 1939, Ratcliff 1941a). Ratcliff (1941a) estimated there were 1,100 elk and 1,400 mule deer in RMNP during the winter of 1940 (Figs. 4, 5) and recommended the numbers of both species be reduced.

It is difficult to determine the relative effects of elk and deer in the early 1940s. Elk and deer were beginning to rebound by the 1940s. The winter diet of mule deer consists largely of upland shrubs, such as sagebrush and bitterbrush, but summer diets are more variable (Kufeld et al. 1973). The reports of numerous deer present in 1940 and apparent heavy sagebrush mortality implies deer were affecting upland shrub communities. The deer population appeared to peak in the 1930s and was well in decline by the 1940s (Fig. 5). Subsequent declines throughout the 20th century were most likely due to increasing competition with elk or a deteriorating forage base caused by the high numbers of deer in previous years.

The complete mortality of young aspen trees and high utilization of willow suggests elk were affecting these vegetation types in certain areas (Dixon 1939, Ratcliff 1941a). It is also likely that the effects of previous cattle grazing activities were still contributing to range conditions (as predicted by Wright et al. 1933). Elk may have still avoided areas in the park during the early 1940s (e.g., Moraine Park still had two large, private ranches), as they were observed to concentrate in Mill Creek and Beaver Meadows (Dixon 1939, Ratcliff 1941a).

Because the elk population was still increasing and range conditions were considered to be deteriorating (Dixon 1939, Ratcliff 1941a), an elk and deer management plan was written for RMNP in 1943 (Condon 1943 cited in Guse 1966). The plan stated “300 elk and 200 deer should be removed from the actual herds of 705 elk and 717 deer counted in April 1943”. The goal of the reductions was to reduce the grazing and browsing effects on native vegetation.

¹ Despite the purchase, there were still significant tracts of core winter range areas that remained private. These were not purchased until the 1960s (Dixon 1931, cited in Guse 1966).




Figure 5. Approximate deer population size in the Estes Valley during winter from 1915 to 2003 (Stevens 1980a , Colorado Division of Wildlife 1999, Conner 2004). There is no way to determine the accuracy of estimates made prior to 1960. Recent counts (2001-2003) were conducted using mark-resight methods (Conner 2004).

Direct reductions were carried out for the first time in December 1944 and January 1945, and removed 301 elk and 113 deer from the park winter range areas. Shortly thereafter, Grater (1945) suggested that the reductions were not necessary, stating that the grasses on the winter range were in fair to good condition and browse plants (such as willow) were not seriously damaged except in localized areas where elk concentrated to feed. Hillary A. Tolson, Assistant Director of the National Park Service, also wrote to RMNP in January 1946 and stated that the Director's office had a "strong dislike" for reduction programs either inside or adjacent to National Park Service units. As a result, further reductions were halted until 1949.

The park's 1945-47 annual reports generally remained optimistic about range conditions. However, in a 1949 report, the park was again focused on reducing the number of elk and deer to improve range conditions: "The major wildlife problem at this time is the overused condition of winter range by deer and elk, it is anticipated that a reduction of elk and deer will be effected during the winter of 1949-50". Such conclusions were generally based on personal, subjective observations; quantitative data on range conditions were not collected again until 1954 by Buttery (1955).²

Direct reductions resumed in 1949-50 and removed 340 elk and 100 deer in RMNP. Annual reductions of approximately 60 elk and 40 deer continued until 1962. Stevens (1980a) provides a summary of population estimates that indicate approximately 350 to 800 elk and 300 to 700 deer remained in the park during the winters between 1950 and 1962. Direct reductions in the park were terminated in 1962 for a variety of reasons. Buttery (1955) concluded upland shrubs and willow were displaying signs of recovery and overall winter range conditions were improving. Regional biologist Jim Cole also considered the range in good enough condition in 1955 to end the reduction program (Wright 1992). However, the primary reason the control program ended was because RMNP, the Colorado Department of Game and Fish, and U.S. Forest Service signed a Memorandum of Understanding (1962) that agreed to undertake a cooperative elk study program that would determine the distribution and migration routes of elk in and around RMNP.³ The state also experimented with a pre-season (now referred to as a late season) hunt in January and February 1963. The goal of these research efforts was to determine if there was any time of year when sufficiently large numbers of elk from the park were located outside park boundaries to allow hunters (versus park rangers) to harvest park herd surpluses (Denney et al. 1967).⁴

Large numbers of elk were only documented to move east of the park boundary (towards Estes Park) once during the five-year study ending in 1967 (Denney et al. 1967). Two reasons likely contributed to this elk migration out of the park. First, the only significant

² A report by Fred Packard, the park's full time wildlife technician, was published in 1947, but it was based on work done prior to the initial reduction programs in 1944-45 (Packard 1947).

³ The Colorado Department of Game and Fish is now known as the Colorado Division of Wildlife.

⁴ The agencies had good motivation to find an alternative to the elk control program within the park. At the time, the Colorado Department of Game and Fish had suspended any assistance with the reduction program because the state wanted the park opened to public hunting, and the National Park Service was under increasing pressure to end control programs in RMNP and Yellowstone (Wright 1992).

snowfall during the study occurred in the park at this time (January to March 1963). Large snowfall in the park causes elk to move to lower elevations outside the park (Lubow et al. 2002). Second, elk had likely developed a tendency to avoid the park during this time of winter because this is when rangers had lethally controlled animals during the previous 10-15 winters. The single large migration also coincided with the first pre-season 1963 hunt which removed more elk (>500) than any previous NPS control effort (Stevens 1980a). However, many of these animals were located over 30 km (>20 miles) from the park border and a NPS trapping report (1967-68) suggested they were not part of the RMNP elk herd. Harvests throughout the rest of the Denney et al. (1967) study were markedly reduced, averaging less than 75 animals per year (Stevens 1980a). Nevertheless, the agencies moved ahead with a long-term management plan to control elk numbers with public harvests outside RMNP boundaries.⁵

The long-term management plan (1967 supplement to the 1962 Memorandum of Understanding) agreed to 1) dispose of surplus elk by public hunting outside of the park, 2) establish a pre-season (i.e., late season) hunt in January and February adjacent to park boundaries, and 3) trap and transplant animals if the public hunt is not effective. When the state was not able to hold a January and February hunt in 1967, the park trapped and transplanted 55 animals during the winter of 1966-67. By the summer of 1967, the park and state believed they could keep the population size stable if the park population was initially brought down to approximately 400 animals (National Park Service trapping report 1967-68). In an effort to lower the population to 400 and hold it steady, the park trapped 175 elk and the state conducted a January and February hunt in 1968. Initial results were encouraging as 268 animals were harvested during this hunt; but these animals were either not from the park population or the park population had been underestimated in previous years because elk numbers appeared to more than double in size during the following years (Fig. 4; Stevens 1980a). Late season hunts outside the park continued today, but the 1967-68 transplants were the last time RMNP directly reduced the herd. This marked the beginning of a management era that has become widely known as “natural regulation” (Huff and Varley 1999).

Natural Regulation

As it did for NPS natural resource management policies for the first six decades of the 20th Century, Yellowstone provided the model and testing ground for a profound change in the way natural resources, particularly animal populations, would be managed in national parks in the remaining portion of the century. These changes came about because of increased public scrutiny to Yellowstone’s existing elk management policy, which like RMNP, involved routinely culling animals to keep the population at a prescribed level. Following contentious congressional hearings regarding culling activities in Yellowstone (U.S. Senate

⁵ Note: Denney et al. (1967) suggested that the marked elk in the study did not routinely move out of the park because they were captured in the wrong area (i.e., Moraine and Horseshoe Park). They suggested that additional summer trapping efforts in the northeast corner of the park would show elk moving out of RMNP during the winter. However, this is largely inconsequential because the issue at hand was elk population size in Moraine Park and Horseshoe Park, not simply finding animals to harvest outside RMNP.

1967), it became clear to the NPS that a politically more appealing alternative was needed. The alternative adopted was to allow the ungulate populations in the park to reach levels that would be limited only by range potential and other ecological factors such as winter severity and hunting outside the park. This paradigm would test the idea that 1) ungulate populations might be limited by food or weather (i.e., naturally regulated) rather than held to an arbitrary level by culling and 2) this would not result in deteriorated or undesirable range conditions. This management era (1968-present), marked by little to no management interventions with ungulate populations, has become widely known in the media and scientific literature as “natural regulation” (Huff and Varley 1999). For the sake of clarity, we too use the term natural regulation.

When natural regulation was postulated it was set up as a hypothesis to be modified as more knowledge was gained. Yellowstone’s Chief Biologist in fact termed the effort “experimental management”, an approach very much akin to what is today referred to as adaptive management (Cole 1971). In keeping with this approach, a long-term monitoring program was to be included and the hypothesis was to be modified as information dictated. However, a monitoring plan was never undertaken and the experimental management became existing management. If it had been, a series of alternative hypotheses would have been undoubtedly formulated. One of these probably would have been that predation limits populations below some level that would be dictated by range potential and a weather-forage interaction (J. Peek personal communication). As it was, the initial hypothesis was aimed at determining whether or not the relatively protected Yellowstone elk population would actually over-graze its range, or if density-dependent population control factors - primarily the interaction of food and climate on reproduction and survival - would act to reduce the population growth rate and stabilize the population before there were undesirable changes in the forage base.

Implicit in the initial natural regulation management scenario was the fact that there would be little or no management intervention with the dynamics of the elk population (Peterson 1999). The Yellowstone natural regulation model was adopted by RMNP in 1968 although it was recognized that conditions at RMNP were considerably different than at Yellowstone. Rocky Mountain National Park never completely advocated natural regulation because winters are mild and the majority of historic winter range areas for elk are outside the park and largely influenced by humans (e.g., Estes Park). The concept of natural regulation as practiced in RMNP always included the understanding that hunting adjacent to the park or some other type of control was necessary to help control the elk population to fulfill the role of extirpated predators (Stevens 1980a).

Science-Based Management

National Park Service Management Policies (NPS 2001) state:

“The Service will adopt park resource preservation, development, and use management strategies that are intended to maintain the natural population fluctuations and processes that influence the dynamics of individual plant and animal

populations, groups of plant and animal populations, and migratory animal populations in parks...Biological or physical processes altered in the past by human activities may need to be actively managed to restore them to a natural condition or to maintain the closest approximation of the natural condition in situations in which a truly natural system is no longer attainable. Prescribed burning and the control of ungulates when predators have been extirpated are two examples. The extent and degree of management actions taken to protect or restore park ecosystems or their components will be based on clearly articulated, well-supported management objectives and the best scientific information available.”

The following section summarizes research that has been conducted on elk population dynamics in the Estes Valley area since the onset of natural regulation. These results allow for an opportunity to evaluate if elk population dynamics and vegetation conditions are within their natural range of variability. This report defines the natural range of variability as elk population processes and vegetation conditions that would exist in the absence of human dominance over the landscape and a viable wolf population in the RMNP area. Wolves are the major predator of elk and have the potential to directly alter ungulate abundance and foraging behavior through predation (Orians et al. 1997, Kunkel and Pletscher 1999, Peterson 1999, Hebblewhite et al. 2002).

We recognize that the natural range of variability as described here is not entirely complete. Historic records and the existence of large, pre-settlement game drives for elk in RMNP indicate Native Americans hunted in the area (reviewed above). The effect of a completely intact predator base – including wolves, grizzly bears, and humans – would likely be greater than wolves alone (e.g., Orians et al. 1997). Thus, the results presented here should be considered conservative, representing the maximum number of elk that would occur in the RMNP area.

The predicted effect of wolves on elk population dynamics in the RMNP area was determined below with the SAVANNA ecosystem model (Coughenour 2002, description above). Elk and vegetation conditions were simulated using the current amount of available habitat in the presence of wolves. Weather was selected randomly using local data from 1949-1998.

Elk Research (1968 – present)

Elk Distribution and Movements – Elk in the RMNP area spend approximately seven months per year on the core and primary winter range (November to May) and three months on summer range (mid June to September). The remaining two months are spent on or in transition between these two ranges (Fig. 3). Within their seasonal ranges, elk move in response to various factors, including weather and hunting (Larkins 1997). Elk respond to hunting by moving from areas that are open to hunting to areas where hunting is prohibited.

The elk population includes three winter subpopulations that exhibit different population dynamics and migration patterns: 1) Moraine Park and Beaver Meadows (referred to as the

Moraine Park subpopulation), 2) Horseshoe Park, and 3) the Town of Estes Park and adjacent National Forest Service areas (referred to as the Estes Park subpopulation) (Fig. 6; Larkins 1997, Lubow et al. 2002). Elk tend to stay within these areas, although 15% (400 to 500 elk) regularly move between subpopulations. The population dynamics of the Moraine Park and Horseshoe Park subpopulations do not differ and will be collectively referred to here as the park subpopulation. Elk population dynamics are different between the park and town subpopulations. There is also a small group of elk that spend winter on the alpine tundra in RMNP (100-200 animals; years 1980 and 2000 estimates), but they are not considered in this report as little is known about their migration patterns or population dynamics.

The elk population exhibits large, seasonal migrations between winter and summer ranges (Fig. 7). The timing of migration depends on weather, in particular snow depth, which determines when forage on the summer range will become available. Generally 75-90% of the population migrate to higher elevations or the Kawuneeche Valley for the summer (Larkins 1997, RMNP unpublished data). Elk from the three major winter range areas exhibit distinct migration routes to and from their summer ranges (Larkins 1997). Elk in the Moraine Park subpopulation migrate over the Ute Trail and to the Kawuneeche Valley, where they calve, and most remain for the summer, although some migrate back to alpine areas for the summer. Elk from Horseshoe Park and Estes Park migrate up the Fall River drainage and spend summer in alpine areas of the park. However, the timing of the migrations of the two subpopulations is different, with Horseshoe Park elk migrating to higher elevations in the Cache la Poudre and Fall River areas to calve, while elk from town calve in Horseshoe Park before proceeding to higher elevations for summer.

Approximately 20% of the primary winter range falls within park boundaries (Fig. 3), extending from Cow Creek on the north to Hollowell Park on the south and east to Hidden Valley. Approximately one-third of elk in the area winter in the park (Lubow et al. 2002); however, park subpopulations make temporary cross-boundary movements out of the park to lower elevations, especially during snowfall events.

Most of the winter range is outside the park in the Town of Estes Park and on adjacent private and U.S. Forest Service Lands (Fig. 3). The Town of Estes Park provides continuous elk winter range with Moraine Park and Horseshoe Park. Elk are attracted to the town area for several reasons, including extensive grasslands, some of which are nutritionally enhanced by fertilization (e.g., golf courses); little to no animal predation; and lack of hunting. In addition, the town lies at a lower elevation than winter range areas in the park, and consequently has lower snow depths and more favorable winter temperatures for elk (Singer et al. 2002). During winter, elk also range east of the Estes valley, with elk numbers and time periods varying depending on habitat conditions, snowfall, and hunting. Elk range north to Crosier Mountain and Glen Haven and east to Drake and Bobcat Gulch and the vicinity of the Meadowdale Ranch (Bear 1989, Larkins 1997). Elk also use Storm Mountain, Jug Gulch, Bear Gulch, and the Pole Hill-Nixon Park areas. Other elk populations use areas east of the Estes Valley (Bear 1989); the extent to which elk from the RMNP area mix with these populations is not known, but it is estimated that up to 1,000 elk from the RMNP population spend several months per winter in these areas (R. Spowart personal communication).




Figure 6. Elk subpopulation boundaries during winter on the east side of Rocky Mountain National Park (RMNP) and vicinity.




Figure 7. Elk migration routes and major stream drainages in the Rocky Mountain National Park (RMNP) area. Elk that winter in Moraine Park tend to follow the Ute Trail over the Continental Divide and spend their summer in the Kawuneeche Valley. Elk that winter in Horseshoe Park and in and around the Town of Estes Park follow the Fall River corridor and summer in alpine areas of RMNP (Larkins 1997). Elk from the RMNP area have also recently been observed to migrate further east during winter (represented in the figure by a ?), but the migration routes and seasonal ranges of these elk have not been studied.

Primary summer range areas in the park are at higher elevations in alpine and subalpine and alpine habitat, and in the Kawuneeche Valley on the west side of the park (Figs. 2, 3). Outside the park, elk summer in the Twin Sisters, Lion Gulch, Pierson Park, Lake Pasture, Big Elk Park, and Big Elk Meadows areas (Larkins 1997, J. George personal communication). At least 10-15% of elk in the RMNP area have been documented to spend summer on the winter range. Recent park surveys indicate that during summer, at least 100 to 200 animals stay on the park winter range areas, and as many as 550 animals stay on town winter range areas (RMNP unpublished data). The only large group of elk (>300 animals) that summer on the winter range generally uses the Meadowdale Ranch and 18-hole golf course on the east end of Estes Park. This group has been in this area since at least the 1970s (Stevens 1980a).

In the fall of 2002, three elk that were radio-collared in the park during summer were observed just west of the town of Loveland, along with a much larger group of elk than the resident population in that area (Figs. 3, 7). This coincided with much smaller population estimates in the Estes Valley than previous years, suggesting that at least 200 to 300 elk that normally wintered in the Estes Valley moved eastward. Following the 2002-2003 winter, two of the radio-collared elk returned to summer in the park, while the third remained with the resident population near Loveland. Since that time, the size of the population in Loveland has remained higher than it was prior to 2001, while population estimates in the RMNP area have been lower, particularly for the town subpopulation. The cause of any potential shift eastward is unknown, although suggestions have included drought, several significant snowfall events, and limited forage due to high elk densities in the Estes Valley. It is unknown whether some or all of these elk still summer in the park and whether these shifts are temporary or long term in nature.

Elk Population Dynamics – The population dynamics of the Moraine Park and Horseshoe Park subpopulations do not differ and are collectively referred to as the park subpopulation (Lubow et al. 2002). This subpopulation has reached carrying capacity and is limited by forage availability (Coughenour 2002, Lubow et al. 2002, Singer et al. 2002). Estimates for the park population over the last 10 years have ranged from approximately 700-1,100 elk. After reductions within RMNP were stopped in 1968, the park subpopulation initially increased at an annual rate of 6.5% and then gradually slowed its rate of growth to reach an average carrying capacity of 1,069 animals in 1991 (Fig. 8; Lubow et al. 2002). This population based carrying capacity estimate is consistent with three other ecological carrying capacity estimates based on either ecosystem simulation modeling or forage biomass and energy considerations. (1) The SAVANNA model predicts that under current management conditions, the park subpopulation will continue to fluctuate between 800-1,100 elk (Coughenour 2002). (2) Forage based estimates by Singer et al. (2002) indicate the population will stabilize at approximately 1,150 elk. (3) Hobbs et al. (1982) made ecological carrying capacity estimates for two years; year one was ~1000 and year two was ~1600. However, population estimates are expected to be maintained at the lower level (a dry year) because it is more limiting (Lubow et al. 2002). Such fluctuations illustrate the difficulty of computing ecological carrying capacity estimates over short time periods and emphasize the year-to-year variation that can occur.




Figure 8. Elk population dynamics (five-year moving average) in Rocky Mountain National Park (RMNP) and the Town of Estes Park (Lubow et al. 2002, RMNP unpublished data). The park subpopulation primarily resides in Moraine Park, Beaver Meadows, and Horseshoe Park. The town subpopulation resides in the Town of Estes Park and on U.S. Forest Service lands to the east. Decreased elk numbers since 2002 are due in part to elk migrations out of Estes Park during winter.

Elk did not winter in the Town of Estes Park in noticeable numbers until the late 1970s (N. T. Hobbs personal communication, cited in Lubow et al. 2002). The town subpopulation increased at annual rates of 11% from 1979 to 1983 and 5.2% from 1991 to 2001. Although the origin of the town population was likely emigration from the park, the subsequent growth of the town subpopulation appeared to be independent of the park subpopulation (Lubow et al. 2002). Population estimates reached a high between 1997 and 2000, with annual estimates ranging from 2,000-2,500 elk. Estimates from 2001 to 2005 have ranged between 1,000-1,400 elk in the Estes Valley area (Fig. 8). These lower estimates coincided with increased numbers in the park in 2001 and increased observations of elk from the Estes Valley near Loveland in years 2002 and 2003 (Fig. 7, see above).

Three carrying capacity estimates have recently been made for the town elk herd. (1) A population based estimate projects an eventual population size of 2869 elk, but has a wide margin of error (i.e., standard error ± 415) which predicts the estimate could fall between 2,454 and 3,284 elk (Lubow et al. 2002). (2) A forage based estimate predicts 3,082 elk will eventually occupy the town sector (Singer et al. 2002). (3) A SAVANNA estimate of the carrying capacity in town to be 1,400-2,000 elk, with an initial overshoot to a population size of 2,400 elk before the stabilization occurs (Coughenour 2002). It is important to note that the latter two estimates were based on vegetation maps from 1996, so development in town may have already changed their current values. The population based estimate ($2,869 \pm 415$) will also continue to change as future development removes or enhances elk habitat. Recent results suggest the elk subpopulation in town is stabilizing (as in Coughenour 2002).

Carrying capacity in the town area in 1996 was estimated to be only 5% less than if the area was still in pristine, natural condition (Coughenour 2002, Singer et al. 2002). Development and the creation of Lake Estes has reduced the land area available for elk foraging, but the creation of golf courses and other irrigated and fertilized areas has enhanced the productivity of remaining lands.

If wolves were still present, elk numbers in the park during winter would be expected to fluctuate between 300-500 elk (Coughenour 2002). Elk numbers in the town area during winter would be expected to fluctuate between 900-1600 elk (Coughenour 2002). SAVANNA indicates that a total of 1,200-2,100 elk would inhabit the primary winter range under current conditions and in the presence of wolves. This is considerably lower than the fluctuations observed over the last 10 years, which ranged from 1,700-3,500 elk in the Estes Valley. Limitation of elk populations by wolves has been observed or predicted in several North American ecosystems (Garton et al. 1990, Boyce 1993, Mack and Singer 1993, Kunkel and Pletscher 1999, Hebblewhite et al. 2002, Singer et al. 2003); and limitation of ungulate populations below K by their major predators has been observed throughout the world (Sinclair 1985, Messier and Crete 1985, Bergerud and Elliot 1986, Gasaway et al. 1992, Orians et al. 1997, Peterson 1999). Although some studies have found ungulate numbers to be limited by forage in the presence of an intact predator community (Sinclair et al. 1985, Skogland 1985, Fryxell 1987, Mduma et al. 1999), we know of no research that indicates elk population size in the RMNP area or other similar ecosystems would be stable at K in the

presence of wolves. Further, as mentioned above, the effect of a completely intact predator base that includes wolves, grizzly bears, and humans would likely be greater than wolves alone (e.g., Orians et al. 1997).

Elk Densities – Elk group sizes during winter can range from a single individual to over 600 animals in Moraine Park and the golf course areas in Estes Park (Larkins 1997). Elk densities on the winter range were estimated from 1994 to 1999 by Singer et al. (2002). Densities reported here represent the average values of 44 aerial surveys conducted in winter during the early morning and afternoon. Elk densities are variable in the park, with high (30 to 65 elk/km²) to very high (66 to 110 elk/km²) concentrations on about 7% of the winter range, centered in Moraine Park (Singer et al. 2002). The remainder of the winter range generally has moderate (10 to 29 elk/km² on 11% of the winter range) to low (<10 elk/km² on 82% of the winter range) densities (Singer et al. 2002). Although elk use lower-density areas of the winter range to rest or as they move between areas, the majority of their time spent foraging is highly concentrated on a small percentage of the winter range. Elk densities on core winter range areas that are greater than 100 elk/km² are the highest concentrations ever documented for a free-ranging population that is not artificially fed.

The average density in Estes Park is 28.5 elk/km². Site-specific density information is not available for Estes Park, but a similar pattern exists with very high concentrations in areas with excellent foraging conditions (e.g. golf courses) and lower concentrations in other areas (e.g. downtown). Elk use essentially all parts of the Estes Valley, with concentrations occurring in the area between Dry Gulch and Devil's Gulch Road, along U.S. Highway 34, the Crocker Ranch area, on both golf courses, and between Fish Creek and Colorado Highway 7 (Larkins 1997). Elk are generally less concentrated on summer range areas.

It is difficult to determine what density of elk is appropriate, but research elsewhere indicates densities of elk in RMNP during winter are outside their natural range of variation. For example, elk densities in willow and grassland habitats of the core winter range (10 to 110 elk/km²) are high compared to other free-ranging populations in North America, where they range from 1 to 36 elk/km² (Houston 1982, Singer et al. 1994, Smith and Anderson 1996, Hebblewhite et al. 2002, Nietvelt 2001). Because there are no major predators of elk in the RMNP area (Singer et al. 2002), elk behavior and distribution is altered because they can optimize their foraging strategies by decreasing their vigilance of predators (e.g., Laundré et al. 2001, Wolff and Van Horn 2003). Elk in Yellowstone, Banff National Park, and Montana have recently been shown to exist at lower densities, increase use of forested areas, and decrease use of grassland habitats when wolves are present (Nietvelt 2001, Ripple et al. 2001, Ripple and Beschta 2004, Wolff and Van Horn 2003, Hebblewhite et al. 2005, Creel et al. 2005). Similar predator avoidance behaviors have also been found in other ungulate species (e.g., Edwards 1983, Ferguson et al. 1988, Dekker et al. 1996, Schmitz et al. 1997, White et al. 1998, White and Feller 2001). These findings strongly suggest that densities of elk on the core winter range grasslands (which contain aspen and willow communities of concern, see below) have exceeded natural values due to lack of a major predator.

Elk Survival – The largest source of mortality for adult elk in the population is hunting; however, elk that spend winter in the park or Town of Estes Park are not affected by hunting unless they leave to use adjacent U.S. Forest Service or private lands where landowners allow hunting. This can occur, particularly in response to heavy snowstorms.

Lubow et al. (2002) estimated survival rates from 1965 to 2001 in the park and 1978 to 2001 in town. During these periods, bull survival increased from 52% to 79% in the park, but remained constant at 42 % in town. Low but increasing male survival in the park indicate that these individuals are also subject to harvest at some times, but vulnerability has declined over time. Lubow et al. (2002) found that adult female survival was 91% for both park and town and concluded that harvest did not appear to have a differential effect on cow survival in the park versus town subpopulations. However, in the past several years, cow harvest has increased, so this may no longer be the case.

A calf mortality study conducted in RMNP and the Estes Valley between 1979 and 1982 found that malnutrition was the most significant source of mortality (35%), followed by hunting (23%), predation by coyotes (17%), and disease (12%) (Bear 1989). Most death from malnutrition occurred in the first few weeks after birth, and was attributed to cows being in poor condition due to weather or resource limitations. As population sizes and densities increased from 1965 to 2001, calf survival to eight months of age declined in both park and town (Lubow et al 2002). Calf survival to 20 months also declined strongly in the park as elk density increased, and was present but less apparent in town. In 2001, calf survival to eight months was 35% in the park and 88% in town. Survival to 20 months was 24% in the park and 73% in the town.

Chronic Wasting Disease – Elk and deer that winter on the east side of RMNP are infected with chronic wasting disease (CWD). No other ungulates (e.g., moose (*Alces alces*), bighorn sheep (*Ovis canadensis canadensis*)) or animals are known to be susceptible to CWD. Chronic wasting disease is a transmissible spongiform encephalopathy that primarily occurs in free ranging deer and elk in northeastern Colorado and southeastern Wyoming (Miller et al. 2000). Cases in free ranging deer or elk have also been documented in South Dakota, Wisconsin, Illinois, New Mexico, Utah, and Saskatchewan. Little is known about the disease, which eventually causes behavioral changes, emaciation, excessive salivation, weakness, and death in infected animals. The disease can remain latent (i.e., no symptoms are apparent) for months and potentially years, and it is terminal in all cases. It is unknown if the disease is native, and if not, how it originated; but it has been present in the area for at least 30 years (Miller et al. 2000).

At present, CWD has only been diagnosed in elk from the RMNP area by examining brain tissue after death. Prevalence estimates of elk inside RMNP have not been determined but are believed to be comparable to surrounding areas. Past prevalence in elk in adjacent areas was estimated to be less than 1% based on surveillance from mandatory elk head submission from hunters (Miller et al. 2000); however, estimates in 2003 and 2004 range from 1% to 3% (Colorado Division of Wildlife 2003, 2004).

There are many uncertainties associated with the disease; for example, it may be able to exist and be transmitted via the environment (e.g., soils). To date, there is no evidence that suggests CWD can affect the population dynamics of elk.

Summary

Elk that winter in the Estes Valley are referred to as park and town subpopulations due to the areas they inhabit and differences in their population dynamics. The park subpopulation reached ecological carrying capacity of approximately 1,000 elk in the early 1990s. Estimates above or below this average carrying capacity estimate are due to annual variability in forage conditions, temporary migrations, timing of park surveys, and weather. SAVANNA estimates 300-500 elk would inhabit the park during winter if wolves were still present.

The town population, which also uses surrounding U.S. Forest Service land, reached a high between 1997 and 2000, with annual estimates ranging from 2,000-2,500 elk. Estimates from 2001 to 2005 have ranged between 1,000-1,400 elk. Carrying capacity estimates for the town herd vary depending on the method used, ranging from 1,400-3,300 elk. SAVANNA estimates 900-1,600 elk would inhabit the town area during winter if wolves were still present.

Thus, SAVANNA indicates 1,200-2,100 elk would inhabit the primary winter range in a tri-trophic system where elk, food, and wolves freely interact. This is considerably lower than the fluctuations observed in the current bi-trophic system, where elk are only limited by their food base and population estimates have ranged from 1,700-3,500 elk over the last 10 years. The ability of wolves to limit elk populations below ecological carrying capacity is supported by research elsewhere in North America. No estimates are available for the natural range of densities that would be expected to occur in the presence of wolves. However, densities on the core winter range areas in the park are the highest concentrations ever documented for a free-ranging elk population and it is highly unlikely such aggregations would occur in the presence of wolves.

FOCAL PLANT COMMUNITIES

Aspen

Aspen Ecology – Aspen covers less than 5% of RMNP, but provides important habitat for a large number of plant, butterfly, and bird species compared to other habitat types the park (Mueggler 1985, Simonson et al. 2001, Turchi et al. 1994). Although some stands in the park are relatively large ($>0.4 \text{ km}^2$ or 0.15 square miles), most are only a few hectares in size.

Aspen trees that are connected by their roots are referred to as a clone. Almost all aspen stands that have been studied in the western U.S. are made up of a single clone or mosaic of clones, and they are believed to almost always reproduce vegetatively (Shepperd and Engelby 1983, Schier 1985, Kay 1997a). Vegetative reproduction is when the interconnected roots

give rise to adventitious suckers that eventually become an aspen tree. Aspen establishment can also occur through sexual propagation (seeds) when conditions are suitable, but this is considered a rare occurrence in the western U.S. (Shepperd and Engelby 1983, McDonough 1985, Kay 1997b). Few aspen trees live more than 200 years (Jones and Schier 1985), but aspen clones are often hundreds or potentially thousands of years old (see also Kay 1997a).

The primary factors that affect the ability of aspen to regenerate are apical dominance, disturbances such as fire, and ungulate browsing. Apical dominance is when auxin, a plant hormone that inhibits shoot and tree growth, is transported from aerial parts of the tree to the roots (Farmer 1962, Schier 1975a, Schier 1985). When the flow of auxin to the roots is cut off, new suckers arise and the existing trees resume growth (Schier 1985). Deteriorating, overmature aspen clones often fail to regenerate because apical dominance is maintained over a shrinking root system (Schier 1975b).

Aspen is a shade intolerant tree that cannot reproduce in the presence of mature conifers. Fire or other types of disturbance (e.g., an avalanche) are thus necessary to facilitate aspen persistence in the landscape by opening up the forest canopy and preventing conifers from overtaking aspen trees. Because aspen initially grow much faster than conifers, canopy opening disturbances allow aspen to expand and maintain their clone until another disturbance can occur. In RMNP, aspen on the core winter range grasslands are not successional to conifers and do not require fire to regenerate in the absence of elk. However, elk are a natural part of the system and fire may still play an integral role in aspen stand establishment by altering elk feeding patterns, plant competition, and releasing nutrients that increase aspen growth. Aspen throughout the rest of the park are largely dependent on fire (or another type of disturbance) to sustain themselves.

Elk browsing can kill or stunt the growth of aspen suckers. When intense enough, this can prevent all suckers within an aspen stand from reaching tree size. This could eventually eliminate the aspen stand because photosynthesis will cease as the older trees die, and the root systems will become malnourished and result in the loss of the clone. However, aspen may also be able to persist for long periods of time in a shrub-like state (e.g., Dobkin et al. 2002) and it is very plausible that this allows aspen to persist until ecological conditions are appropriate for trees to become established. Thus, the presence of aspen trees may be a periodic event on the winter range made plausible by persistence in a shrub-like state when trees cannot become established.

Elk also strip the bark off of live trees. The nutritional value of bark is unknown, but aspen barking by elk is not believed to be related to food shortages (in RMNP, aspen were barked when only 350 elk, ~10% of current numbers, were estimated to be on the winter range). Bark stripping does not usually kill aspen, but it can create inoculation sites for pathogens that lead to aspen mortality (Hinds 1985).

Aspen in RMNP – Concerns about the effects of elk on aspen in the park are specific to the core winter range areas (e.g., Moraine Park, Beaver Meadows, and Horseshoe Park) and the Kawuneeche Valley (Suzuki et al. 1999). These are the most visible stands in the park by

virtue of their proximity to the road system and park entrances, however they occupy less than 2% of the winter range and make up less than 1% of the aspen in the park. The large majority (>99%) of aspen in the park are not located on the primary winter range or Kawuneeche Valley and are not adversely affected by elk browsing (Suzuki et al. 1999, Kaye et al. 2005).

Research has shown that elk browsing on the winter range has prevented aspen suckers from maturing into trees (i.e., a height ≥ 2.5 m or 8 feet in height) since at least 1970 (Baker et al. 1997, Olmsted 1997, Suzuki et al. 1999). Olmsted (1979) and Baker et al. (1997) found an inverse correlation between aspen regeneration on the core winter range and elk population size. The latter study concluded aspen stands only produced a new cohort of trees when there was no town subpopulation and the park subpopulation was less than 600 animals.

The results of Olmsted (1979) and Baker et al. (1997) suggest that elk population size is the primary factor controlling aspen regeneration and cohort establishment on the core elk winter range. However, Stevens (1980a) did not find a correlation between aspen leader use and elk population size, and suggested that elk distribution may be equally or even more important to successful aspen regeneration than overall population size. Stevens (1980a) states: “Olmsted [1979] utilized the regeneration of aspen in certain stands as an indication of a relief from browsing during the 1950’s, [but] these stands were near roads where the maximum direct reduction efforts on elk took place and therefore received little sustained use”. Baker et al. (1997) worked in the same area as Olmsted (1979) and is subject to the same concern, namely that it is impossible to determine if elk population size or density is a more important predictor of aspen utilization rates. Weisberg and Coughenour (2003), who found densities were a more accurate predictor of aspen regeneration and growth than overall population size, give tacit support to the Stevens (1980a) hypothesis.

There are two 35-year-old aspen exclosures on the elk winter range, and both contain multiple age classes of aspen. The ratio of live to dead trees is twice as high for stands inside these exclosures versus outside. Average stocking rates inside exclosures was 245,000 suckers/km² and outside exclosures was 47,300 suckers/km² (Baker et al. 1997).

Existing aspen trees on the park’s winter range are declining rapidly as they die of old age. Between 1975-76 and 1995-96, Olmsted (1997) found large trees decreased by 42%. This indicates that the existing aspen trees on the park’s core winter range are over-mature and in danger of losing their largest aboveground component (i.e., trees). All research indicates that if current elk densities persist on the core winter range, there may be a complete loss of aspen trees (Baker et al. 1997, Olmsted 1997, Suzuki et al. 1999, Coughenour 2002, Weisberg and Coughenour 2003).

The concern about the loss of aspen on the core winter range is largely due to the acceptance that the aspen stands are clones that only regenerate vegetatively and have been present for hundreds if not thousands of years. Another possibility is that the clones on the core winter range were established from seed relatively recently. Such events (i.e., recent aspen regeneration by seed) in the western U.S. have been associated with large fire events (e.g.,

Romme et al. 1995, Romme et al. 2001). However, elk herbivory killed all the suckers before an aspen stand or clone could become established (Romme et al. 1995). Aspen trees and stands may have been established in the absence of herbivory. The montane region of the northern Colorado Front Range and RMNP experienced extensive fires during the 1800s (Rowdabaugh 1978, Veblen and Lorenz 1991, Mast et al. 1998, Veblen et al. 2000). There were also no elk present in RMNP by 1880, so it is plausible that aspen on core winter range areas of RMNP are seedlings that persisted because of increased fire frequency and elk extirpation.

Baker et al. (1997) cite limited photographic evidence from the early 1900s that show aspen stands were present on the winter range prior to elk extirpation. However, the one photo they provide is clearly not on the core elk winter range because it is in a dense stand of conifers; Suzuki et al. (1999) found that many such stands on the primary winter range (in the conifers) are still able to reproduce. There are also no elk scars on the bark, which likely would have been present if the trees existed prior to elk extirpation. Baker et al. (1997) and Olmsted (1979) also aged aspen trees on the elk winter range. None of the estimated dates were prior to elk extirpation (1880), but this could also be a function of aspen longevity, which typically does not extend beyond 100 to 150 years of age (Jones and Schier 1985).

The SAVANNA model was used to simulate aspen cover on the core elk winter range. In all model runs with elk, aspen cover on the winter range was initiated at current levels and declined. Even when wolves were present and limited the elk population, aspen declined substantially. Only when elk were removed from the model was aspen able to persist or increase on the core elk winter range (Coughenour 2002). However, neither elk densities nor distribution was modeled in SAVANNA. This, along with the fact that aspen have regenerated and established new cohorts of trees during the 20th century, again suggests that predation or some other factor (e.g., past park service shooting programs) that influences elk distribution has played a key role in allowing aspen on the core winter range to recruit new trees. For example, previous research has found that elk will decrease their use of areas that are heavily hunted or has predators (Altmann 1956, Ripple et al. 2001). Such effects may allow aspen to successfully regenerate on the winter range in RMNP, but it is unclear how long elk avoidance of a given area may persist.

A separate SAVANNA modeling effort has been conducted to more intensively examine the effects of elk density on aspen regeneration (Weisberg and Coughenour 2003). Although this effort suggested similar results as Coughenour (2002), it did find that aspen were able to produce new cohorts in the presence of lower elk densities, dependent on the amount of time elk spent feeding in aspen stands. When elk used aspen for approximately eight months of the year, aspen could regenerate when there were fewer than 5 elk/km². If elk were assumed to only use aspen for three months of the year, aspen could regenerate in the presence of as many as 14 elk/km².

Baker et al. (1997) and Singer et al. (1998a) cite successful aspen reproduction in ungulate exclosures as evidence that recent changes in climate or fire absence are not responsible for aspen declines in these locations. However, because herbivory is a natural component of the

RMNP ecosystem, the interaction of these effects with other natural components of the ecosystem cannot be fully evaluated inside exclosures. For example, if it was found that even a small number of elk at a low density prevented aspen from regenerating it would suggest that the stands are either not a natural component of the system or other key factors to aspen recruitment (e.g., fire regimes, climate) may have been altered. Until such hypotheses are tested, it is not possible to completely eliminate the role of other factors such as fire, climate, or altered foraging patterns due to the presence of predators.

Willow

Three woody shrubs that occur on the winter range in RMNP are of primary interest from the viewpoint of elk herbivory: willow, bitterbrush, and sagebrush. The general distribution of these shrubs on the winter range is shown in Figure 9.

Willow Ecology – Willow establishment occurs by sexual and asexual means. Sexual propagation is by seed. Willow seeds are only viable for a few days and can only take root in wet areas with unvegetated or disturbed soils with good light availability (Cottrell 1993, Cottrell 1995). Willows in RMNP establish themselves on three principal landforms; point bars, abandoned beaver ponds, and abandoned channels or ox-bows (Cooper et al. 2003, Gage and Cooper 2004a). The latter two types are both associated with the presence of beaver in an area, suggesting beaver are not only important for asexual propagation (see next paragraph), but sexual propagation as well.

Asexual propagation (vegetative) occurs when existing willow roots or a willow stem cut by beaver gives rise to new shoots to become a new plant. Shoots from the roots of a willow are very important because they allow existing willow plants and root systems to maintain themselves for long periods of time (i.e., 100 years or longer). Beaver cuttings also allow willows to colonize areas that are suitable for willow growth but not seedling establishment, and may be important on a landscape scale (Cottrell 1995, see also Gage and Cooper 2004b). A potential long-term result of smaller beaver populations in the park is a decrease in willow cover (Cottrell 1995).

Willow growth and height in the intermountain west is often determined by large ungulate and beaver browsing (Singer et al. 1994), as well as site conditions such as soil type, length of growing season, nutrient concentrations, and water table height (Cottrell 1995, Peinetti 2000). Willows have developed two primary defenses against the effects of browsing: first, they exhibit rapid vertical growth rates that extend growth beyond the height of browsing (Bryant et al. 1983); and second, willow can produce defense compounds that make them less palatable to large ungulates (Singer et al. 1994). Neither defense is likely to be effective if ungulate consumption levels are too great, especially if intense herbivory is combined with beaver activity or poor site conditions.

Willow in RMNP – Willow is the dominant woody shrub on almost all wet meadow or riparian areas in RMNP. It is an important food source for elk (Hobbs et al. 1981, Singer et al. 2002) and beaver (Baker and Cade 1995), and supports a large number of bird and




Figure 9. Woody shrub distribution on the elk winter range in Rocky Mountain National Park (RMNP).

butterfly species on the core elk winter range relative to other habitat types (e.g., Connor 1993, Simonson et al. 2001). There are primarily three species of willow on the elk winter range; *Salix geyeriana*, *S. monticola*, and *S. planifolia* (Zeigenfuss et al. 2002). Park-wide, *S. monticola* and *S. planifolia* tend to be co-dominant below 2900 m (~9500 feet), while *S. planifolia* dominates above this elevation (Cottrell 1995). The remainder of this section covers what is currently known about willow and the effects of elk on the core winter range in RMNP. Little is known about willow communities in the Kawuneeche Valley or alpine tundra areas (see the “Alpine Tundra” section below). Willow in the Kawuneeche Valley area appear to be heavily browsed by elk and moose, but widespread willow measures have not been made and the relative effects of each ungulate cannot be determined at this time.

Willow Distribution in RMNP – Reports from the late 1800s suggest Moraine Park used to have more extensive willow and riparian shrub cover than it did in the 1930s and 1940s; it was apparently reduced by 1900 because many areas were drained and cut down for haying purposes as the area was settled (Gysel 1960).⁶ There was also considerable development (e.g., lodges, a golf course) in Moraine Park, Beaver Meadows, and Horseshoe Park as late as the 1960s (Buchholtz 1983). However, these effects were not quantified, and only the recent history of willow distribution is known for Moraine Park and Horseshoe Park.

Over the past 50 to 60 years, riparian shrub cover (assumed to primarily be willow) has declined by approximately 20% in Moraine Park and Horseshoe Park. Willow declines in Moraine Park are visibly correlated to a large reduction (69%) in surface water, caused by an almost complete loss (>90% decline) of the area’s beaver population since 1940 (Packard 1947, Peinetti et al. 2002, Zeigenfuss et al. 2002). Beaver are highly effective at building dams which create ponds and additional stream channels. Because willow growth and survival in RMNP are primarily dependent on ground water from streams and snowmelt instead of rainfall (Alstad et al. 1999), it is not surprising that large expanses of willow have died where streams have become totally dry and water tables have undergone dramatic decreases. Many stream channels in Moraine Park that were filled with water and bordered by live willow in 1937 are now dry with large, dead willow on the old stream banks.

Willow declines in Horseshoe Park are largely due to the Lawn Lake Flood (Peinetti et al. 2002), an unnatural flood event that occurred when a man-made dam that was constructed in 1903 breached in 1982. This flood caused 831,000 m³ of water to cascade down Roaring River and into Fall River, which runs through the middle of Horseshoe Park. The largest flood-related effect in Horseshoe Park was the deposition of over 750,000 tons of rock at the confluence of Roaring River and Fall River (Jarrett and Costa 1993). This resulted in an alluvial fan which created a new lake directly upstream. Existing willow in these areas were either destroyed by the alluvial fan or flooded out by the new lake (too much water can also kill a willow due to anaerobic soil conditions). Downstream, the flood was dispersed into meadow areas and was not large enough to significantly alter the hydrology of Fall River and the Horseshoe Park area (Jarrett and Costa 1993). However, Peinetti et al. (2002) attribute some loss of willow in downstream areas to direct removal by flood waters. Horseshoe Park

⁶ Moraine Park was originally named Willow Park (Buchholtz 1983).

has also experienced a large decrease (47%) in surface water, due in part to a 90% decline in the beaver population (Zeigenfuss et al. 2002). This has also likely contributed to the decline in willow, though not to the extent as in Moraine Park.

Willow Growth and Size in RMNP – Although surface water is correlated with willow presence on the winter range (Peinetti et al. 2002), the size and morphology of existing willow plants on the elk winter range in RMNP is primarily determined by elk browsing. Elk browsing levels on the winter range were found to significantly reduce willow height (Peinetti et al. 2001, Zeigenfuss et al. 2002), volume (Peinetti et al. 2001), and the number of leaves per stem (Peinetti et al. 2001) when compared to willow inside ungulate exclosures. Elk browsing was found to have a greater effect over the long-term: Schoenecker et al. (2004) found willow volume and height to be 98% greater inside versus outside a 35 year-old ungulate exclosure located in Beaver Meadows.

Areas once dominated by tall willows (at least two meters in height) on the core winter range are now dominated by short willows (less than a meter). Peinetti et al. (2002) suggests the decline in tall willows is due to the combined effects of a drastic decrease in surface water in Moraine Park and the Lawn Lake Flood in Horseshoe Park followed by elk browsing levels that are preventing small willow plants from growing beyond one meter (~3 feet). At least two other factors are facilitating a further conversion to a short willow community. First, as the few beaver that still exist in Moraine Park and Horseshoe Park select and cut down tall and relatively unbrowsed willow for food (Baker et al. 2004), elk browse the plants and keep them in a short stature (less than 30 cm). This creates a lack of food for beaver and results in a feedback mechanism that reduces beaver and willow populations (Baker et al. 2005). Second, antler rubbing during the fall can also tear down a tall willow plant. These shorter willows are then suppressed by current elk browsing levels.

Singer et al. (2002) observed elk consumption and density to be strongly correlated to willow growth, morphology, and size. Willow growth and size were optimized when elk consumed approximately 21% of the biomass and was negatively effected when elk consumed more than 37% of the biomass. Optimal willow growth refers to the maximum observed willow production in the presence of elk herbivory, which was greater than ungrazed willows. Negative willow growth represents a 40% decline in most components of willow growth from maximum growth levels. Elk densities ≥ 32 elk/km² generally resulted in a 40% decline in willow size and growth parameters. However, the relationship between willow consumption rates, elk densities, and willow growth varies greatly across the winter range (e.g., Peinetti et al. 2002, Singer et al. 2002, Zeigenfuss et al. 2002) and meaningful increases in willow growth may require elk densities lower than 15 elk/km². Increases in willow growth at or below 30 elk/km² are not highly correlated and difficult to ascertain from Singer et al. (2002) due to a lack of data for willow stands with 10-30 elk/km². Willow growth is also site specific; Peinetti et al. (2001) and Zeigenfuss et al. (2002) found differences in willow growth rates between Moraine Park and Horseshoe Park.

The effects of elk on willow production (i.e., new biomass grown by the plant each year) have not been clearly shown. Comparing browsed and unbrowsed plants in exclosures,

Zeigenfuss et al. (2002) found elk significantly reduced willow production after four years. Peinetti et al. (2001) did not find this; browsed plants were smaller, but similar in overall biomass production because they produced longer, thicker shoots in the lower levels of the plant canopy. The discrepancy between these studies is likely due to differences in the time of study and methods utilized. More importantly, both studies clearly found elk herbivory to greatly reduce willow size and vigor (Peinetti et al. 2001, Zeigenfuss et al. 2002).

Shallower water tables (i.e., closer to ground level and more beneficial to the roots of a plant) were found to have a positive correlation to willow size, but the effect was minimal when compared to elk browsing effects (Singer et al. 2002, Zeigenfuss et al. 2002).⁷ In general, water tables near streams on the elk winter range do not fall below one meter during the growing season (Zeigenfuss et al. 2002, Cooper et al. 2003). The roots of established willow plants (at least 3-4 years old) should be able to easily reach these ground water depths. However, Menezes et al. (2002) suggested that, in non stream-side plants, the level of elk browsing during the mid to late 1990s was decreasing the ability of willow roots to reach groundwater sources (note: this was based on isotopic differences between stream-side and non stream-side water signatures that were considered significantly different by the author based on P-value of 0.099). Menezes et al. (2002) and Peinetti et al. (2001) suggest that this is due to willows on the winter range overcompensating C to their aboveground willow parts (stems, leaves) in response to high levels of herbivory, resulting in less C allocation to the belowground plant components (roots) and a decreased ability of the plants to reach deeper water sources. Stream-side plants were still able to reach groundwater because they were located in areas with more shallow water tables (Menezes et al. 2002). However, the relative effects of elk and water are exhibited by the finding that many stream side plants with excellent water availability are in poor condition due to the effects of elk browsing (Peinetti et al. 2002, Zeigenfuss et al. 2002, Cooper et al. 2003).

Willow Reproduction in RMNP – Cooper et al. (2003) found that willow seeds are currently the principal method of willow establishment on elk winter range areas in RMNP, with little to no reproduction occurring via willow stem fragments. There are currently very few beaver on the elk winter range and it is possible that greater population numbers could increase the role of vegetative establishment. It is also notable that two of the three landforms that create suitable seed sites in RMNP are created by beaver activity (abandoned beaver ponds and abandoned channels or ox-bows). An increase in beaver would likely increase the reproductive capabilities of willow either directly through vegetative propagation or indirectly through creating appropriate sites for seedling establishment.

Even though seedling establishment is currently the principal method of willow reproduction on the elk winter range, it is severely inhibited by a lack of seed rain and seedling survival, primarily due to the effects of herbivory (Gage and Cooper 2005). Even when seedlings have

⁷ Note that the previous study by Peinetti et al. (2002), which suggested declines in surface water are correlated with declines in willow distribution, examined the entire winter range. The studies in this section (Singer et al. 2002, Zeigenfuss et al. 2002) examined existing willow plants near streams. Thus, these areas have not been largely affected by the decline in beaver because water tables remain elevated throughout the year (due to their proximity to water sources).

been established in recent decades, few to no willow plants have grown beyond a meter (~3 feet) (Peinetti et al. 2002, Zeigenfuss et al. 2002, Cooper et al. 2003). Cooper et al. (2003) found no relationship between willow age and height, and thus small, single willow stems that appear to be newly established or young plants cannot be assumed to be recent colonization (as suggested by Peinetti et al. 2002).

The SAVANNA model predicted willow cover on the winter range in the park to be almost three times greater prior to European American settlement or undisturbed conditions (higher water tables, with wolves) versus current conditions (lower water tables, no wolves, greater levels of herbivory). Undisturbed willow cover was found to be similar to the maximum amount of potential willow cover (no elk or deer). These results suggest willow cover did not exhibit an increase when elk were extirpated, and that initial declines in willow likely started when areas such as Moraine Park were developed (Gysel 1960). Willow declines continued over the last 50-60 years due to greater foraging pressure from elk and loss of the areas beaver population. These declines and complete losses of willow riparian areas on the core winter range are expected to continue until elk population size or density is reduced in willow riparian areas on the core winter range (Gage and Cooper 2005).

Willow cover has also been found to be highly dependent on water table levels. The model assumed that, primarily due to the loss of beaver, current water table conditions are lower than they were historically. Under such conditions, even if the elk population were reduced and all willow riparian areas were fenced off, willow would only be able to regain half of its (pre-1860) level of cover (Coughenour 2002). This indicates a need for a diversity of management actions to attain an even partial recovery of willow on core winter range areas, a conclusion supported by several empirical studies conducted in RMNP (Baker et al. 2004, Baker et al. 2005, Gage and Cooper 2005).

Bitterbrush and Sagebrush

Bitterbrush and Sagebrush Ecology – Bitterbrush and sagebrush are found on arid plains and slopes throughout the Rocky Mountains. Natural disturbances such as fire and browsing can greatly affect the growth and reproduction of both shrubs. Bitterbrush has a greater tolerance to fire because it can resprout from previously existing plants if the fire is not too intense (Blaisdell and Mueggler 1956). Conversely, sagebrush plants are easily killed by fire and can have trouble re-establishing themselves because they only reproduce by seed (Kershaw et al. 1998, Wambolt 1998, Wambolt et al. 2001). Recovery rates to previous levels for both species following fire can range from just a few years resulting in a rejuvenated shrub community to over 30 years (Bunting et al. 1985, Wambolt et al. 2001).

Bitterbrush is also more resistant to browsing than sagebrush due to plasticity in growth rates and resource allocation traits (Bilbrough and Richards 1993). Both plant species are very important diet components for deer in RMNP, and to a lesser extent elk (bitterbrush makes up 3-6% of elk diet), because they maintain higher protein levels than grasses (Kufeld et al.

1973, Stevens 1980a, Hobbs et al. 1981).⁸ If large ungulate browsing follows fire events, it can often result in long-term or permanent shrub loss (Wambolt et al. 2001). Small mammals have also been found to feed extensively on bitterbrush seeds (Martin et al. 1951) and can have a large, detrimental effect on bitterbrush recruitment (Clements and Young 1996).

Bitterbrush and Sagebrush in RMNP – The distribution of bitterbrush and sagebrush on the winter range at the time of settlement is unknown. By the time shrub communities in RMNP were first studied in the 1930s they had been significantly altered. Market hunting during the 1860s and 70s significantly reduced elk and deer numbers, likely allowing upland shrubs to expand, but fires in ponderosa pine forests during the 19th century (Veblen et al. 2000, Ehle 2001) and an expanding deer population in the early 1900s (Stevens 1980a) may have also greatly decreased shrub abundance.

By the 1930s, managers in RMNP concluded that the deer population was having a large, negative effect on upland shrubs (Grater 1945). Deer started to decline in the 1940s, and this was further facilitated by National Park Service reduction programs. Gysel (1960) suggested that sagebrush continued to decline even as deer were reduced. No quantitative data was ever collected on upland shrubs so even less is known about any changes in bitterbrush production or distribution.

Zeigenfuss et al. (2002) did not find herbivory by elk and mule deer to affect bitterbrush production or size through four years of study. Resprouting of bitterbrush was observed in all sites after prescribed fire treatments. Fire reduced bitterbrush size in the first and second year after burning and production in the second year, suggesting additional plant mortality took place during the first year after the fires (Zeigenfuss et al. 2002). Nesvacil and Olmsted (2003) found that bitterbrush canopy area, volume, and estimated annual production remained low six and seven years post-burn, while total shrub canopy area, volume and annual production only differed due to grazing ($P < 0.10$). This suggests that bitterbrush can resprout adequately after burning, but the current levels of herbivory in RMNP are impeding regeneration and growth (Nesvacil and Olmsted 2003).

Schoenecker et al. (2002) found upland shrub production, primarily sagebrush, to be 67% greater inside versus outside 35 year-old exclosures. Similarly, grass biomass was reduced by 32% outside upland shrub exclosures.

Herbaceous Communities on the Winter Range

Up to 70% of an elk's diet in RMNP during winter is graminoids (Hobbs et al. 1981, Singer et al. 2002). However, the effects of elk on herbaceous production on the core winter range are equivocal. Zeigenfuss et al. (2002) observed herbaceous production in willow communities to be significantly reduced (18-29%) in 1998 (the last year of their study); but they also observed herbaceous production was higher in grazed versus ungrazed sites in 1994, 1995, and 1996 (this was statistically significant in 1994 and 1995, $P < 0.10$). The difference

⁸ Upland shrubs are particularly important to deer because they are the majority (up to 75%) of their diet (Kufeld 1973). See the Ungulate section for more information.

between 1998 and the other years could be due to the fact that all other previous years had above average precipitation. Studies in Yellowstone have found that elk only affect herbaceous communities in years of drought (Singer et al. 1998b). Thus, elk may have greater effects in RMNP when precipitation is average or below-average.

Annual herbaceous consumption rates averaged 55% in willow communities and 60% in upland shrub communities on the core winter range. The majority of consumption occurred during summer in willow areas and winter in upland areas (Singer et al. 2002)⁹. Singer et al. (2002) reviewed herbaceous plant consumption rates for a variety of ecosystem types around the world. Mountain bunchgrass steppe and mixed grass prairie communities, the most comparative systems to RMNP, were able to withstand consumption rates of 40%, but not 60%. This indicates the consumption rates in RMNP may be large enough to alter herbaceous communities in RMNP (Singer et al. 2002). Herbaceous plants in willow communities are particularly vulnerable because the majority of grazing is occurring during the growing season (Augustine and McNaughton 1998).

Recent experimental studies have documented a number of individual plant species that are being affected by herbivory. In willow riparian areas, Zeigenfuss et al. (2002) found grazed sites had more goldenrod species (*Solidago* spp.) and exclosed sites (ungrazed) had more bluebell (*Mertensia ciliata*) after four years. There was also more bare ground (4.6% greater) and grass cover (10.8% greater) on grazed versus ungrazed willow sites, but these differences were not statistically significant (Singer et al. 2002). In upland bitterbrush sites, Zeigenfuss et al. (2002) found ungrazed sites had a higher percent cover of prairie sage (*Artemisia ludoviciana*) and sulphur buckwheat (*Eriogonum umbellatum*) after four years. In 35 year-old exclosure sites, which covered a variety of habitat types, grazed areas had 6.4% more bare ground than ungrazed areas, but differences were not statistically significant (Singer et al. 2002).

Using David R. Steven's long-term data set, Zeigenfuss et al. (1999) found the non-native grass *Phleum pratense* (known as timothy) increased significantly (by 54%) in meadow areas on the winter range from 1968-1992. *Phleum commutatum* is highly palatable to elk, very productive, and has the ability to take over large areas of land because it is resilient to grazing and other types of disturbance. Other significant changes over this time include an increase in the coverage of forbs, *Carex* spp., lichens, and mosses, and bare ground in grassland areas.

There is little other evidence to suggest that non-native or invasive plant species are being greatly facilitated by elk grazing. Studies by Singer et al. (2002) and Zeigenfuss et al. (2002) found few significant differences in plant species composition due to herbivory in willow riparian and upland bitterbrush habitat types. Schell and Stohlgren (1997) and Stohlgren et al. (1999) examined plots inside and outside of several 35 year-old exclosures on the elk winter range in RMNP (and other areas) and did not find grazing to accelerate the spread of exotic plant species. They concluded soil quality, climate, and disturbance have much greater

⁹Note: During certain times of the year, especially the spring, deer also consume herbaceous plants. However, they are primarily browsers (Kufeld et al. 1973) and are not likely having a large effect on herbaceous communities.

effects on plant species composition than grazing. Stohlgren et al. (1999) found that elk, even when present at a very high density, reduce non-native plant species coverage. However, this study was conducted at a much larger scale than RMNP and it is difficult to ascertain if such conclusions hold true for an elk population at extremely high densities (>100 elk/km²).

Coniferous Forest Communities in RMNP

There are three major forest communities in RMNP: 1) ponderosa pine, with some Douglas fir and lodgepole pine, found only on the east side of the park ranging from approximately 2,450 to 2,800 m (~8,000 to 9,200 feet) in elevation; 2) lodgepole pine ranging from 2,500 to 3,100 m (8,200 to 10,200 feet); and 3) subalpine forests, consisting of Engelmann spruce (*Picea engelmannii*) and/or subalpine fir (*Abies lasiocarpa*), ranging from 2,800 to 3,500 m (~8,000 to 11,500 feet) (Fig. 2). The latter two, higher elevation forest types, are primarily affected by fire and climate and are generally considered to be within their natural range of variability (Sherriff et al. 2001). Elk populations have a minimal effect on these lodgepole and subalpine forests as they are not an important food source in RMNP (Hobbs et al. 1981, Larkins 1997, Singer et al. 2002).

Ponderosa pine forests make up a large proportion of the elk winter range in RMNP. Fire has historically played a major role in the maintenance of healthy ponderosa pine forests throughout the Rocky Mountains (Arno 1980). Periodic fires - both lightning caused and those set by Native Americans - were believed to be the major factor underlying the mosaic of forest ages and densities that were typical of ponderosa pine communities in RMNP prior to the arrival of European Americans.

Fire return intervals for ponderosa pine forests in RMNP vary widely, averaging between 20 and 60 years but ranging from as little as one to as much as 300 years (Rowdabaugh 1978, Mast et al. 1998, Ehle 2001). Ehle (2001) found small fires to occur every one to 120 years and large fires to occur every 200 to 300 years. Large fire events have a longer time interval because the trees must attain a certain size and density before they can support crown fires that spread over a large area. Ehle (2001) concluded that ponderosa pine communities in RMNP historically underwent large fire events that cause extensive tree mortality and led (directly or indirectly) to regeneration, followed by small surface fires that caused small patches of tree turnover and created a heterogeneous landscape over time. This is consistent with the findings of Veblen et al. (2000), who examined upper elevation ponderosa pine forests adjacent to RMNP. The fire regime suggested by Veblen et al. (2000) and Ehle (2001) would have resulted in a wide variety of stand ages and densities in the ponderosa pine forests in and around RMNP. This contrasts with low elevation ponderosa pine forests along the Colorado Front Range and in the southwest U.S., where forests were maintained in an open park-like forest structure due to frequent low-intensity surface fires that affect a large proportion of the forest (Veblen et al. 2000).

Past fire scars from ponderosa pine forests in and adjacent to RMNP appear to show an increase in fire activity in the 1800s largely due to logging activities and other human

ignitions (Rowdabaugh 1978, Mast et al. 1998). Because of these activities, we cannot be sure that pictures taken at this time reflect the conditions that existed prior to European American settlement (Clements 1910, Veblen and Lorenz 1991). Fire increases during the 19th century followed by the fire suppression efforts of the 20th century have generally worked in combination to create many dense, single aged stands (Veblen et al. 2000). Ehle (2001) concludes that a more heterogeneous forest would more closely resemble the forests found prior to European American settlement. However, long periods of time without small or large fires may also be possible in RMNP and dense stands are not necessarily unnatural (Veblen et al. 2000, Ehle 2001).

Although ponderosa pine forests make up a large proportion of the elk winter range in RMNP, dense even-aged stands provide little forage for elk (Hobbs et al. 1981, Singer et al. 2002) and likely have little effect on elk populations. More heterogeneous ponderosa pine savannahs would likely produce additional forage for elk, but it is difficult to quantify the effect they would have. It has been suggested that fire suppression is allowing ponderosa pine trees to expand into and overtake some of the meadows that make up the core of the elk winter range and their primary forage base. This could potentially lead to a significant decrease in elk forage, but no formal research has been conducted on this subject. Historic and current photos do show that stands of ponderosa pine are more dense, and it may appear that they now occupy former meadows. However, the pure grassland areas (e.g., Moraine Park, Horseshoe Park, Beaver Meadows) that make up the primary feeding areas for elk on the winter range have not been encroached on by trees (Veblen and Lorenz 1991).

Alpine Tundra

A complete description of plant types found in alpine tundra areas can be found above and in Appendix 1. Very little work has been done on the effects of elk on plant species composition in alpine tundra areas. Zeigenfuss et al. (1999) found willow declined in height and coverage from 1968 to 1992, but herbaceous coverage remained stable. Stevens (1980b) reported that all the major cover species, except curly sedge (*Carex rupestris*), decreased in cover and frequency in the 1970s. It cannot be determined if such changes are due to elk because ungrazed areas were not available for comparison. However, Singer et al. (2002) recently found that willow can constitute over 50% of an elk's diet during the summer in the alpine areas of RMNP, suggesting that the effects of elk in alpine areas need to be further explored.

Summary

It is clear that elk do not affect the majority (>95%) of aspen in RMNP. Persistence of aspen throughout the park is primarily dependent on fire or other disturbances that open up coniferous forest canopy and allow aspen stands to recruit new trees. On the core winter range, however, elk are the proximate factor preventing new aspen tree establishment. There is no definitive evidence that aspen are a long-term feature of the core winter range, but research conducted elsewhere indicates that aspen stands in the western U.S. have been present for hundreds of years. Under current levels of herbivory, all of the research indicates

that aspen stands on the core winter range will be extirpated or reduced to a shrub-like state. There remains uncertainty as to what population size or density of elk is compatible with aspen regeneration and growth on the core winter range. At a minimum, the number of elk on the core winter range will need to be less than 600. Modeling indicates the density of elk will need to be reduced to $\leq 4\text{--}15\text{ elk/km}^2$ to achieve any tree recruitment on the core winter range.

Willow distribution on the core winter range of RMNP has been negatively affected since the area was settled. Recent declines are attributed to reductions in surface water, due to a continual decline of the area's beaver population; and high levels of browsing by an elk population at ecological carrying capacity. New willow are unable to be established due to the limited distribution of mature seed producing plants, lack of suitable seed sites, and high levels of elk herbivory. Vegetative reproduction is severely limited by too few beaver in the area. Established willow plants cannot survive or grow to produce seeds due to the levels of elk browsing. Thus, as the remaining tall, seed-producing plants are cut down by beaver, elk suppress their growth and lead to further willow and beaver declines. At a minimum, elk densities need to be reduced from current levels of over 100 elk/km^2 in some areas to $\leq 15\text{--}30\text{ elk/km}^2$ to allow willow to grow beyond the reach of elk and become viable, seed producing plants. This alone, however, is not enough to ensure willow reproduction or expansion on the willow range; beaver activities and surface water flows also need to be restored to create appropriate seed and vegetative reproduction sites and facilitate willow growth.

Herbivory by elk and deer has not been found to have a large, negative impact on mature, existing bitterbrush and sagebrush plants. However, in combination with the current levels of herbivory, fire can have severe, negative effects on both species. Sagebrush is intolerant of fire regardless of subsequent browsing levels. Bitterbrush can adequately resprout following a fire; however, fire will have a large, negative impact on bitterbrush until browsing pressures are alleviated. The relative effects of elk and deer on upland shrubs on the winter range are unknown.

Herbaceous consumption rates by elk on the winter range in RMNP are extremely high and exceed many areas in North America. Research to date indicates that this has not led to a widespread, increased incidence of invasive species in herbaceous communities on the winter range. However, the large increase in the non-native timothy should be a concern to managers. Several other trends have also been attributed to elk grazing: prairie sage, sulphur buckwheat, and bluebell have decreased; and goldenrod species and bare ground have increased.

Elk populations have a minimal effect on the coniferous forests of RMNP. None of the trends in alpine tundra areas can be directly attributed to elk due to the lack of experimental data; however, the high percentage of willow in the diet of elk during summer in alpine areas and negative correlation between increased elk numbers and decreased willow height and coverage is reason for concern and should be further studied.

UNGULATE INTERACTIONS

Mule deer, bighorn sheep, and moose also inhabit RMNP. These ungulates are spatially segregated from each other for most of the year: moose primarily occur on the west side of the park during the summer in the vicinity of the Kawuneeche Valley; mule deer occur throughout the park in summer, but in winter are only found on the east side of the park near or overlapping with elk winter range areas; and bighorn sheep are found in several mid to high elevation areas throughout the course of the year (Fig. 10). Elk considerably outnumber these ungulates in and near the park (at a ratio of 100 elk: 17 deer: 17 bighorn sheep: 1 moose) and overlap extensively with all of them due to their distribution, population size, and large seasonal migrations (Figs. 3, 7). This section reviews the status of mule deer, bighorn sheep, and moose in the park and discusses how elk interact and potentially compete with these ungulates.

Ungulates may compete for forage if two species overlap in diet and range use, and if one or both of their levels of consumption are large enough to limit available forage (e.g., Hobbs et al. 1996a, Hobbs et al. 1996b). If competitive effects are strong enough and favor a particular ungulate at the expense of another, this can affect the production, distribution, or population size of another species (Hobbs et al. 1996a, Hobbs et al. 1996b, Forsyth and Hickling 1997). Such interspecific competition for resources may be entirely natural and appropriate in some ungulate grazing systems. However, if a competitively dominant species becomes overabundant, it could lead to changes considered to be outside the natural range of variation.

Mule Deer

The number of mule deer that inhabited the RMNP area before 1915 is not known, but it is clear that mule deer were heavily hunted and suffered large declines throughout Colorado and the RMNP area during the late 1800s and early 1900s (Stevens 1980a, Colorado Division of Wildlife 1999). Colorado responded to these declines by closing the state to hunting in 1913. With the creation of RMNP, which also instituted programs to control predators and stop poaching, deer numbers in the park increased to as many as 1,400 in the winter of 1938 (Fig. 5; Stevens 1980a).¹⁰

The number of deer on the east side of RMNP appeared to start declining in 1939 or 1940, declines that were further facilitated by the NPS deer and elk reduction program from 1943 to 1968. Deer were reduced by NPS to “improve range conditions” on the east side of the park. Deer numbers appeared to continue to decline even after reductions were terminated in 1968 (Fig. 5). This is not unique to the RMNP area, as the entire western U.S. has observed a continual decrease in mule deer numbers since at least the 1970s (Colorado Division of Wildlife 1999). Estimates conducted from 2001 to 2003 during winter indicate the mule deer

¹⁰ Stevens (1980a) notes that there is no way to determine the accuracy of this estimate because there is no documentation on how the surveys were conducted. However, it is clear there were many more deer during the 1930s than the late 1800s and early 1900s.




Figure 10. Primary areas of use for moose, bighorn sheep, and mule deer in Rocky Mountain National Park (RMNP).

population in the Estes Valley decreased from 833 (95% C.I.; 659-1,048) to 561 (498-632) (Fig. 5; Conner 2004). Recent population estimates from 2004 and 2005 indicate deer have not declined further (M. Watry personal communication).

Competition with Elk – Kufeld et al. (1973) summarized the diet of mule deer in the Rocky Mountains and found it is largely composed of browse (shrubs and trees) and forbs throughout the year, with a low to moderate use (~20%) of graminoids (grasses) in the spring. In the winter, deer must rely heavily on browse and cannot subsist on dry, senescent grasses (Colorado Division of Wildlife 1999). Conversely, the diet of elk in RMNP can encompass a wide variety of species in a wide variety of habitats, and they can greatly increase their intake of browse species when grasses are unavailable or the nutritional content of grasses becomes less than browse (Hobbs et al. 1981). Such conditions commonly occur during winter, a drought, or when an ungulate reaches its food limited carrying capacity (Hobbs et al. 1981, Forsyth and Hickling 1997). This is important because it is during these periods when deer exhibit their greatest need for browse (Kufeld et al. 1973). Elk are also larger than deer and can displace deer from preferred feeding areas (Colorado Division of Wildlife 1999).

Because elk are competitively dominant, at carrying capacity on the core winter range, and overlap considerably with mule deer during winter (Figs. 3, 10), it is likely that elk are having a negative impact on mule deer production and population size (Colorado Division of Wildlife 1999; see also Hobbs et al. 1996a, Hobbs et al. 1996b, Forsyth and Hickling 1997). This is supported by Coughenour (2002) and the observation that deer numbers and use continued to decline while elk numbers and use increased in the best deer habitats (upland shrub) after the RMNP control program was terminated (Zeigenfuss et al. 1999). In general, elk and deer population estimates over the last 50-75 years from RMNP are inversely correlated (Figs. 4, 5, 8).¹¹

The SAVANNA model indicated that under natural conditions and with wolves present in RMNP, deer populations on elk winter range areas would be limited to approximately 200 animals. The model also predicts that without wolves, deer would be limited to a similar population size through competition with elk (Coughenour 2002). However, the latter scenario (limitation by competition with elk) results in deer that are not as healthy as a predator limited population.

Chronic Wasting Disease – Live deer can be tested for the disease by taking a tonsillar biopsy in the field and analyzing the sample in a lab. This has allowed RMNP and the Colorado Division of Wildlife to determine that prevalence of the disease inside and adjacent to the park is 5.4% (Conner 2004).

Simulation modeling predicts CWD has the potential to cause drastic population reductions in deer (Miller et al. 2000, Gross and Miller 2001). However, this does not mean the

¹¹ It is important to note that the initial decline in the mule deer population size in the late 1930s and early 1940s is believed to be due to an overpopulation of mule deer reducing their own forage base (Stevens 1980a). However, continued declines throughout the 20th century in conjunction with an increasing elk herd suggest that elk may now be competing with and affecting mule deer.

population declines in mule deer over the last 50-70 years are due to CWD, as mule deer have declined throughout Colorado and the western U.S., including areas where CWD have not been documented (Colorado Division of Wildlife 1999).

Bighorn Sheep

Prior to 1880 bighorn sheep were reported to be abundant and occupy a much greater area in RMNP than they currently do (Ratcliff 1941*b*, Packard 1939). Ratcliff (1941*b*) suggested there may have been as many as 4000 sheep in the park during the early 1800s, but there is no way to assess the accuracy of this estimate. Initial population declines began in the late 1800s due to market hunting and continued sporadically through the 1950s due to a number of factors, including competition with livestock, disease, and development (Packard 1939, Packard 1946, Goodson 1978).

Bighorn populations were apparently stable or increasing from the 1960s to the 1980s. Bighorn population size on the west-side (Never Summer and Continental Divide areas) was estimated to be 124 in 1975-76 and between 452-502 in the late 1980s. Population size on the east-side (i.e., Mummy Range) was estimated to be 81 in 1975-76 and 200 in the late 1980s (Bailey 1990, Goodson and Stevens 1991).

There is anecdotal evidence of pneumonia induced dieoffs occurring in both herds during the mid 1990s, and recent research and park observations support this assertion. West- and east-side bighorn populations in 2003 were estimated to be 289 and 78, respectively, and both herds tested positive for multiple pathogens believed to be related to sheep dieoffs due to disease (McClintock 2004). In addition, a young ram from the Mummy Range herd was observed coughing and later found dead due to pneumonia in summer 2001, and low lamb counts have occurred since the mid 1990s (RMNP unpublished data). Such findings are typical of herds undergoing or previously affected by pneumonia epizootics, which are considered the greatest threat to bighorn sheep populations in North America (Bailey 1986; Coggins 1988; Ryder et al. 1992; Cassirer et al. 1996; Jorgenson et al. 1997, Monello et al. 2001).

Competition with Elk – Bighorn sheep are primarily found at higher elevations in the park (Figs. 2, 10) and elk only use these areas during the summer and fall (Figs. 3, 7). Capp (1967) and Harrington (1978) examined summer range use by elk and bighorn sheep in alpine areas of RMNP and found bighorns and elk were spatially segregated: bighorns tended to use steep, rocky slopes while elk used open meadow areas. Capp (1967) and Singer et al. (2002) found elk in alpine areas to primarily consume forb and browse (*Salix* spp.) species in the summer, while bighorns primarily consume grasses (Capp 1968). Similarly, Singer and Norland (1994) found no evidence that elk have had any effects on bighorn sheep populations in Yellowstone since the onset of natural regulation.

Competition in RMNP cannot be completely ruled out as the population size and range use of elk has increased considerably since the study periods of Capp (1967) and Harrington (1978), sheep and elk diets overlap (Singer and Norland 1994), and elk are capable of displacing

sheep from preferred feeding areas (Goodson 1978). Other unevaluated factors, such as increased forest density due to fire suppression, may also be contributing to loss of habitat and lower bighorn numbers. However, competition with elk and other unevaluated factors are likely of little importance compared to disease issues.

Moose

Moose in Colorado are at the southern limit of their distribution in the U.S. Historically, moose were not common in Colorado or RMNP. There is only one recorded historic occurrence of a moose in RMNP; Estes (1939) reported killing a moose in Moraine Park in 1860. In 1978-79, the Colorado Division of Wildlife introduced 24 moose to the North Park area, located about 30 km west of RMNP, in order to establish a viable resident moose population for hunting purposes. The estimated population size reached 100 by 1986, and a moose was first observed in RMNP in 1980, two years after the first animals were released. Since then, moose observations in RMNP have continued to increase, with the first winter observations in 1985. Currently, the North Park population is estimated at 600 moose.

Moose primarily use RMNP and overlap with elk populations during summer, with an estimated 100-120 animals using areas in or near the Kawuneeche Valley. Only about 60-70 of these moose stay all summer. Almost all moose leave the Kawuneeche Valley for North Park (to the W of RMNP) by the end of September each year, although a few animals do remain in the Kawuneeche Valley area during winter. Moose use the North Park area because it is at a lower elevation, has warmer weather, and better forage conditions than the Kawuneeche Valley area (J. Dungan personal communication).

While moose and elk may compete for forage in summer, most studies have found that moose avoid competition via spatial separation. A study in the Gallatin River Valley in Montana found moose distribution was primarily above and below the major elk distributions and indicated that moose were not tolerant of large groups of elk (Peek and Lovaas 1968). Jenkins and Wright (1987) found that winter moose and elk distribution was mediated in part by snow depth, with moose being able to use areas with deeper snow than elk in Glacier National Park, Montana.

Studies conducted in other areas suggest bears and wolves may limit the size and density of moose populations and their localized effects on vegetation (Bergerud et al. 1983, Gasaway et al. 1983, Ballard et al. 1987, Boutin 1992, McLaren and Peterson 1994, Peterson 1999; but see Boutin 1992). The absence of wolves in RMNP could result in higher densities of moose with greater effects on the plant community.

Summary

Mule deer numbers in the Estes Valley have declined dramatically since the 1940s. Currently 500-600 deer inhabit the area. Several lines of evidence suggest that elk are negatively affecting deer through interspecific competition: elk are competitively dominant; at carrying capacity and thus likely having an affect on winter browse species relied on by deer; and,

after ungulate reductions ended, deer numbers continued to decline while elk numbers increased. SAVANNA also indicates deer populations in RMNP are suppressed by high elk numbers via interspecific competition.

No research suggests that bighorn sheep and moose compete with elk at a level that has population consequences. Disease is considered the most important factor influencing bighorn sheep population dynamics in North America and RMNP. Moose primarily use the Kawuneeche Valley area in summer and leave during winter. There is little evidence to suggest they compete with elk in RMNP, but moose numbers would likely be lower if wolves were present.

EFFECTS OF ELK ON BIODIVERSITY

This section outlines the most likely effects elk could be having on the diversity, distribution, or population dynamics of animals or plants in the park. It focuses on birds, plants, butterflies, and beaver because these groups and their interactions with elk have been studied in RMNP. Effects on these species are usually indirect, in that elk may influence them through competition for resources or alteration of habitat.

Birds

Over 250 bird species have been observed in the RMNP area. The large majority of these birds are seasonal residents or migrants; only 26 species are considered to be common, year-round inhabitants of the park. The most common birds observed in the park can generally be split into four categories; ptarmigan (a grouse species), songbirds and neotropical migrants, raptors, and waterfowl. This report only address the first two groups because data is available to evaluate potential relationships with elk and they are most likely to be affected by elk herbivory effects.

Ptarmigan – White-tailed ptarmigan (*Lagopus leucurus*) are a member of the grouse family (Tetraonidae) and are known for their seasonal changes in plumage which vary from pure white in the winter to a mottled brown in the summer. White-tailed ptarmigan (hereafter ptarmigan) in Colorado primarily inhabit elevations above treeline and during the winter are only found in areas where willow is a dominant or co-dominant plant species. Their diet is dominated by willow buds, leaves, and twigs from October to June, but is more diverse during the summer and includes herbaceous and browse species (May and Braun 1972). It has been suggested that the population size of grouse species inhabiting arctic or subarctic regions fluctuate in a cyclic fashion over the course of 7-10 years (Williams 1954), but this has not been definitively shown for white-tailed ptarmigan (Choate 1963, Braun et al. 1991).

White-tailed ptarmigans have been studied since 1965 at two alpine study sites adjacent to Trail Ridge Road in RMNP. During the 1970s, combined population numbers for these areas fluctuated but remained above 100 birds. Since the early-to-mid 1980s yearly fluctuations have continued, but the overall population declined to approximately 50-60 birds by 1999 (Braun et al. 1991, K. Giesen and C. Braun unpublished data). This decline was concurrent

with a decline in willow cover and an increase in elk population size, prompting Braun et al. (1991) to suggest “heavy use of willow by elk in early winter and early spring constrains ptarmigan breeding densities by reducing [the] amount of food (willow buds) available to ptarmigan in late winter during the early breeding period when ptarmigan establish [breeding] territories”. Conversely, Wang et al. (2002) found no relationship between elk population size and ptarmigan population growth rates. They did find a significant relationship between weather and ptarmigan, suggesting periodic increases in winter temperature over the last two decades have been negatively associated with ptarmigan population growth rate. However, the data presented in Braun et al. (1991) and Wang et al. (2002) is correlational and cannot be used to infer cause and effect relationships between elk, willow, and ptarmigan. More research is needed to determine if elk, climate, or both factors are influencing ptarmigan populations.

Songbirds – At least 150 songbirds have been observed in the park. About 1/3 of these are neotropical migrants, defined as birds that spend their winter south of the United States or Tropic of Cancer. This classification and definition is taken from Connor (1993), who included the neotropical migrant lists of both Colorado Partners in Flight and Cunningham (1988). Neither list includes raptors such as the peregrine falcon (*Falco peregrinus*), even though they winter in the tropics. Many neotropical migrants and songbirds also breed in RMNP (Johnsgard 1986).

Songbirds and neotropical migrant diversity is greatest in aspen, riparian willow, and ponderosa pine habitat in RMNP (Connor 1993, Turchi et al. 1994), which combined make up 9% of the total park area. The core elk winter range areas in the park are very important for songbirds because it contains the majority (55%) of these habitat types. Within RMNP, the only other large, continuous portions of such habitat is in the Kawuneeche Valley, which contains 9% of the park’s aspen and willow riparian habitat (source: RMNP GIS database).

Turchi et al. (1994) found bird species richness to be significantly higher in aspen than conifer habitat, and percent shrub cover (0.5-2 m or 1.5 to 7 feet in height) within aspen stands to be the single most important predictor variable for bird species richness in RMNP. Lloyd (1997) surveyed bird species present in aspen stands on and off the winter range, and found the mountain chickadee (*Poecile gambeli*) to be more abundant in stands located off the winter range. He attributed this to the fact that mountain chickadees tend to inhabit coniferous forests (Johnsgard 1986), which are much more abundant around aspen stands off versus on the winter range. Zaninelli and Leukering (1998) and Duberstein (2001) suggested live aspen trees are more important to cavity nesting birds than dead trees.

Several studies have suggested that different bird species use different sizes and densities of willow and aspen; i.e., short and tall willow are both important (Zaninelli and Leukering 1998, Leukering and Carter 1999, Duberstein 2001). Anderson (2002) found a negative relationship between elk use and understory volume of aspen stands and a positive relationship between understory volume and bird abundance in the Jackson Hole area. In willow stands, the abundance of several willow-dependent bird species was reduced in willow stands that were proximate to elk feeding areas. Anderson (2002) attributed these

differences to the effects of elk herbivory. Willow-dependent bird species negatively affected by proximity to elk herbivory included willow flycatchers (*Empidonax traillii*), yellow warblers (*Dendroica petechia*), MacGillivray's warblers (*Oporornis tolmiei*), fox sparrows (*Passerella iliaca*), and song sparrows (*Melospiza melodia*). Thus, a localized conversion to short willow in core winter range areas will likely result in localized decreases of bird abundance and biodiversity.

The inability of upland shrubs to regenerate following fire due to elk herbivory has been found to negatively affect green-tailed towhee (*Pipilo chlorurus*) nest survival (Jehle 2004).

Butterflies

Simonson et al. (2001) examined butterfly diversity in six different habitat types on the elk winter range. They found butterfly diversity, richness, and uniqueness was highest in aspen and wet meadow habitat types. The aspen habitat included two species listed as rare by the state of Colorado. Other habitat types included in the study were dry meadow, ponderosa pine, lodgepole pine, and burned ponderosa pine. Interestingly, Simonson et al. (2001) found butterfly species richness exhibited a strong correlation to native as well as non-native plant species richness. The decreased coverage of sulphurflower in grazed sites (Singer et al. 2002) is a concern because it is an important host plant for a variety of butterfly species (R. Bray personal communication).

Beaver

Beaver Ecology – Beaver are a keystone species that have profound effects on ecosystem structure and function (Naiman et al. 1988). Beaver modify their environment by cutting aspen and willow for food and construction material, and building dams that raise the water table, trap sediment, and increase nitrogen availability to willow (Naiman et al. 1988, Baker and Hill 2003). Beaver dams also slow current velocity, increase deposition and retention of sediment and organic matter in their ponds, reduce downstream turbidity on floodplains, increase the area of soil-water interface, elevate the water table, change the annual stream discharge rate by retaining precipitation runoff during high flows and slowly releasing it during low flows, alter stream gradients by creating a stair-step profile, and increase resistance to disturbance (Naiman et al. 1988). Beaver foraging can alter species composition, density, growth, and distribution of woody vegetation.

Willow species provide an important food source for beaver. Willow leaves are high in protein content and are readily eaten during the summer. The bark of willow stems may be the only source of winter food for beaver that live in locations where surface water freezes during winter (Baker and Cade 1995). Beaver are central place foragers that cut and remove entire stems at or near the ground surface. They often cut all stems from preferred shrubs growing near their winter food caches, dams, and lodges, but become more selective as foraging distances increase (Baker and Hill 2003).

Beaver cutting stimulates sprouting from below the cut. Beaver affect willow growth and height in the intermountain west along with ungulate browsing (Singer et al. 1994) and site conditions such as soil type, length of growing season, nutrient concentrations, and water table height (Cottrell 1995, Peinetti 2000).

Beaver in RMNP – Beaver populations have declined dramatically in RMNP since 1940. While multiple factors have likely contributed to beaver declines, the most significant cause appears to be trapping that occurred in 1941 through 1949 (Baker et al. 2004). Along the Big Thompson river in Moraine Park, populations steadily declined from an estimate of over 300 beavers in 1940 (Packard 1947) to 12 beavers in 1994 to 1998 (Zeigenfuss et al. 2002). Beaver surveys and aerial photographs taken in 1999 revealed only one beaver colony in Moraine Park. Declines or a complete absence of beaver has been observed throughout drainages on the east side of the park. Most of the current beaver populations occur in areas with low willow use by elk. Beaver are largely absent from areas with heavy use by elk, with the east slope population in the park currently estimated to be ~40 beaver (Baker et al. 2004). Similarly, in the Kawuneeche Valley, beaver numbers were estimated to be about 600 in 1949, and only 30 in 1999 (C. Westbrook unpublished data).

As mentioned previously, the >90% decline in beaver numbers in Moraine Park is correlated with a large reduction (69%) in surface water and willow declines over the last 50 to 60 years (Peinetti et al. 2002, Zeigenfuss et al. 2002). An estimated 90% decline in the beaver population in Horseshoe Park has contributed to a large decrease (47%) in surface water that has also likely contributed to the decline in willow (Zeigenfuss et al. 2002). The lack of beaver is accelerating willow declines by inhibiting the development of appropriate sites for willow seedling establishment and limiting recharge of the shallow aquifers in Moraine Park and Horseshoe Park.

Baker et al. (2004) found that beaver in the park prefer relatively tall, unbrowsed willow and select against short, hedged willow. The interaction of beaver cutting and elk browsing strongly suppresses compensatory growth in willow, altering the structure and function of the willow community and facilitating a conversion from tall willow to a short willow community. As the few beaver that still exist in Moraine Park and Horseshoe Park cut down the tall willow for food, elk browse the plants and keep them short (less than 30 cm; Baker et al. 2005).

The decline in beaver has also reduced the reproductive capabilities of willow. Beaver are important to willow reproduction in several ways. Two of the three landforms that provide suitable sites for willow establishment by seed in the park — abandoned beaver ponds and abandoned channels or ox-bows — are associated with the presence of beaver in an area (Cooper et al. 2003). In addition, because there are currently very few beaver on the elk winter range, little to no reproduction occurs via willow stem fragments. Beaver cuttings allow willows to colonize areas that are suitable for willow growth but not seedling establishment, and may be important on a landscape scale (Cottrell 1995). Existing willow roots or stems cut by beaver can give rise to new shoots to become a new plant, which allows willow to be maintained for long periods of time (100 years or longer).

Summary

There is little evidence that elk herbivory is negatively affecting white-tailed ptarmigan in alpine tundra areas. However, several factors strongly suggest more research is needed before the effects of elk can be discounted, including the negative correlation between elk and ptarmigan numbers, high percentage of alpine willow in elk summer diets, and possible declines in alpine willow coverages. Increased winter temperatures have also been suggested to negatively affect ptarmigan population growth rates and must also be considered in any future research.

Bird and butterfly diversity has been found to be highest in aspen, willow riparian, and ponderosa pine habitat on the primary winter range. No studies have shown a direct effect of elk on these animal species in RMNP, but the large impact of elk on these habitats on the core winter range, which contains the majority of these habitats, should be a management concern. In particular, these habitats support a distinct biological community at these elevations that are not found elsewhere in the park because almost all other aspen and willow stands are located at higher elevations in either coniferous forests or alpine tundra. The mechanism by which elk are suggested to decrease bird abundance is via decreased live trees in aspen stands and decreased structural complexity of shrubs in aspen and willow habitat. If current trends continue it could lead to the local exclusion of several cavity nesting and willow-dependent bird species.

Beaver declines in RMNP are believed to initially be due to trapping that began in the 1940s. Beaver declines over the last 50-60 years contributed to a loss of water and willow on the core winter range. Subsequent declines or lack-of-recovery of beaver is attributed to competition with elk for food, in particular tall willow, which has a limited distribution on the core winter range. As remaining tall willow on the core winter range are cut for food by beaver, their subsequent regrowth is suppressed by elk. This is leading to further declines in the beaver and willow populations.

NUTRIENT CYCLES

Nutrient cycles are considered important to ecosystem sustainability because depletion of nitrogen (N) and soil organic matter reduces long-term plant productivity (Vitousek 1982) and could result in a change in plant species composition (Schoenecker et al. 2004).

Schoenecker et al. (2002) used empirical measures and an ecosystem model (CENTURY) to examine elk and their effect on the overall sustainability of winter range areas in RMNP. They found N mineralization rates were lower in grazed versus ungrazed short willow sites ($P < 0.07$), as were nitrate pools ($P < 0.10$). Schoenecker et al. (2004) also suggested elk are transferring N away from aspen and at least some willow communities (e.g., short willow communities). Nitrogen cycles in tall willow areas were not affected, and elk grazing did not affect N cycles or pools in the upland shrub areas (Schoenecker et al. 2004).

CENTURY modeling indicated soil N concentrations on the winter range under the current level of herbivory is expected to decline by less than 2% over the next 50 years (Schoenecker et al. 2002). If the elk herd continues to increase, this could increase the loss of N to 2% or slightly more, but it is unknown what effects such losses would have (if any) on the biodiversity, abundance, or status of plant species on the core winter range.

Other empirical results have found little evidence of major effects of elk on nutrient cycles or soil N concentrations on the core winter range in RMNP. Elk have been found to slightly increase or decrease some soil and plant C and N concentrations, but no large, consistent effects have been observed (Binkley et al. 2003, Menezes et al. 2001, Zeigenfuss et al. 2002).

The SAVANNA model predicted soil carbon (C) and N pools to remain relatively stable under a wide range of grazing intensities and elk abundance. Coughenour (2002) states: "There was a very slow, nearly imperceptible decline over long time periods (50-100 years)...the sizes of the soil organic matter C and N pools were also very large relative to annual plant uptake and grazing offtake, which provided a large buffering capacity". However, Coughenour (2002) acknowledges that a more detailed analysis of grazing impacts on soil fertility under different grazing conditions or spatial locations would be useful.

CONCLUSIONS

The limited historical, archeological, and ethnographic information that is available suggests that ungulates, including elk, were historically abundant in the RMNP area, but actual population sizes or densities from this time cannot be determined. Current research indicates that elk population size and density in the Estes Valley has exceeded the natural range of variability that would be expected in the presence of an intact predator community. There is no single number of elk that should be considered natural. Instead, a range of elk numbers that correspond to differences in annual food availability and predation by wolves is the appropriate reference for natural conditions. This is referred to as a tri-trophic equilibrium among elk, food, and wolves. Currently, a bi-trophic equilibrium exists among elk and their food base. This has led to a winter subpopulation in RMNP that is at ecological carrying capacity and a subpopulation in the town area that is at or approaching carrying capacity. Under this bi-trophic system, elk population estimates for the Estes Valley over the last 10 years have ranged from 1,700-3,500 elk. Current research indicates 1,200-2,100 elk would exist in the area under a tri-trophic system. This is supported by the literature and should be considered a conservative estimate of the natural range of variability; i.e., the greatest number of elk that would have been present. This is due to the fact that the above population estimates for elk in a tri-trophic system did not consider the effects of two additional and potentially major predators that were present in the RMNP ecosystem prior to European settlement, humans and grizzly bears.

No estimates are available for what range of densities should be considered natural, but there are several strong indications that densities in the current bi-trophic system are greater than they would be under a tri-trophic system. To our knowledge, elk densities on the core winter range are the highest concentrations ever documented for a free-ranging population that is not

artificially fed. There is also a growing body of evidence that indicates wolves can have a large impact on elk distribution and group size. The lack of wolves in the RMNP area has likely allowed for prolonged use and aggregation on the core winter range grasslands where predation pressure by wolves and behavioral adaptations by elk would be greatest.

The population size and density of elk in the Estes Valley is directly affecting several vegetation communities, most notably aspen and willow on the core winter range areas. It cannot be established with certainty that aspen are a long-term natural feature of the core winter range, but we consider it prudent to conserve the remaining stands until this can be determined through future research and monitoring efforts. Research indicates the best way to ensure aspen persistence on the core winter range is to either drastically reduce elk densities or exclude herbivory until new trees that are large enough to withstand herbivory have been recruited into the population. Fencing is the most practical option because it may be difficult to achieve the necessary elk densities (≤ 4 to 15 elk/km²) while still allowing elk populations to fluctuate within the natural range of variability.

The question of whether or not aspen are a natural feature of the winter range can be answered in one of two ways. First, the effects of elk can be monitored after aspen stands on the core winter range are restored via fencing and elk population dynamics are within their natural range of variability. Second, genetic or carbon dating techniques may be able to be used to determine how long the current aspen clones have been present on the winter range. Aspen should be considered a long-term, natural feature of the core winter range if rejuvenated aspen stands are able to persist in the presence of elk populations within their natural range of variability or if it can be determined that aspen clones were present prior to European settlement of the area.

Historical accounts and current research indicates willow distribution on the core winter range is noticeably reduced from pre-European settlement times. Elk browsing is currently the proximate factor preventing willow growth and reproduction. Recent declines and management efforts needed to restore willow are confounded by a beaver population that has declined significantly over the last 50 years and cannot become re-established due to the effects of elk. The lack of beaver in the area has led to a loss of surface water and suitable sites for willow reproduction in the area. Further, as beaver cut down tall willow for food, elk suppress its regrowth and lead to additional declines in beaver and willow.

Full restoration of willow on the winter range will require not only a reduction in elk browsing pressure but increased beaver activity as well. However, willow must be restored before beaver can return to the area. Research indicates that the maximum density of elk on core winter range areas will need to be reduced to ≤ 15 - 30 elk/km² to allow willow to grow beyond the height of herbivory, produce seeds, and provide suitable food for beaver. Restoration efforts will need to be closely monitored for several reasons. First, maximum elk densities may need to be lower than 15 - 30 elk/km², as evidence for this range resulting in increased willow growth is limited. Second, fencing to prevent elk browsing or possibly food supplementation to reduce the effects of beaver foraging may be necessary to allow the willow-elk-beaver interaction to achieve a sustainable level.

The above reasons for the restoration of elk dynamics and vegetation conditions are based on the conclusion that elk and their associated effects are outside the natural range of variation. The other major reason for restoration is to sustain or increase a variety of plant and animal species that are directly or indirectly affected by elk. The current effects of elk herbivory and subsequent reductions in the structural complexity of aspen and willow habitat is likely to have adverse effects on mule deer, willow-dependent and cavity-nesting birds, butterflies, and individual plant species such as sulphur buckwheat that inhabit the core winter range. It also remains to be determined if elk are negatively affecting alpine willow and ptarmigan populations, as no experimental research or studies on ptarmigan ecology or mortality have been conducted.

We recognize that questions remain about the role of elk in the RMNP ecosystem. Research and modeling efforts used to predict the specific effects of wolves on the population dynamics and distribution of elk is relatively recent, with the majority of work being conducted during the last 10-15 years. However, the best available research to date indicates that elk population size and densities in the Estes Valley are outside their natural range of variability and this is resulting in a number of adverse effects in the RMNP ecosystem. From an ecological perspective, there are clear tradeoffs between continuing the current management course of nonintervention and initiating active management to control elk dynamics and restore vegetation communities of concern. Research indicates that nonintervention will continue to result in an elk population at or near ecological carrying capacity, declines of aspen and willow and few or no beaver on the core winter range, and a localized decline or loss of plant and animal species. The effects of management intervention depend on the population and density goals for the elk population and specific management actions undertaken. Research indicates that if elk densities were low enough to allow willow to escape the effects of elk browsing, it would result in a trophic cascade that would increase willow distribution, beaver populations, and habitat for a variety of songbirds. Underscoring the importance of timely management intervention is the fact that as conditions in aspen and willow habitat on the winter range continue to deteriorate, restoration efforts will become increasingly difficult or impossible to achieve.

ACKNOWLEDGEMENTS

We thank J. Peek and D. Houston for their very helpful and insightful reviews.

LITERATURE CITED

- Anderson, E. M. 2002. Influences of elk on upland aspen, riparian willow, and associated landbirds in and near Jackson Hole, Wyoming. M.S. Thesis. University of Wyoming, Laramie.
- Alstad, K. P., J. M. Welker, S. A. Williams, and M. J. Trlica. 1999. Carbon and water relations of *Salix monticola* in response to winter browsing and changes in surface water hydrology: an isotopic study using $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$. *Oecologia* 120:375-385.
- Altmann, M. 1956. Patterns of herd behavior in free-ranging elk of Wyoming, *Cervus canadensis nelsoni*. *Zoologica* 41:116-143.
- Arno, S. F. 1980. Forest fire history of the northern Rockies. *Journal of Forestry* 78:460-465.
- Augustine, D. J., and S. J. McNaughton. 1998. Ungulate effects on the functional species composition of plant communities: herbivore selectivity and plant tolerance. *Journal of Wildlife Management* 62:1165-1183.
- Bailey, J. A. 1986. The increase and dieoff of Waterton Canyon bighorn sheep: biology, management and dismanagement. Biennial Symposium of the Northern Wild Sheep and Goat Council 5:325-340.
- Bailey, J. A. 1990. Management of Rocky Mountain bighorn sheep herds in Colorado. Colorado Division of Wildlife Special Report No. 66. Colorado Division of Wildlife. Fort Collins, Colorado.
- Baker, B. W., and B. S. Cade. 1995. Predicting biomass of beaver food from willow stem diameters. *Journal of Range Management* 48:322-326.
- Baker, B. W., and E. P. Hill. 2003. Beaver (*Castor canadensis*). Pages 288-310 in G. A. Feldhamer, B. C. Thompson, and J. A. Chapman, eds. *Wild mammals of North America: Biology, Management, and Conservation*. Johns Hopkins University Press, Baltimore, MD.
- Baker, B. W., H. C. Ducharme, D. C. S. Mitchell, T. R. Stanley, and H. R. Peinetti. 2004. Why aren't there more beaver in Rocky Mountain National Park? *Proceedings of the Annual Conference of the Colorado Riparian Association* 17:85-90.
- Baker, B. W., H. C. Ducharme, D. C. S. Mitchell, T. R. Stanley, and H. R. Peinetti. 2005. Interaction of beaver and elk herbivory reduces standing crop of willow. *Ecological Applications* 15:110-118.
- Baker, W. L., J. A. Munroe, and A. E. Hessel. 1997. The effects of elk on aspen in the winter range of Rocky Mountain National Park. *Ecography* 20:155-165.

- Ballard, W. B., J. S. Whitman, and C. L. Gardner. 1987. Ecology of an exploited wolf population in south-central Alaska. Wildlife Monographs No. 98.
- Bear, G. D. 1989. Seasonal distribution and population characteristics of elk in Estes Valley, Colorado. Colorado Division of Wildlife Special Report No. 65.
- Benedict, J. B. 1992. Footprints in the snow: high altitude cultural ecology of the Colorado Front Range, U.S.A. Arctic and Alpine Research 24:1-16.
- Benedict, J. B. 1996. The game drives of Rocky Mountain National Park. Center for Mountain Archeology Research Report No. 7.
- Bergerud, A. T., and J. P. Elliot. 1986. Dynamics of caribou and wolves in northern British Columbia. Canadian Journal of Zoology 64:1515-1529.
- Bergerud, A. T., W. Wyett, and J. B. Snider. 1983. The role of wolf predation in limiting a moose population. Journal of Wildlife Management 47:977-988.
- Bilbrough, C. J., and J. H. Richards. 1993. Growth of sagebrush and bitterbrush following simulated winter browsing: mechanisms of tolerance. Ecology 74:481-492.
- Binkley, D., F. Singer, M. Kaye, and R. Rochelle. 2003. Influence of elk grazing on soil properties in Rocky Mountain National Park. Forest Ecology and Management 185:239-247.
- Blaisdell, J. P., and W. F. Mueggler. 1956. Sprouting of bitterbrush (*Purshia tridentata*) following burning or top removal. Ecology 37:365-377.
- Boutin, S. 1992. Predation and moose population dynamics: a critique. Journal of Wildlife Management 56:116-127.
- Boyce, M. S. 1993. Predicting the consequences of wolf recovery to ungulates in Yellowstone National Park. Pages 234-269 in R. S. Cook, ed. Ecological issues on reintroducing wolves into Yellowstone National Park. National Park Service Scientific Monographs 92/22.
- Braun, C. E., D. R. Stevens, K. M. Giesen, and C. P. Melcher. 1991. Elk, white-tailed ptarmigan and willow relationships: a management dilemma in Rocky Mountain National Park. the North American Wildlife Conference Transactions:56:74-85.
- Brunswig, R. H., Jr. 2001. Lawn Lake (5LR318): results of an archaeological mitigation research project at a high altitude prehistoric site in Rocky Mountain National Park. Final report to the National Park Service.
- Brunswig, R. H., Jr., and J. Doerner. 2001. The Lawn Lake site (5LR318): new evidence for high altitude late Paleoindian adaptations and Paleolandscapes of Colorado's southern

Rockies in the early Holocene. Paper presented at the 5th Biennial Rocky Mountain Anthropological Conference, Waterton Lakes National Park, Alberta, Canada, September 20-23, 2001.

Bryant, J. P., F. S. Chapin, III, and D. R. Klein. 1983. Carbon/nutrient balance of boreal plants in relation to vertebrate herbivory. *Oikos* 40:357-368.

Buchholtz, C. W. 1983. Rocky Mountain National Park: a history. Colorado Associated University Press, Boulder.

Buckland, S., D. Anderson, K. Burnham, and J. Laake. 1993. Distance Sampling: Estimating Abundance of Biological Populations. Chapman and Hall, New York, NY.

Bunting, S. C., L. F. Neuenschwander, and G. E. Gruell. 1985. Fire ecology of antelope bitterbrush in the northern Rocky Mountains. Pages 48-57 in J. E. Lotan and J. K. Brown, eds. Fire's effects on wildlife habitat: symposium proceedings. USDA Forest Service General Technical Report INT-186. Intermountain Research Station. Ogden, Utah.

Butler, W. B. 1997. Rocky Mountain National Park research design for archeology. Final report to National Park Service and Colorado State Historic Preservation Office.

Buttery, R. 1955. Range conditions and trends resulting from winter concentrations of elk in Rocky Mountain National Park, Colorado. M.S. Thesis. Colorado State University, Fort Collins.

Capp, J. C. 1967. Competition among bighorn sheep, elk and deer in Rocky Mountain National Park, Colorado. M.S. Thesis. Colorado State University, Fort Collins.

Capp, J. C. 1968. Bighorn sheep, elk, and mule deer range relationships. A review of the literature. Final report to the Rocky Mountain Nature Association.

Cassells, E. S. 1995. Hunting the open high country: prehistoric game driving in the Colorado alpine tundra. Ph.D. Dissertation, Department of Anthropology, University of Wisconsin, Madison.

Cassirer, E. F., L. E. Oldenburg, V. L. Coggins, P. Fowler, K. Rudolph, D. L. Hunter, and W. J. Foreyt. 1996. Overview and preliminary analysis of a bighorn sheep dieoff, Hells Canyon 1995-1996. Biennial Symposium of the Northern Wild Sheep and Goat Council 10:78-86.

Caughley, G. 1976. Wildlife management and the dynamics of ungulate populations. Pages 183-246 in T. H. Coaker, ed. Applied Biology. Academic Press, London, UK.

Caughley, G. 1981. Overpopulation. p. 7-19. In: P. A. Jewell, and S. Holt. eds. Problems in management of locally abundant wild mammals. Academic Press, NY.

Choate, T. S. 1963. Habitat and population dynamics of white-tailed ptarmigan in Montana. *Journal of Wildlife Management* 27:684-699.

Clarke, J. A., C. E. Olmsted, and K. Larkins. 1994. Patterns of habitat use, movement, and distribution of elk adjacent to the eastern boundary of Rocky Mountain National Park. Final report to the National Park Service.

Clements, C. D., and J. A. Young. 1996. Influence of rodent predation on antelope bitterbrush seedlings. *Journal of Range Management* 49:31-34.

Clements, F. E. 1910. The life history of lodgepole burn forests. USDA Forest Service Bulletin 79. Government Printing Office, Washington D. C.

Coggins, V. L. 1988. The Lostine Rocky Mountain bighorn sheep die-off and domestic sheep. *Biennial Symposium of the Northern Wild Sheep and Goat Council* 6:57-64.

Cole, G. F. 1971. An ecological rationale for the natural or artificial regulation of native ungulates in parks. *North American Wildlife Conference Transactions*: 36:417-425.

Colorado Division of Wildlife. 1999. Declining mule deer populations in Colorado: reasons and responses. A report to the Colorado Legislature.

Colorado Division of Wildlife. 2003. Prevalence and distribution of chronic wasting disease in Colorado, 2002-2003. Available on the internet at <http://wildlife.state.co.us/CWD/pdf/PrevalenceSummary8-031.pdf>.

Colorado Division of Wildlife. 2004. Prevalence and distribution of chronic wasting disease in Colorado, 2003-2004. Available on the internet at http://wildlife.state.co.us/CWD/pdf/PrevalenceSummary_9-14-04.pdf.

Condon, D. D. 1943. Recommendations and an outline for the management of the elk and deer population in Rocky Mountain National Park. National Park Service Report.

Connor, J. 1993. Neotropical migrant bird survey for Rocky Mountain National Park. Rocky Mountain National Park Resource Management Series #17.

Conner, M. M. 2004. Estimation of mule deer population size, survival, and a description of movements in Rocky Mountain National Park. Final report to the National Park Service.

Cooper, D. J., J. Dickens, and E. Gage. 2003. Constraints on, and opportunities for, riparian willow establishment, Rocky Mountain National Park, Colorado. Final report to the National Park Service.

- Cooper, D. J., S. W. Woods, R. A. Chimner, and L. H. MacDonald. 2000. Effects of the Grand Ditch on wetlands of the Kawuneeche Valley, Rocky Mountain National Park, Colorado. Final report to the National Park Service.
- Cottrell, T. R. 1993. The comparative ecology of *Salix planifolia* and *Salix monticola* in Rocky Mountain National Park. Ph.D. Dissertation, Colorado State University, Fort Collins.
- Cottrell, T. R. 1995. Willow colonization of Rocky Mountain mires. *Canadian Journal of Forest Research* 25:215-222.
- Coughenour, M. B. 2002. Elk in the Rocky Mountain National Park ecosystem B a model based assessment. Final report to the National Park Service.
- Creel, S., J. Winnie, Jr., B. Maxwell, K. Hamlin, and M. Creel. 2005. Elk alter habitat selection as an antipredator response to wolves. *Ecology* 86:3387-3397.
- Cunningham, R. L. 1988. North American migratory birds and the National Park System. Some interpretive thoughts. Final report to the National Park Service.
- Dekker, D. G., W. Bradford, and J. R. Gunson. 1996. Elk and wolves in Jasper National Park, Alberta from historical times to 1992. Pages 85-94 in L. N. Carbyn, S. H. Fritts, and D. R. Scrip, eds. *Ecology and conservation of wolves in a changing world*. Canadian Circumpolar Institute, University of Alberta, Edmonton.
- Denney, R. N., R. J. Boyd, E. Bucknall, W. Schuett, P. F. Gilbert, J. Cooney, L. D. Hibbs, G. D. Bear, and R. B. Gill. 1967. Research project segment: elk seasonal movements, Rocky Mountain National Park. Job Completion Report (State of Colorado).
- Dobkin, D. S., F. J. Singer, and W. S. Platts. 2002. Ecological condition and avian response in willow, aspen, and cottonwood communities of the National Elk Refuge, Jackson, Wyoming. Final report by High Desert Ecological Research Institute.
- Dixon, J. S. 1939. Special report on the elk problem at Rocky Mountain National Park. Final report to the National Park Service.
- Duberstein, J. N. 2001. The effects of elk on nest site selection in cavity-nesting birds in Rocky Mountain National Park. Final report to the National Park Service.
- Edwards, J. 1983. Diet shifts in moose due to predator avoidance. *Oecologia* 60:185-189.
- Ehle, D. S. 2001. Spatial and temporal patterns of disturbance and ponderosa pine forest structure in Rocky Mountain National Park. M.S. Thesis. University of Wyoming, Laramie.
- Estes, M. 1939. The memoirs of Estes Park. Reprint of 1939 Colorado Magazine article.

Farmer, R. E., Jr. 1962. Aspen root sucker formation and apical dominance. *Forest Science* 8:403-410.

Ferguson, J. W., A. T. Bergerud, and R. Ferguson. 1988. Predation risk and habitat selection in the persistence of a remnant caribou population. *Oecologia* 76:236-245.

Forsyth, D. M., and G. J. Hickling. 1998. Increasing Himalayan tahr and decreasing chamois densities in the eastern Southern Alps, New Zealand: evidence for interspecific competition. *Oecologia* 113:377-382.

Frost, H., G. Storm, M. Batcheller, and M. Lovallo. 1997. White-tailed deer management at Gettysburg National Military Park and Eisenhower National Historic Site. *Wildlife Society Bulletin* 25:462-469.

Fryxell, F. M. 1928. The former range of the bison in the Rocky Mountains. *Journal of Mammalogy* 9:129-139.

Fryxell, J. M. 1987. Food limitation and demography of a migratory antelope, the white-eared kob. *Oecologia* 72:83-91.

Gage, E. A., and D. J. Cooper 2004a. Constraints on willow seedling survival in a Rocky Mountain floodplain. *Wetlands* 24:908-911.

Gage, E. A., and D. J. Cooper 2004b. Controls on willow cutting survival in a montane riparian area. *Journal of Range Management* 57:597-600.

Gage, E. A., and D. J. Cooper, 2005. Patterns of willow seed dispersal, seed entrapment, and seedling establishment in a heavily browsed montane riparian ecosystem. *Canadian Journal of Botany* 83:678-687.

Garton, E. O., R. L. Crabtree, B. B. Ackerman, and G. Wright. 1990. The potential impact of a reintroduced wolf population on the northern Yellowstone elk population. Pages 3-59 to 3-91 in J. D. Varley and W. G. Brewster, eds. *Wolves for Yellowstone? A report to the United States Congress, vol. 2, research and analysis*. National Park Service, Yellowstone National Park.

Gasaway, W. C., K. T. Gasaway, and H. H. Berry. 1983. Persistent low densities of plains ungulates in Etosha National Park, Namibia: testing the food-regulating hypothesis. *Canadian Journal of Zoology* 74: 1556-1572.

Gasaway, W. C., R. D. Boertje, D. V. Grangaard, D. G. Kelleyhouse, R. O. Stephenson, and D. G. Larsen. 1992. The role of predation in limiting moose at low densities in Alaska and Yukon and implications for conservation. *Wildlife Monographs* No. 120.

Goodson, N. J. 1978. Status of bighorn sheep in Rocky Mountain National Park. M.S. Thesis. Colorado State University, Fort Collins.

Goodson, N. J., and D. R. Stevens. 1991. Population dynamics, home ranges, and habitat use of bighorn sheep on the Continental Divide and in the Never Summer Mountains. Final report to the National Park Service.

Grater, R. K. 1945. Report on the elk-deer reduction program of 1944-1945 at Rocky Mountain National Park. National Park Service Report.

Griggs, R. F. 1956. Competition and succession on a Rocky Mountain fellfield. *Ecology* 37:8-20.

Gross, J. E., and M. W. Miller. 2001. Chronic wasting disease in mule deer: a model of disease dynamics, control options, and population consequences. *Journal of Wildlife Management* 65:205-215.

Guse, N. G., Jr. 1966. An administrative history of and elk herd. M.S. Thesis. Colorado State University, Ft. Collins.

Gysel, L. W. 1960. An ecological study of the winter range of elk and mule deer in the Rocky Mountain National Park. *Journal of Forestry* 58:696-703.

Harrington, F. A. 1978. Ecological segregation of ungulates in alpine and subalpine communities. Ph.D. Dissertation, Colorado State University, Fort Collins.

Hebblewhite, M., C. A. White, C. G. Nietvelt, J. A. McKenzie, T. E. Hurd, J. M. Fryxell, S. E. Bayley, and P. C. Paquet. 2005. Human activity mediates a trophic cascade caused by wolves. *Ecology* 86:2135-2144.

Hebblewhite, M., D. H. Pletscher, and P. C. Paquet. 2002. Elk population dynamics in areas with and without predation by recolonizing wolves in Banff National Park, Alberta. *Canadian Journal of Zoology* 80:789-799.

Hinds, T. E. 1985. Diseases. Pages 87-106 in N.V. DeByle and R. P. Winokur, eds. *Aspen: ecology and management in the western United States*. USDA Forest Service General Technical Report RM-119. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado.

Hobbs, N. T., D. L. Baker, J. E. Ellis, and D. M. Swift. 1981. Composition and quality of elk winter diets in Colorado. *Journal of Wildlife Management* 45:156-171.

Hobbs, N. T., D. L. Baker, J. E. Ellis, D. M. Swift, and R. A. Green. 1982. Energy- and nitrogen-based estimates of elk winter-range carrying capacity. *Journal of Wildlife Management* 46:12-21.

- Hobbs, N. T., D. L. Baker, G. D. Bear, and D. C. Bowden. 1996a. Ungulate grazing in sagebrush grassland: mechanisms of resource competition. *Ecological Applications* 6:200-217.
- Hobbs, N. T., D. L. Baker, G. D. Bear, and D. C. Bowden. 1996b. Ungulate grazing in sagebrush grassland: effects of resource competition on secondary production. *Ecological Applications* 6:218-227.
- Hobbs, N. T., D. Ojima, J. Loomis, S. Weiler, M. McDuff, D. M. Theobald, A. P. Covich, J. S. Baron, M. B. Coughenour, D. J. Cooper, L. Landrum, J. Dickens. 2003. An integrated assessment of the effects of climate change on Rocky Mountain National Park and its gateway community. Unpublished report to the Environmental Protection Agency.
- Houston, D. B. 1982. The northern Yellowstone elk. Macmillan Publishing Co., New York.
- Huff, D. and J. Varley. 1999. Natural regulation in Yellowstone National Park's northern range. *Ecological Applications* 9:17-29.
- Jarrett, R. D., and J. E. Costa. 1993. Hydrology and geomorphology of the 1982 Lawn Lake dam failure, Colorado. Pages 1-17 in H. E. McCutchen, R. Herrmann, and D. R. Stevens, eds. *Ecological effects of the Lawn Lake Flood of 1982, Rocky Mountain National Park*. Scientific Monograph NPS/NRROMO/NRSM-93/21.
- Jehle, G. 2004. Avian response to prescribed fire in the upland shrub/ponderosa pine system of Rocky Mountain National Park, Colorado. M.S. Thesis, Colorado State University, Ft. Collins.
- Jenkins, K. J. and R. G. Wright. 1987. Dietary niche relationships among cervids relative to winter snowpack in northwestern Montana. *Canadian Journal Zoology* 65:1397-1401.
- Johnsgard, P. A. 1986. *Birds of the Rocky Mountains*. Colorado Associated University Press, Boulder.
- Jones, J. R., and G. A. Schier. 1985. Growth. Pages 19-24 in N.V. DeByle and R. P. Winokur, eds. *Aspen: ecology and management in the western United States*. USDA Forest Service General Technical Report RM-119. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado.
- Jorgenson, J.T., M. Festa-Bianchet, J-M. Gaillard, and W. D. Wishart, W.D. 1997. Effects of age, sex, disease, and density on survival of bighorn sheep. *Ecology* 78:1019-1032.
- Kay, C. E. 1997a. The condition and trend of aspen, *Populus tremuloides*, in Kootenay and Yoho National Parks: Implications for ecological integrity. *The Canadian Field-Naturalist* 111:607-606.

Kay, C. E. 1997*b*. Is aspen doomed? *Journal of Forestry* 95:4-11.

Kaye, M. W., D. Binkley, and T. J. Stohlgren. 2005. Effects of conifers and elk browsing on quaking aspen forests in the central Rocky Mountains, USA. *Ecological Applications* 15:1284-1295.

Kershaw, L., A. MacKinnon, and J. Pojar. 1998. *Plants of the Rocky Mountains*. Lone Pine Publishing, Edmonton.

Kiener, W. 1939. Sociological studies of the alpine vegetation on Longs Peak. Ph.D. Dissertation, University of Nebraska, Lincoln.

Kufeld, R. C., O. C. Wallmo, and C. Feddema. 1973. Foods of the Rocky Mountain mule deer. USDA Forest Service Research Paper RM-111. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado.

Kunkel, K. E., and D. H. Pletscher. 1999. Species-specific population dynamics of cervids in a multipredator ecosystem. *Journal of Wildlife Management* 63:1082-1093.

Larkins, K. F. 1997. Patterns of elk movement and distribution in and adjacent to the eastern boundary of Rocky Mountain National Park. M.S. Thesis. University of Northern Colorado, Greeley.

Laundré, J.W., L. Hernández, and K. B. Altendorf. 2001. Wolves, elk, and bison: reestablishing the "landscape of fear" in Yellowstone National Park, U.S.A. *Canadian Journal of Zoology* 79:1401-1409.

Leukering, T., and M. Carter. 1999. Abundance and nesting success of common breeding birds in willow riparian areas in Rocky Mountain National Park. Final report to the National Park Service.

Lloyd, J. 1997. The effect of elk herbivory on vegetation and songbirds of aspen communities in Rocky Mountain National Park. Final report to the National Park Service.

Lubow, B. C., F. J. Singer, T. L. Johnson, and D. C. Bowden. 2002. Dynamics of interacting elk populations within and adjacent to Rocky Mountain National Park. *Journal of Wildlife Management* 66:757-775.

Mack, J. A., and F. J. Singer 1993. Using pop-II models to predict effects of wolf predation and hunter harvests on elk, mule deer, and moose on the northern range. Pages 49-74 *in* R. S. Cook, ed. *Ecological issues on reintroducing wolves into Yellowstone National Park*. National Park Service Scientific Monographs 92/22.

- Martin, A. C., H. S. Zim, and A. L. Nelson. 1951. A guide to wildlife food habits. Dover Publications, Inc., New York.
- Mast, J. N., T. T. Veblen, and Y. B. Linhart. 1998. Disturbance and climatic influences on age structure of ponderosa pine at the pine/grassland ecotone, Colorado Front Range. *Journal of Biogeography* 25:743-755.
- May, T. A., and C. E. Braun. 1972. Seasonal foods of adult white-tailed ptarmigan in Colorado. *Journal of Wildlife Management* 36:1180-1186.
- McClintock, B. T. 2004. The mark-resight method: an application to bighorn sheep in Colorado and a simulation study evaluating estimators. M.S. Thesis. Colorado State University, Fort Collins.
- McDonough, W. T. 1985. Sexual reproduction, seeds, and seedlings. Pages 25-28 *in* N.V. DeByle and R. P. Winokur, eds. *Aspen: ecology and management in the western United States*. USDA Forest Service General Technical Report RM-119. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado.
- McLaren, B. E., and R. O. Peterson. 1994. Wolves, moose, and tree rings on Isle Royale. *Science* 266: 1555-1558.
- McLaughlin, J. S. 1931. First annual spring wildlife report, RMNP. National Park Service Report.
- McShea, W. H. B. Underwood, and J. Rappole. 1997. Deer management and the concept of overabundance. Pages 1-7 *in* W. McShea, H. B. Underwood, and J. Rappole, eds. *The Science of Overabundance: Deer Ecology and Population Management*. Smithsonian Institution Press, Washington, D.C.
- Mduma, S. A. R., A. R. E. Sinclair, and R. Hilborn. 1999. Food regulates the Serengeti wildebeest. *Journal of Animal Ecology* 68:1101-1122.
- Menezes, R. S. C., E. T. Elliot, and J. M. Welker. 2002. Isotopic evidence of the effects of herbivory and landscape position on plant nitrogen sources in a riparian ecosystem. *In* F. J. Singer, ed. *Ecological evaluation of the abundance and effects of elk in Rocky Mountain National Park, Colorado, 1994-1999*. Final report to the National Park Service.
- Menezes, R. S. C., E. T. Elliot, D. W. Valentine, and S. A. Williams. 2001. Carbon and nitrogen dynamics in elk winter ranges. *Journal of Range Management* 54:400-408.
- Messier, F., and M. Crete. 1985. Moose-wolf dynamics and the natural regulation of moose populations. *Oecologia* 65:503-512.

Miller, M. W., E. S. Williams, C. W. McCarty, T. R. Spraker, T. J. Kreeger, and E. T. Thorne. 2000. Epidemiology of chronic wasting disease in free-ranging cervids. *Journal of Wildlife Diseases* 36:676-690.

Monello, R. J., D. L. Murray, and F. Cassirer. 2001. Ecological correlates of pneumonia epizootics in bighorn sheep. *Canadian Journal of Zoology* 79:1423-1432.

Mueggler, W. F. 1985. Vegetative associations. Pages 45-55 in N.V. DeByle and R. P. Winokur, eds. *Aspen: ecology and management in the western United States*. USDA Forest Service General Technical Report RM-119. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado.

Naiman, R. J., C. A. Johnston, and J. C. Kelley. 1988. Alteration of North American streams by beaver. *Bioscience*. 387:753-762.

National Park Service. 2001. Management Policies. United States Department of Interior, Washington D.C. Available on the internet at <http://www.nps.gov/policy/mp/policies.html>.

Nesvacil, K., and C. E. Olmsted. 2003. The effects of prescribed fire and ungulate herbivory 6 and 7 years post-burn in the upland bitterbrush (*Purshia tridentata*) communities of Rocky Mountain National Park, Colorado. Proceedings - 2002 Fire Conference: Managing Fire and Fuels in the Remaining Wildlands and Open Spaces of the Southwest, December 2-5, 2002, San Diego, California. USDA Forest Service General Technical Report.

Nietvelt, C. G. 2001. Herbivory interactions between beaver (*Castor canadensis*) and elk (*Cervus elaphus*) on willow (*Salix* spp.) in Banff National Park, Alberta. M.S. Thesis. University of Alberta, Edmonton.

Olmsted, C.E. 1979. The ecology of aspen with reference to utilization by large herbivores in Rocky Mountain National Park. Pages 89-97 in M. S. Boyce and L. D. Hayden-Wing, eds. *North American elk: ecology, behavior and management*. University of Wyoming Press, Laramie.

Olmsted, C.E. 1997. Twenty years of change in Rocky Mountain National Park elk winter range aspen. Unpublished Report to the National Park Service.

Orians, G., D. A. Cochran, J. W. Duffield, T. K. Fulla, R. J. Gutierrez, W. M. Hanemann, F. C. James, P. Karieva, S. R. Kellert, D. Klein, B. N. McLellan, P. D. Olson, and G. Yaska. 1997. Wolves, bears and their prey in Alaska. National Academy of Press, Washington, D.C.

Packard, F. M. 1939. Notes on a conversation of Fred M. Packard with Mr. Abner Sprague at Estes Park Colorado on August 7, 1939, regarding the early history of the bighorn and other game animals in Rocky Mountain National Park. Unpublished Report to the National Park Service.

Packard, F. M. 1946. An ecological study of the bighorn sheep in Rocky Mountain National Park. *Journal of Mammalogy* 27:3-28.

Packard, F. M. 1947. A survey of the beaver population of Rocky Mountain National Park, Colorado. *Journal of Mammalogy* 28:219-227.

Peek, J. M. and A. Lovaas. 1968. Differential distribution of elk by sex and age on the Gallatin winter range. *Journal Wildlife Management* 32:553-557.

Peinetti, H. R. 2000. Riparian willow dynamics and their interaction with environmental and biological factors in the elk winter range of Rocky Mountain National Park (Colorado) B a multi-scale analysis. Ph.D. Dissertation, Colorado State University, Fort Collins.

Peinetti, H. R., M. A. Kalkhan, and M. B. Coughenour. 2002. Long-term changes in willow spatial distribution on the elk winter range of Rocky Mountain National Park (USA). *Landscape Ecology*: 17:341-354.

Peinetti, H. R., R. S. C. Menezes, and M. B. Coughenour. 2001. Changes induced by elk browsing in the aboveground biomass production and distribution of willow (*Salix monticola* Bebb): their relationships with plant water, carbon, and nitrogen dynamics. *Oecologia* 127:334-342.

Peterson, R. O. 1999. Wolf-moose interaction on Isle Royale: the end of natural regulation. *Ecological Applications* 9:10-16.

Ratcliff, H. M. 1941*a*. Winter range conditions in Rocky Mountain National Park. *North American Wildlife Conference Transactions*: 6:132-139.

Ratcliff, H. M. 1941*b*. Progress report on the bighorn sheep study, Rocky Mountain National Park. Final report to the National Park Service.

Ripple, W. J., and R. L. Beschta. 2004. Wolves, elk, willows, and trophic cascades in the upper Gallatin Range of Southwestern Montana, USA. *Forest Ecology and Management* 200:161-181.

Ripple, W. J., E. J. Larsen, R. A. Renkin, and D. W. Smith. 2001. Trophic cascades among wolves, elk and aspen on Yellowstone National Park's northern range. *Biological Conservation* 102:227-234.

Robisch, E. and R. G. Wright. 1995. A survey of ungulate management in selected U. S. national parks. *Natural Areas Journal* 15:117-123.

Romme, W. H., L. Floyd-Hanna, D. D. Hanna, and E. Bartlett. 2001. Aspen's ecological role in the west. Pages 243-259 in W. D. Shepperd, D. Binkley, D. L. Bartos, T. J. Stohlgren, and L. G. Eskew, compilers. Sustaining aspen in western landscapes: symposium proceedings. Proceedings RMRS-P-18. Rocky Mountain Research Station. Ft. Collins, Colorado.

Romme, W. H., M. G. Turner, L. L. Wallace, and J. S. Walker. 1995. Aspen, elk, and fire in northern Yellowstone National Park. *Ecology* 76: 2097-2106.

Rowdabaugh, K. M. 1978. The role of fire in the ponderosa pine-mixed conifer ecosystems. M.S. Thesis. Colorado State University, Fort Collins.

Ryder, T. J., E. S. Williams, K. W. Mills, K. H. Bowles, and E. T. Thorne. 1992. Effect of pneumonia on population size and lamb recruitment in Whiskey Mountain bighorn sheep. Biennial Symposium of the Northern Wild Sheep and Goat Council 8:136-146.

Schell, L. D., and T. J. Stohlgren. 1997. Effect of elk grazing on native and non-native plant species diversity in Beaver Meadows. Final report to the Rocky Mountain Elk Foundation.

Schier, G. A. 1975a. Promotion of sucker development on *Populus tremuloides* root cuttings by and antiauxin. *Canadian Journal of Forest Research* 5:338-340.

Schier, G. A. 1975b. Deterioration of aspen clones in the middle Rocky Mountains. USDA Forest Service Research Paper INT-170. Intermountain Forest and Range Experiment Station, Ogden, Utah.

Schier, G. A. 1985. Vegetative regeneration. Pages 29-33 in N.V. DeByle and R. P. Winokur, eds. Aspen: ecology and management in the western United States. USDA Forest Service General Technical Report RM-119. Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado.

Schmitz, O. J., A. P. Beckerman, and K. M. O'Brien. 1997. Behaviorally mediated trophic cascades: effects of predation risk on food web interactions. *Ecology* 78:1388-1399.

Schoenecker, K. A., F. J. Singer, R. S. C. Menezes, L. C. Zeigenfuss, and D. Binkley. 2002. Sustainability of vegetation communities grazed by elk in Rocky Mountain National Park. In F. J. Singer, ed. Ecological evaluation of the abundance and effects of elk in Rocky Mountain National Park, Colorado, 1994-1999. Final report to the National Park Service.

Schoenecker, K. A., F. J. Singer, L. C. Zeigenfuss, D. Binkley, and R. S. C. Menezes. 2004. Effects of elk herbivory on vegetation and nitrogen processes. *Journal of Wildlife Management* 68:837-849.

Shepperd, W. D., and O. Engelby. 1983. Rocky Mountain aspen. U.S. Forest Service Agricultural Handbook 445:77-79.

- Sherriff, R. L., T. T. Veblen, and J.S. Sibold. 2001. Fire history in high elevation subalpine forests in the Colorado Front Range. *Ecoscience* 8:369-380.
- Simonson, S. E., P. A. Opler, T. J. Stohlgren, and G. W. Chong. 2001. Rapid assessment of butterfly diversity in a montane landscape. *Biodiversity and Conservation* 10:1369-1386.
- Sinclair, A. R. E. 1985. Does interspecific competition or predation shape the African ungulate community? *Journal of Animal Ecology* 54:899-918.
- Sinclair, A. R. E., H. Dublin, and M. Borner. 1985. Population regulation of the Serengeti wildebeest: a test of the food hypothesis. *Oecologia* 65:266-268.
- Singer, F. J., and J. E. Norland. 1994. Niche relationships within a guild of ungulate species in Yellowstone National Park, Wyoming, following release from artificial controls. *Canadian Journal of Zoology* 72:1383-1394.
- Singer, F. J., D. M. Swift, M. B. Coughenour, and J. D. Varley. 1998*b*. Thunder on the Yellowstone revisited: an assessment of management of native ungulates by natural regulation, 1968-1993. *Wildlife Society Bulletin* 26:375-390.
- Singer, F. J., G. Wang, and N. T. Hobbs. 2003. The role of ungulates and large predators on plant communities and ecosystem processes in national parks. Pages 444-486 *in* R. G. Anthony and C. Zabel, eds. *Large mammal communities in U.S. western coniferous forests*. Cambridge University Press, Cambridge.
- Singer, F. J., L. C. Mack, and R. C. Cates. 1994. Ungulate herbivory of willows on Yellowstone's northern winter range. *Journal of Range Management* 47: 435-443.
- Singer, F. J., L. C. Zeigenfuss, B. Lubow, and M. J. Rock. 2002. Ecological evaluation of the appropriate number of ungulates in U.S. National Parks: a case study. *In* F. J. Singer, ed. *Ecological evaluation of the abundance and effects of elk in Rocky Mountain National Park, Colorado, 1994-1999*. Final report to the National Park Service.
- Singer, F. J., L. C. Zeigenfuss, R. G. Cates, and D. T. Barnett. 1998*a*. Elk, multiple factors, and persistence of willows in national parks. *Wildlife Society Bulletin* 26:419-428.
- Skogland, T. 1985. The effects of density-dependent resource limitation on the demography of wild reindeer. *Journal of Animal Ecology* 54:359-374.
- Smith, B. L., and S. H. Anderson. 1996. Patterns of neonatal mortality of elk in northwest Wyoming. *Canadian Journal of Zoology* 74:1229-1237.
- Sprague, A. E. 1925. Game more plentiful now than 50 years ago in Estes Park region. *Estes Park Trail*, June 5, 1925.

Stevens, D.R. 1980*a*. The deer and elk of Rocky Mountain National Park; A 10-year study. National Park Service Report ROMO-N-13.

Stevens, D. R. 1980*b*. Impact of elk on alpine tundra in Rocky Mountain National Park. *Proceedings of High Altitude Vegetation Workshop* 4:228-244.

Stohlgren, T. J., L. D. Schell, and B. V. Heuvel. 1999. How grazing and soil quality affect native and exotic plant diversity in Rocky Mountain grasslands. *Ecological Applications* 9:45-64.

Suzuki, K., H. Suzuki, D. Binkley, and T. J. Stohlgren. 1999. Aspen regeneration in the Colorado Front Range: differences at local and landscape scales. *Landscape Ecology* 14:231-237.

Toll, O. W. 1962. Arapaho names and trails: a report of a 1914 pack trip. Privately published.

Turchi, G. M., P. L. Kennedy, D. Urban, and D. Hein. 1994. Bird species richness in relation to isolation of aspen habitats. *Wilson Bulletin* 107:463-474.

United States Senate. 1967. Hearings before a Subcommittee on Appropriations, U. S. Senate: First Session on Control of Elk Populations in Yellowstone National Park. Washington, D. C. GPO.

Veblen, T. T., T. Kitzberger, and J. Donnegan. 2000. Climatic and human influences on fire regimes in ponderosa pine forests in the Colorado Front Range. *Ecological Applications* 10:1178-1195.

Veblen, T. T., and D. C. Lorenz. 1991. The Colorado Front Range: a century of ecological change. University of Utah Press, Salt Lake City.

Vitousek, P. 1982. Nutrient cycling and nutrient use efficiency. *American Naturalist* 119:553-572.

Wambolt, C. L. 1998. Sagebrush and ungulate relationships on Yellowstone's northern range. *Wildlife Society Bulletin* 26:429-437.

Wambolt, C. L., K. S. Walhof, and M. R. Frisina. 2001. Recovery of big sagebrush communities after burning in south-western Montana. *Journal of Environmental Management* 61:243-252.

Wang, G., N. T. Hobbs, K. M. Giesen, H. Galbraith, D. S. Ojima, and C. E. Braun. 2002. Relationships between climate and population dynamics of white-tailed ptarmigan in Rocky Mountain National Park, Colorado, USA. Final report to National Park Service.

Weisberg, P.J., and M. B. Coughenour. 2003. Model-based assessment of aspen responses to elk herbivory in Rocky Mountain National Park, U.S.A. *Journal of Environmental Management* 32:152-169.

White, C. A., and M. C. Feller. 2001. Predation risk and elk-aspen foraging patterns. Pages 61-80 *in* W. D. Shepperd, D. Binkley, D. L. Bartos, T. J. Stohlgren, and L. G. Eskew, eds. *Sustaining aspen in western landscapes: symposium proceedings, 13-15 June, Grand Junction, CO. USDA Forest Service Proceedings RMRS-P-18. Rocky Mountain Research Station. Fort Collins, Colorado.*

White, C. A., C. E. Olmsted, and C. E. Kay. 1998. Aspen, elk, and fire in the Rocky Mountain national parks of North America. *Wildlife Society Bulletin* 26:449-462.

Willard, B. E. 1963. Phytosociology of the alpint tundra of Trail Ridge, Rocky Mountain National Park, Colorado. Ph.D. Dissertation, University of Colorado, Boulder.

Williams, G. R. 1954. Population fluctuations in some northern hemisphere game birds (Tetraonidae). *Journal of Animal Ecology* 23:1-34.

Wolff, J. O., and T. Van Horn. 2003. Vigilance and foraging patterns of American elk during the rut in habitats with and without predators. *Canadian Journal of Zoology* 81:266-271.

Wright, G. M., J. S. Dixon, and B. H. Thompson. 1933. *Fauna of the National Parks of the United States: a preliminary survey of faunal relations in National Parks.* U.S. Government Printing Office, Washington D.C.

Wright, R. G. 1992. *Wildlife research and management in the National Parks.* University of Illinois Press, Chicago.

Zaninelli, J., and T. Leukering. 1998. Aspen: a critical component of cavity-nesting bird habitat in Rocky Mountain National Park. Final report to the National Park Service.

Zeigenfuss, L. C., F. J. Singer, and D. Bowden. 1999. Vegetation responses to natural regulation of elk in Rocky Mountain National Park. Biological Science Report USGS/BRD/BSRC1999-0003. U.S. Government Printing Office, Denver, Colorado.

Zeigenfuss, L. C., F. J. Singer, S. Williams, and T. Johnson. 2002. Influences of herbivory and water on willow in elk winter range. *Journal of Wildlife Management* 66:788-795.

Appendix 1. Common plant species found in Rocky Mountain National Park covertypes. Common names are given when available.

Cover Type/Distribution	Common Plant Species Found
Spruce/Fir	Overstory Species:
The spruce/fir cover type generally occurs between 9,000 feet and 11,600 feet. Lower elevations consist of a dense spruce/fir overstory, while higher elevations (> 10,000 feet) are usually made up of spruce/fir islands with grassy openings.	Subalpine Fir (<i>Abies lasiocarpa</i>)
	Engelmann Spruce (<i>Picea engelmannii</i>)
	Lodgepole Pine (<i>Pinus contorta</i>)
	Understory Species:
	Heartleaf Arnica (<i>Arnica cordifolia</i>)
	Alpine Mountain-Sage (<i>Artemisia scopulorum</i>)
	American Bistort (<i>Bistorta bistortoides</i>)
	Marsh Reedgrass (<i>Calamagrostis canadensis</i>)
	(<i>Carex arapahoensis</i>)
	Pyrenean Sedge (<i>Carex pyrenaica</i>)
	Ross' Sedge (<i>Carex rossii</i>)
	One-Flowered Erigeron (<i>Erigeron simplex</i>)
	Short-Leaved Fescue (<i>Festuca ovina brach.</i>)
	Richardson Geranium (<i>Geranium richardsonii</i>)
	Drummond's Rush (<i>Juncus drummondii</i>)
	Small Flowered Woodrush (<i>Luzula parviflora</i>)
	Brunt-Fruited Sweet-Cicely (<i>Osmorhiza depauperata</i>)
	Mountain Timothy (<i>Phleum alpinum</i>)
	Sticky Jacob's-Ladder (<i>Polemonium delicatum</i>)
	Wintergreens (<i>Pyrola</i> spp.)
	(<i>Senecio crassulus</i>)
	Arrow-Leaved Groundsel (<i>Senecio triangularis</i>)
	Sibbaldia (<i>Sibbaldia procumbens</i>)
	Star-Flowered False Solomon's-Seal (<i>Smilicina stellata</i>)
	Clovers (<i>Trifolium</i> spp.)
	Whortleberry, Blueberries, Huckleberries (<i>Vaccinium</i> spp.)
	Alpine Speedwell (<i>Veronica wormskjoldii</i>)
Lodgepole Pine	Overstory Species:
The lodgepole pine cover type generally occurs between 8,000 and 10,000 feet. Lodgepole is typically found on most any slope or aspect above 8,500 feet, but is successional to spruce/fir forests higher elevations.	Lodgepole Pine (<i>Pinus contorta</i>)
	Douglas Fir (<i>Pseudotsuga menziesii</i>)
	Subalpine Fir (<i>Abies lasiocarpa</i>)
	Engelmann Spruce (<i>Picea engelmannii</i>)
	Understory Species:
	Common Bearberry (<i>Arctostaphylos uva-ursi</i>)
	Heartleaf Arnica (<i>Arnica cordifolia</i>)
	(<i>Berberis repens</i>)
	Elk sedge (<i>Carex geyeri</i>)
	Ross' Sedge (<i>Carex rossii</i>)
	Fireweed (<i>Epilobium angustifolium</i>)
	(<i>Jamesia americana</i>)
	Common Juniper (<i>Juniperus communis</i>)
	(<i>Oreochrysum parryi</i>)

Appendix 1 (continued). Common plant species found in Rocky Mountain National Park covertypes. Common names are given when available.

Cover Type/Distribution	Common Plant Species Found
Lodgepole Pine (continued)	Understory Species (continued): Blue-Mist Penstemon (<i>Penstemon virens</i>) Leafy cinquefoil (<i>Potentilla fissa</i>) Canada buffaloberry (<i>Shepherdia canadensis</i>) Spike Trisetum (<i>Trisetum spicatum</i>) Low bilberry (<i>Vaccinium myrtilus</i>) Whortleberry (<i>Vaccinium scoparium</i>)
Ponderosa Pine The ponderosa pine cover type is only found on the eastern side of the park and occupies a wide variety of habitats below 9,000 feet.	Overstory Species: Ponderosa Pine (<i>Pinus ponderosa</i>) Lodgepole Pine (<i>Pinus contorta</i>) Limber Pine (<i>Pinus flexilis</i>) Trembling Aspen (<i>Populus tremuloides</i>) Douglas Fir (<i>Pseudotsuga menziesii</i>) Understory Species: (<i>Agropyron albicans</i>) Sun Sedge (<i>Carex heliophila</i>) Ross' Sedge (<i>Carex rossii</i>) Sulphur Buckwheat (<i>Eriogonum umbellatum</i>) Western wallflower (<i>Erysimum asperum</i>) (<i>Geranium fremontii</i>) Common Juniper (<i>Juniperus communis</i>) (<i>Muhlenbergia filiculmis</i>) Blue mist penstemon (<i>Penstemon virens</i>) (<i>Purshia tridentata</i>)
Douglas Fir The douglas fir cover type is found mostly on the east side of the park from the lowest elevations to just over 9,000 feet. It is most prevalent on steep, north facing slopes.	Overstory Species: Douglas Fir (<i>Pseudotsuga menziesii</i>) Ponderosa Pine (<i>Pinus ponderosa</i>) Lodgepole Pine (<i>Pinus contorta</i>) Understory Species: Ross' Sedge (<i>Carex rossii</i>) (<i>Fragaria americana</i>) (<i>Jamesia americana</i>) Common Juniper (<i>Juniperus communis</i>) Rocky Mountain ninebark (<i>Physocarpus monogynus</i>) Leafy cinquefoil (<i>Potentilla fissa</i>)
Limber Pine The limber pine cover type is prominent on the eastern side of the park, and ranges from 9,000 to 11,000 feet. This is a minor cover type found on steep, windwept areas that have a southern or western exposure.	Overstory Species: Limber Pine (<i>Pinus flexilis</i>) Subalpine Fir (<i>Abies lasiocarpa</i>) Engelmann Spruce (<i>Picea engelmannii</i>) Lodgepole Pine (<i>Pinus contorta</i>) Understory Species: Common Bearberry (<i>Arctostaphylos uva-ursi</i>) Fendler's Sandwort (<i>Arenaria fendleri</i>)

Appendix 1 (continued). Common plant species found in Rocky Mountain National Park covertypes. Common names are given when available.

Cover Type/Distribution	Common Plant Species Found
Limber Pine (continued)	Understory Species (continued): Marsh Reedgrass (<i>Calamagrostis canadensis</i>) Ross' Sedge (<i>Carex rossii</i>) (<i>Festuca ovina</i>) Northern Gentian (<i>Gentianella amarella</i>) Common Juniper (<i>Juniperus communis</i>) Alpine Sandwort (<i>Minuartia obtusiloba</i>) Sticky Jacob's-Ladder (<i>Polemonium viscosum</i>) Spotted Saxifrage (<i>Saxifraga bronchialis</i>) Lance-Leaved Stonecrop (<i>Sedum lanceolatum</i>) Spike-Like Goldenrod (<i>Solidago spathulata</i>) Whiproot Clover (<i>Trifolium dasyphyllum</i>)
Aspen The aspen cover type occurs from 8,000 feet to treeline (about 11,600 feet), but is most common at intermediate elevations on a variety of aspects.	Overstory Species Trembling Aspen (<i>Populus tremuloides</i>) Colorado Blue Spruce (<i>Picea pungens</i>) Lodgepole Pine (<i>Pinus contorta</i>) Understory Species: Yarrow (<i>Achillea lanulosa</i>) Creeping Oregon-Grape (<i>Berberis repens</i>) Elk Sedge (<i>Carex geyeri</i>) Smooth Wildrye (<i>Elymus glaucus</i>) White Geranium (<i>Geranium richardsonii</i>) Prairie Rose (<i>Rosa woodsii</i>) Fendler's Meadowrue (<i>Thalictrum fendleri</i>)
Riparian The riparian cover type occupies a variety of areas from the lowest elevations in the park to treeline (about 11,600 feet). Most riparian cover occurs in valley-bottoms, low-lying forest openings, and on high elevation snow areas.	Overstory Species Trembling Aspen (<i>Populus tremuloides</i>) Mountain Willow (<i>Salix monticola</i>) Flat-Leaved Willow (<i>Salix planifolia</i>) (<i>Salix wolfii</i>) Understory Species: Marsh Reedgrass (<i>Calamagrostis canadensis</i>) Mountain Marsh-Marigold (<i>Caltha leptosepala</i>) Water Sedge (<i>Carex aquatilis</i>) Small-Winged Sedge (<i>Carex microptera</i>) Beaked Sedge (<i>Carex ultriculata</i>) Tufted Hairgrass (<i>Deschampsia cespitosa</i>) Field Timothy (<i>Phleum pratense</i>)
Meadows The meadow cover type is characterized by dry grasslands and shrublands at low elevations on both sides of the park.	Various grasses (<i>Agropyron</i> spp.) Big Sagebrush (<i>Artemisa tridentata</i>) Timber Oatgrass (<i>Danthonia intermedia</i>) Parry's Oatgrass (<i>Danthonia parryi</i>) (<i>Festuca thurberi</i>) Long-Haired Needlegrass (<i>Stipa comata</i>)