


A Biodiversity Conservation Assessment for Lake Superior



Volume 2: Regional Unit Summaries

Prepared by the Superior Work Group (SWG) of the Lake Superior Lakewide Action and Management Plan (LAMP)

Final Draft: June 2013

Recommended Citation

Lake Superior Lakewide Action and Management Plan (LAMP) - Superior Work Group. 2013. Lake Superior Biodiversity Conservation Assessment: Regional Unit Summaries. 222pp.

Disclaimer

This report reflects the best efforts of the preparers (Dan Kraus and Megan Ihrig) to accurately represent and interpret the available expertise and information on Lake Superior and the views and opinions of project participants. Every effort to ensure the accuracy of the information contained in this study has been taken. We welcome suggestions for improvements.

Data Sources

For a full list of the Data Sources and Methods used to create the Tables and Figures throughout this Regional Unit Summaries report please see Appendix A: Spatial Data Catalogue and Methods.

Volume 1: Lakewide Assessment & Volume 2: Regional Summaries

Please note that this report includes two volumes. Volume 1 includes an assessment of lakewide biodiversity target health and threats. Volume 2 contains regional summaries and maps that are presented in this document. It is recognized that many regions contain additional information and mapping on biodiversity and threats that could not be fully reflected in this report. Wherever possible, regional and local data and spatial information on biodiversity targets and threats has been noted in the text.

Acknowledgements

This report has been prepared by a project Steering Committee from the Superior Work Group (SWG) of the Lake Superior Lakewide Action and Management Plan (LAMP) with coordination and support from the Nature Conservancy of Canada. The Steering Committee members included:

Environment Canada: Rob Hyde

Environmental Protection Agency: Elizabeth LaPlante

Great Lakes Indian Fish and Wildlife Commission: Ann McCammon Soltis, Jennifer Vanator

Michigan Department of Environmental Quality: Matt Preisser

Minnesota Department of Natural Resources: John Jereczek

National Park Service: Joan Elias (retired)

Nature Conservancy of Canada: Megan Ihrig, Dan Kraus,

Ontario Ministry of Natural Resources: Marilee Chase, Neil Dawson, Sue Greenwood

Parks Canada: Ray Boudreau

U.S. Fish and Wildlife Service: Henry Quinlan

USDA Forest Service: Mark Fedora

USDA Natural Resources Conservation Service: Dana Raines

Wisconsin Department of Natural Resources: Cherie Hagen

The Steering Committee acknowledges the valuable input provided by the following reviewers:

Tyler Kaspar	1854 Treaty Authority
Andrew Ecclestone	Anishinabek/Ontario Fisheries Resource Centre
Julie Van Stappen	Apostle Islands National Lakeshore
Cyrus Hester	Bad River Band of the Lake Superior Tribe of Chippewa Indians
Naomi Tillison	Bad River Band of the Lake Superior Tribe of Chippewa Indians
Tom Gorenflo	Chippewa Ottawa Resource Authority; St. Marys River
Mike Ripley	Chippewa Ottawa Resource Authority; St. Marys River
Laurie Wood	Environment Canada
Katheryne O'Connor	Environment Canada/ Canadian Wildlife Service
Daryl Seip	Environment Canada/ Canadian Wildlife Service
Scott Millard	Emeritus, Department of Fisheries and Oceans
Tom Pratt	Fisheries and Oceans Canada
Nancy Schuldt	Fond du Lac Band of Lake Superior Chippewa, Environmental Program
Seth Moore	Grand Portage Band of Chippewa
Lucinda Johnson	Great Lake Environmental Indicators Project/ University of Minnesota
Sigrid Smith	Great Lakes Environmental Assessment and Mapping Project
Erin Johnston	Keweenaw Bay Indian Community
George Beck	Lac Vieux Desert Band of Lake Superior Chippewa Indians
Anne Hokanson	Michigan Department of Environmental Quality
Stephanie Swart	Michigan Department of Environmental Quality
William Taft	Michigan Department of Environmental Quality
David Caroffino	Michigan Department of Natural Resources
Amy Clark Eagle	Michigan Department of Natural Resources
Christopher Hoving	Michigan Department of Natural Resources
Glenn Palmgren	Michigan Department of Natural Resources
Phil Schneeberger	Michigan Department of Natural Resources
Shawn Sitar	Michigan Department of Natural Resources
Sue Tangora	Michigan Department of Natural Resources
Bruce Carlson	Minnesota Department of Natural Resources
Maya Hamady	Minnesota Department of Natural Resources
Ethan Perry	Minnesota Department of Natural Resources
Annie Bracey	Natural Resources Research Institute, University of Minnesota Duluth
Gerald Niemi	Natural Resources Research Institute, University of Minnesota Duluth

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

Tom Beechey	Nature Conservancy of Canada
James Duncan	Nature Conservancy of Canada
Nick Lapointe	Nature Conservancy of Canada
Chris Maher	Nature Conservancy of Canada
Brenda Koenig	Ontario Ministry of Natural Resources
Christine Drake	Parks Canada
Cavan Harpur	Parks Canada
Chantal Vis	Parks Canada
Matthew Hudson	Sigurd Olson Environmental Institute, Northland College
Mary Khoury	The Nature Conservancy
Doug Pearsall	The Nature Conservancy
Linda Wire	University of Minnesota
Sue Eggert	U.S. Forest Service
Dale Higgins	U.S. Forest Service Chequamegon-Nicolet National Forest
Gary Cypinski	U.S. Dept. of Interior, Fish & Wildlife Service
Brian Huberty	U.S. Dept. of Interior, Fish & Wildlife Service
Janet Keough	U.S. Environmental Protection Agency
Daniel Yule	U.S. Geological Survey
Gary Caspar	University of Wisconsin Milwaukee Field Station
Kate Barrett	Wisconsin Department of Natural Resources
William Blust	Wisconsin Department of Natural Resources
Diane Daulton	Wisconsin Department of Natural Resources
Andrew Fayram	Wisconsin Department of Natural Resources
Lynelle Hanson	Wisconsin Department of Natural Resources
Martin Jennings	Wisconsin Department of Natural Resources
Frank Koshere	Wisconsin Department of Natural Resources
Steve LaValley	Wisconsin Department of Natural Resources
Ryan Magana	Wisconsin Department of Natural Resources
Ryan O'Connor	Wisconsin Department of Natural Resources
Paul Piszczek	Wisconsin Department of Natural Resources
Peter Stevens	Wisconsin Department of Natural Resources
Fred Strand	Wisconsin Department of Natural Resources
Scott Toshner	Wisconsin Department of Natural Resources

Mapping and spatial analysis was completed by Andrea Hebb and Gary White of the Nature Conservancy of Canada (NCC). Sarah Perkins from Battelle kindly supported collecting expert information in the webinars.


Special thanks to the amazing NCC team that supported the project website and communications with our Lake Superior experts and project administration: Christine Beevis-Trickett, Quincin Chan, James Duncan, Craig Easton, Shane Eisenberg, Teva Harrison, Dana Kleniewski, Lynda Kosalle, Chris Maher, Laura Mousseau, Luc Thomas & Chuck Van Kempen.

Funding for this project is from Environment Canada and the Nature Conservancy of Canada.

Guide to the Lake Superior Regional Unit Summaries

Volume Two of the Lake Superior Biodiversity Conservation Assessment provides more detailed information and mapping on 20 regional units (Figure 1). These units are based on quaternary watershed boundaries that were grouped based on coastal environments identified from Lake Superior. In addition to the watershed and coast, each regional unit includes associated inshore and nearshore waters. In some regions, offshore waters were also incorporated to include islands. One unit that encompasses all of the Lake Superior offshore waters was also included. Maintaining the open waters as a single unit was recommended by the Aquatic Community Committee/Lake Superior Technical Committee.

Figure 1.0: Lake Superior Regional Units



This regional information is intended to compliment the lakewide assessment provided in Volume One, and to provide additional detail to support the development of place-based conservation actions. Information in the regional summaries was based on a review of the literature and expert input. Spatial information was calculated for this project using the data sources outlined in the data catalogue (see Appendix A in this volume). It is recognized that many regions contain additional information and mapping on biodiversity and threats that could not be fully reflected in this report. Wherever possible, regional and local data and spatial information on biodiversity targets and threats has been included in the text.

This introduction provides some detail on the different sections of the regional unit summaries.

Healthy Waters Report Card

The grades provided in the report card, and conditions and trend table are intended to denote relative condition/health and stresses for each biodiversity target in the region based on available condition and stress indices (see below). **These grades are intended to help highlight where the biodiversity targets are likely in better or worse health than the lakewide average, and to inform discussion about priority areas for conservation and restoration.** This automated assignment on the relative regional health of biodiversity targets was subject to expert review, and in cases where the experts felt the grade did not reflect actual conditions, the results were overridden with this expert input. Expert opinions were also reflected in the trends section of the conditions and trend table.


For each regional unit, a regional average of all stress/condition indices was calculated based on the individual scores of each sub-unit within the region (see Volume One) For example in Regional Unit 1 (Goulais), each of the 92 quaternary watersheds has a watershed stress index value (in this case ranging from 0 to 0.754, max=1). The regional score is based on an average of these sub-units. For the watershed stress index and Great Lakes Cumulative Stress, the average was subtracted from 1. For the Coastal Condition Index, the average was subtracted from the maximum possible score. These regional average values were then applied to the biodiversity targets. For some targets only one average index was used (e.g. the average value of the watershed stress index was used for the tributary and watershed target). For other targets, the condition/stress is likely reflected by a combination of the indices (e.g. embayment health), and the average of multiple indices was applied. The final score/grade is an average of the score for all biodiversity targets.

By the Numbers Table

This table provides a summary of information on land/water cover, coastal features, condition and land ownership and protected areas for each region. Where applicable, this information is put into context from a regional and Lake Superior-wide perspective. For example, the Goulais regional unit has 92.2 km of sand beaches. This amount of sand beaches make up almost 30% of the total coastline for the Goulais region, and represents 14.7% of all sand beaches on Lake Superior. The notes column provides some clarification on the different calculations for some attributes.

Coastal and Watershed Features Map Series


This map series depicts land cover, sub-watershed boundaries, coastal habitats (e.g. cobble beach, coastal wetlands) and fish spawning areas. The shaded areas denote current and historic spawning areas for lake trout and lake whitefish. The point data generally reflect more accurate locations of current spawning areas. This map is intended to provide an overview of the biodiversity targets in the region.



Condition Map Series


This map series details the stress/condition indices: Watershed Stress Index, Great Lakes Cumulative Stress and the Coastal Condition Index. This map (in combination with the report card) is intended to provide information on the health of the biodiversity targets and highlight potential issue areas.

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries



Important Habitat Sites and Areas Map Series

This map series shows the Important Habitat Sites and Important Habitat Areas mapped by the Lake Superior LAMP (Lake Superior Binational Program [LSBP] Habitat Committee 2006), in the context of the protected areas and land ownership and management (e.g. public lands, national parks) in each regional unit. Additional important habitat areas may occur in some regions, and not all categories of land protection and ownership are reflected. Box 1 provides the criteria that were used to select these sites. Conserving or restoring these system components are of highest priority to maintaining Lake Superior biodiversity, recognizing that other important habitat areas inevitably exist, most notably in remote expanses of Lake Superior's east and north shores where habitats are largely undisturbed.



Box 1: Criteria for the Identification of Biodiversity Features in the Lake Superior Watershed

Ecosystems

1. Large, relatively unfragmented areas most representative of the Lake Superior basin ecosystem that support natural community assemblages where ecosystem dynamics are intact or can be restored.
2. Nationally significant ecosystems. Areas that have wildlife and plant habitat values that go beyond local values in that they provide substantial benefits that extend beyond the basin.
3. Old Growth Forest. Tracts of varying size supporting native old growth forest. Tracts that with restoration and proper management could support high quality, native old growth forest.
4. Coastal shore or coastal wetland ecosystems. Sites that have, or with restoration could develop, high quality, diverse ecosystems that are representative of the interacting communities unique to the Lake Superior shoreline.
5. Areas that support high biological and ecological diversity. Sites that support, or with restoration could support the compositional, functional, and structural elements associated with diverse ecosystems.
6. Habitats that contribute to, or with restoration could contribute to maintaining ecosystem integrity on a landscape scale. These areas could include buffering communities around currently protected ecosystems, core areas within a managed area, or may be connecting corridors between important habitat sites.

Communities

1. Rare communities. Communities that are of high quality, or have high restoration potential, or are critically endangered. Examples include: calcareous fens, beach dunes, interdunal wetlands, red clay wetland complexes, bedrock beaches and cliffs.
2. Plant and wildlife habitats that are rare in the Lake Superior basin, or are rare globally.
3. Plant and wildlife habitats that occur only in the Great Lakes basin.
4. Communities that are, or that with restoration could be, outstanding representatives of the natural (i.e. pre-settlement) ecosystem.

Species

1. Sites (large or small) that serve as habitat for vulnerable, endangered, threatened or special concern species (or candidate species) during any stage of their life cycle. Currently occupied habitats and sites with potential for future colonization or reintroduction are included. Prioritization of potential sites depends on status of the species (i.e. rarity at global, sub-national, and basin scales), likelihood of occupation and the quality (or restoration potential) of the site.
2. Sites that serve, or with restoration may serve, vital functions in the life cycle of species named in appropriate planning documents (e.g. Lake Superior Ecosystem Objectives, Fish-Community Objectives for Lake Superior, Tribal resource plans, etc.)
3. Habitats required for the conservation of migratory wildlife (e.g. neotropical migrant birds, migratory fish, etc.), including staging areas, migration corridors and routes.
4. Spawning and nursery grounds for reptiles, amphibians, fish, or aquatic invertebrates. Colonial water bird nesting sites.
5. Habitats that can contribute to the conservation of species most likely to be at risk from human activity.
6. Habitats that support species that provides important ecological functions (e.g. nutrient cycling or chemical detoxification.)

Table of Contents

Guide to the Lake Superior Regional Unit Summaries	3
1. Goulais	9
2. Agawa/ Michipicoten – Magpie.....	20
3. White and Pic	30
4. Michipicoten Island.....	39
5. Little Pic.....	47
6. Jack Pine/ Nipigon.....	57
7. Black Sturgeon.....	67
8. Dog/Arrow.....	77
9. Baptism-Brule.....	87
10. Isle Royale.....	99
11. Beaver-Lester.....	108
12. St. Louis/ Cloquet.....	118
13. Beartrap-Nemadji	132
14. Bad-Montreal	144
15. Black-Presque Isle /Ontonagon	157
16. Keweenaw Peninsula/ Sturgeon.....	166
17. Dead-Kelsey	177
18. Betsy-Chocolay	187
19. Waiska/ Tahquamenon	197
20. Lake Superior Open Waters.....	206
References	211
Appendix A.....	220

1. Goulais

HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	B	COASTAL WETLANDS	B
EMBAYMENTS & INSHORE	B	COASTAL TERRESTRIAL	A
TRIBUTARIES & WATERSHEDS	B	OVERALL B+	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	Ecologically desirable status; requires little intervention for maintenance
B Good	Within acceptable range of variation; may require some intervention for maintenance.
C Fair	Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.
D Poor	Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.
Unknown	Insufficient information.



View of Lake Superior shoreline from Batchawana Bay. Photo credit: Sue Greenwood/ Ontario Ministry of Natural Resources

Summary/ Description

The Goulais regional unit is located in Ontario on the eastern shore of Lake Superior, and extends from the international boundary at the St. Marys River in the south to near the Montreal River Harbour in the north. This regional unit is 5,929.95 km² in size, including the associated nearshore waters. A number of provincial parks, conservation reserves and enhanced management areas are located in this regional unit. The largest community in the area, Sault Ste. Marie, Ontario is located near the southern end of this regional unit. Other communities in this regional unit include Goulais River, Havilland, Harmony Bay, Batchawana Bay, Obadjiwan (Batchewana¹) First Nation and Searchmont. The jurisdictional area of the Sault Ste. Marie Region Conservation Authority overlap with a small portion of this regional unit. The Goulais regional unit contains one tertiary watershed (Goulais) and 10 quaternary watersheds. The watersheds are dominated by forests, with only small areas of developed and agricultural lands. The coast is characterized by sand and cobble beaches, and includes some of the largest sand beaches on the Ontario side of Lake Superior. This region has more private lands than most other Ontario regions, and only 10% of the coast is in protected areas.

¹ The Batchawana spelling is used when referring to geographical places, while the Batchewana spelling is used when referring to the Obadjiwan First Nation.

TABLE 1.1: Goulais BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
<i>Agriculture</i>	35.46	0.60	1,287.56	
<i>Developed</i>	9.72	0.16	351.99	
<i>Forest</i>	4,428.42	74.68	96,789.30	
<i>Associated Nearshore Waters</i>	1,343.51	22.6	15,951.22	
<i>Other</i>	69.69	1.18	7,452.78	
<i>Water (inland)</i>	43.15	0.73	8,283.49	
Total Area	5,929.95	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
<i>Coastline (km)</i>	308.1	NA	5.42	Based on SOLEC shoreline
<i>Sand Beaches (km)</i>	92.30	29.96	14.66*	*% of Lake Superior Total Sand Beaches
<i>Coastal Wetlands (km²)</i>	30.10	7.53*	3.06**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
<i>Natural Cover in Coastal Zone (km²)</i>	375.21	93.91*	6.79**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
<i>Number of Islands</i>	145	NA	5.5	
Condition	Region	Region %	% of Lake Superior Total	
<i>Population Density (persons/km²)</i>	0.44	NA		
<i>Road Density (km/km²)</i>	0.68	NA		
<i>Number of Dams and Barriers</i>	1,627	NA	6.9	
<i>Artificial Shoreline (km)</i>	15.23	4.06	6.86	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
<i>Private</i>	1,470.78	32.21	4,566.52	Regional area based on landmass
<i>Public/Crown</i>	2,696.54	59.05	4,566.52	
<i>Tribes/ First Nations</i>	6.61	0.14	4,566.52	
<i>Parks & Protected Areas (total)</i>	396.18	8.68	4,566.52	
<i>Parks & Protected Areas (coast)</i>	39.12	9.79*	399.55**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- The Goulais regional unit contains sites of Important Habitat for both lake trout and lake whitefish. Important Habitat sites for lake trout are found off the coast of the Goulais region, in the nearshore zone (Lake Superior Binational Program Habitat Committee 2006) (Figure 1.1).
- Goulais Bay and Batchawana Bay are noted as Lake Superior embayments which are important for lake sturgeon (Auer 2003). In the Goulais regional unit these embayments and the nearshore zone, which provides corridors for movement, are identified as critical management areas for lake sturgeon in the Lake Superior basin (Auer 2003).

Coastal Zone and Islands

- Ile Parisienne is noted as an Important Habitat site for lake whitefish (Lake Superior Binational Program Habitat Committee 2006). Important spawning areas for lake trout and lake whitefish are noted in the shoals and waters around the island (Figure 1.1). Most of the island (991 hectares) and a marine zone which extends 1.6 km from the shore of Ile Parisienne into Whitefish Bay are designated as a provincial conservation reserve, which is 4,669 hectares in total size (OMNR 2005a).
- The waters near the islands of Sandy Islands Provincial Park are Important Habitat for both lake trout and lake whitefish (Lake Superior Binational Program Habitat Committee 2006) (Figure 1.1).
- The Goulais River Beach Ridges Conservation Reserve protects an area containing ancient beach ridge landforms. The majority of the site is made up of wetlands, including bogs, fens, swamps and marshes; these wetlands may be of provincial significance. Moose and a number of bird species are often found in the wetlands. Upland forests have developed on the raised beach dunes that are interspersed between the wetlands (OMNR 2005b).
- This regional unit also contains Important Habitat Areas and Important Habitat Sites. Many of the Important Habitat Sites are located near the shore, while some are located further inland (Lake Superior Binational Program Habitat Committee 2006) (Table 1.3, Figure 1.3).
- Portions of the 2,937 hectare Great Lakes Coast – Sault Ste. Marie Enhanced Management Area (EMA) are located within the Goulais Region. This EMA contains islands and parcels of Crown land, located along Lake Superior's eastern coast, and the north shore of the St. Marys River and Lake Huron. This EMA contains geographically diverse areas, including provincially significant wetlands and habitat for fish and wildlife species. Many recreational activities are permitted throughout the area (OMNR 2007a).

Tributaries and Watersheds

- A number of tributaries in the Goulais region were historically used by lake sturgeon for spawning, including the Batchawana River, Chippewa River and Goulais River; the current status of the lake sturgeon populations in these rivers is being investigated. The OMNR is actively engaging in research of lake sturgeon populations in the Batchawana, Chippewa and Goulais Rivers, and Batchawana and Goulais Bays (S. Greenwood, pers. comm., March 11 2013). The Batchawana River population status is extant, while the population trajectory is unknown. The Chippewa River population status and population trajectory are both unknown. The Goulais River population status is extant, while the population status is unknown (Golder Associates Ltd. 2011); however there is not recent evidence of natural reproduction in the Gravel River (Lake Superior Lake Sturgeon Work Group 2012, unpublished data).
- Two additional tributaries in the Goulais regional unit, the Harmony River and Stokely Creek, were identified as historical spawning tributaries for lake sturgeon. The population status for both tributaries is extirpated (Golder Associates Ltd. 2011). There is some uncertainty as to whether these tributaries, which can be described as shallow and flashy, ever supported lake sturgeon spawning. A naming error in a chart for the area, which refers to the Chippewa River as the Harmony River, could have caused some of this uncertainty (S. Greenwood, pers. comm., February 3 2013).
- A Lake Sturgeon Rehabilitation Plan for Lake Superior (Auer 2003) identifies the Batchawana, Chippewa and Goulais Rivers as three of the seventeen tributaries to Lake Superior in which there should be a focus on lake sturgeon rehabilitation.
- The Goulais and Batchawana Rivers are noted historically as important walleye spawning rivers. Walleye stocks were almost extirpated in the 1960's (Hoff 2002) and have not made significant recovery since that time (S. Greenwood, pers. comm., March 11 2013)

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

- The Goulais River is approximately 67 kilometres long, and is classified as a provincial park. The river supports a self-sustaining brook trout population. Some shoreline wetlands are present (OMNR 2006b).
- The Goulais River Beach Ridges Conservation Reserve protects the Goulais River environment, which is described as exceptional in quality (OMNR 2005b). Cranberry Creek is located in the western section of the Goulais River Beach Ridges Conservation Reserve. This creek provides habitat for brook trout and rainbow trout (OMNR 2005b).
- Brook trout populations which are self-sustaining are found in the Batchawana River (OMNR 2006a).

Figure 1.1: Goulais - Coastal and Watershed Features




TABLE 1.2: Goulais CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	NA
Nearshore ¹	B (0.62)	Unknown
Embayments and Inshore ^{1,2}	B (0.67)	Unknown
Coastal Wetlands ^{2,3}	B (0.762)	Unknown
Islands ⁴	A	Unknown
Coastal Terrestrial ³	A (0.955)	Unknown
Tributaries and Watersheds ²	B (0.71)	Unknown

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 1.2: Goulais - Condition



Important Issues & Threats

- The rehabilitation of lean lake trout in nearshore waters of the east end of the lake has not progressed to the same extent as the remainder of the lake. The Ontario Ministry of Natural Resources (OMNR) continues to make efforts to negotiate an agreement with First Nation communities in order to reduce harvests. In the interim, rehabilitation efforts through fish stocking have been discontinued (S. Greenwood, pers. comm., May 27 2013).
- The presence of Emerald Ash Borer (EAB) has been detected within the boundaries of the nearby City of Sault Ste. Marie, Ontario. The entire area of the city is now regulated, meaning a number of regulated articles (e.g. firewood of all species; trees, nursery stock or other materials from the Ash (*Fraxinus*) genus) cannot be moved without permission (CFIA 2012).
- The Prince Wind Farm is a 300 hectare 126 turbine wind farm on the south shore of Goulais Bay. The Prince Wind Farm is operated by Brookfield Renewable Power (Canadian Wind Energy Association 2008). Several other wind farms are in development.
- The St. Marys River Area of Concern (AOC) is located between two regional units, the Goulais and Waika/ Tahquamenon regional units. A number of point and nonpoint sources have contributed to beneficial use impairments, and sediments are contaminated with arsenic, cadmium, chromium, copper, cyanide and lead. Ten beneficial use impairments were identified in the St. Marys River AOC (U.S. EPA 2013a).

Conservation In Action

Parks & Protected Areas

- Algoma Headwaters Provincial Park
- Goulais River Provincial Park
- Batchawana River Provincial Park
- Batchawana Bay Provincial Park
- Pancake Bay Provincial Park
- Sandy Islands Provincial Park
- Lake Superior Provincial Park Addition
- Goulais River Beach Ridges Conservation Reserve
- Ile Parisienne Conservation Reserve
- Great Lakes Coast - Sault Ste. Marie Enhanced Management Area

Existing Programs & Projects


- Peregrine falcons have been reintroduced to Ontario, including in the Lake Superior basin, following their extirpation as a breeding species in Ontario in the early 1960s (Ontario Peregrine Falcon Recovery Team 2010). The Ontario Ministry of Natural Resources (OMNR), Canadian Wildlife Service and many naturalist organizations and corporations have been involved in the re-establishment of peregrine falcons across Ontario (Ontario Peregrine Falcon Recovery Team 2010). Project Peregrine is a project of the Thunder Bay Field Naturalists, supported by the OMNR. Established in 1989, Project Peregrine now conducts an intensive monitoring program with volunteers and OMNR staff in a number of areas throughout the Ontario portion of the Lake Superior Basin (Thunder Bay Field Naturalists No Date, Ontario Peregrine Falcon Recovery Team 2010). In the 2010 Ontario Peregrine Falcon survey, 72 of the 119 identified territories in Ontario were located in the Lake Superior basin. In Ontario the Lake Superior basin is the location of the highest quality of cliff nesting sites and supports the highest increase in the number of territories, and the highest density of cliff nesting birds (Chikoski and Nyman 2011). Peregrine falcons born and banded in Ontario have also been observed nesting in Minnesota, Wisconsin and Michigan (Redig et

al. 2010 as cited in Chikoski and Nyman 2011). Peregrine falcons are listed as Special Concern by the Committee on the Status of Species at Risk in Ontario (COSSARO) and the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) (OMNR 2013a, COSEWIC 2011a).

TABLE 1.3: Goulais IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
ON-005	Site	Batchawana Island	Coastal wetland; rare animal habitat, migratory bird habitat
ON-006	Site	Batchawana River	Fish spawning area
ON-007	Site	Batchawana Bay	Staging area and brood habitat for migratory waterfowl
ON-007	Area	Batchawana Bay	Staging area and brood habitat for migratory waterfowl
ON-015	Site	Bojack and Bone	Nesting site for water birds
ON-019	Site	Chippewa River	Excellent moose habitat, little access
ON-024	Site	Deadman's Cove	Wetland, fen with rare plants
ON-033	Site	Flowerpot Islands	Colonial water birds
ON-041	Area	Goulais River Delta	Fish spawning area, rare species habitat
ON-044	Site	Harmony River	Fish spawning habitat (largest run of rainbow smelt in area) feeding area for heron
ON-050	Site	King Mountain	High biodiversity value; old growth forest pockets; representative landscape
ON-056	Site	Marlette's Bay	Waterfowl staging area, brood habitat
ON-070	Site	Chippewa River Mouth	Fish spawning habitat
ON-092	Site	Pancake River	Fish spawning area
ON-102	Site	Robertson Cliffs	Rare animal habitat
ON-104	Site	Sand Point	Yellow perch habitat; perch numbers declining
ON-126	Site	Turkey Lakes	Old growth maple and birch; high biodiversity value (wildlife); roadless area; fish spawning area
ON-128	Site	Upper St. Mary's River	Fish spawning area
ON-128	Area	Upper St. Mary's River	Fish spawning area
ON-130	Site	Whiskey Point	Significant staging area for waterfowl
ON-132	Site	Wily Lake	Fish spawning area
ON-136	Site	Gros Cap Corridor	Migratory fish habitat; commercial fishery; colonial water bird habitat; Gros Cap reef

Figure 1.3: Goulais – Important Habitat Sites and Areas



2. Agawa/ Michipicoten – Magpie

HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	B	COASTAL WETLANDS	A-
EMBAYMENTS & INSHORE	B	COASTAL TERRESTRIAL	A+
TRIBUTARIES & WATERSHEDS	A	OVERALL A	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



Sunset over Lake Superior shoreline in Lake Superior Provincial Park. Photo credit: Ethan Meleg

Summary/ Description

The Agawa/ Michipicoten – Magpie unit is located in Ontario on the eastern shore of Lake Superior, and contains the easternmost portion of the Lake Superior basin. Including the associated nearshore waters, this regional unit is 14,413.9 km² in size. This regional unit has its southern boundary near the Montreal River, while its northern boundary is near Pilot Harbour. Lake Superior Provincial Park, a 155,647 ha natural environment class park, is located in this regional unit, along the Lake Superior shore (OMNR 2006d). Communities in this regional unit include Dubreuilville, Hawk Junction, Montreal River, Missanabie, Michipicoten First Nation, Michipicoten River and Wawa. The Agawa/ Michipicoten-Magpie unit combines two tertiary watersheds, the Agawa and the Michipicoten-Magpie, and contains 22 quaternary watersheds. The watersheds are dominated by forests - developed and agricultural lands are very limited. The coast is characterized by rocky shores and cliffs. Coastal wetlands and sand beaches are very rare in this region. Lake Superior Provincial Park and several other parks protect almost two-thirds of the coast in this region.

TABLE 2.1: Agawa/ Michipicoten- Magpie BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
<i>Agriculture</i>	6.40	0.04	1,287.56	
<i>Developed</i>	2.03	0.01	351.99	
<i>Forest</i>	12,076.90	83.79	96,789.30	
<i>Associated Nearshore Waters</i>	826.48	5.73	15,951.22	
<i>Other</i>	939.41	6.52	7,452.78	
<i>Water (inland)</i>	562.70	3.90	8,283.49	
Total Area	14,413.9	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
<i>Coastline (km)</i>	346.1	NA	6.09	Based on SOLEC shoreline
<i>Sand Beaches (km)</i>	35.20	10.17	5.59*	*% of Lake Superior Total Sand Beaches
<i>Coastal Wetlands (km²)</i>	2.97	0.82*	0.30**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
<i>Natural Cover in Coastal Zone</i>	354.58	97.24*	6.42**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
<i>Number of Islands</i>	291	NA	11.0	
Condition	Region	Region %	% of Lake Superior Total	
<i>Population Density (persons/km²)</i>	0.02	NA		
<i>Road Density (km/km²)</i>	0.21	NA		
<i>Number of Dams and Barriers</i>	1,270	NA	5.4	
<i>Artificial Shoreline (km)</i>	0.73	0.20	0.33	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
<i>Private</i>	2,770.11	20.41	13,571.78	Regional area based on landmass
<i>Public/Crown</i>	7,652.45	56.39	13,571.78	
<i>Tribes/ First Nations</i>	32.21	0.24	13,571.78	
<i>Parks & Protected Areas (total)</i>	3,244.76	23.91	13,571.78	
<i>Parks & Protected Areas (coast)</i>	226.22	62.04*	364.63**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- The Agawa/ Michipicoten – Magpie regional unit contains sites of Important Habitat for both lake trout and lake whitefish. Important Habitat sites for lake trout are found off the coast of the Agawa/ Michipicoten – Magpie region, in the inshore and nearshore zones (Lake Superior Binational Program Habitat Committee 2006) (Figure 2.1).
- Michipicoten Bay is noted as a Lake Superior embayment important for lake sturgeon (Auer 2003). In the Agawa/ Michipicoten-Magpie regional unit this embayment and the nearshore zone, which provides corridors for movement, are identified as critical management areas for lake sturgeon in the Lake Superior basin (Auer 2003).
- An offshore cobble bar complex is located at Montreal River Provincial Nature Reserve. This cobble bar formed 6,000 years ago, when glacial lakes were present. Under present day conditions the cobble bar is 60 metres above Lake Superior water levels (OMNR 2006e).

Coastal Zone and Islands

- Leach Island, South Lizard Island and Rowe Island (islands which are part of Lake Superior Provincial Park) are noted as an Important Habitat site for lake trout (Lake Superior Binational Program Habitat Committee 2006) (Figure 2.1).
- The Agawa/ Michipicoten-Magpie regional unit contains Important Habitat Areas, including one area which extends along the coast and into the White and Pic regional unit. Several Important Habitat Sites are also found in this region, including along the shore, inland, and on Leach Island and Montreal Island (part of Lake Superior Provincial Park) (Lake Superior Binational Program Habitat Committee 2006) (Table 2.3, Figure 2.3).
- The McGregor Cove natural heritage area contains arctic coastal disjunct species (OMNR 2006c) and the Lake Superior shoreline at Montreal River Provincial Nature Reserve also contains two kilometres of habitat suitable for arctic alpine plant species (OMNR 2006e).
- The shoreline of Lake Superior at Montreal River Provincial Nature Reserve is noted to be a high energy cobble beach (OMNR 2006e).
- Driftwood Beach at Michipicoten Provincial Park is considered a significant landscape feature (OMNR 2006g).
- 170 taxa have been recorded in Michipicoten Provincial Park (Thompson 1994 as cited in OMNR 2004a). Two of the plant species were provincially significant, and six of the species were regionally significant (Thompson 1994 as cited in OMNR 2004a).
- 34 migratory bird species and 4 year round resident bird species have been recorded in Michipicoten Provincial Park (OMNR 2004a).

Tributaries and Watersheds

- The fish community on the lower Michipicoten River is largely comprised of introduced species, including rainbow trout, chinook salmon, coho salmon and pink salmon. Less abundant are native fish species, including walleye, lake trout, lake sturgeon and brook trout (Eason 2003 as cited in OMNR 2004a).
- Historically 21 tributaries in Lake Superior had lake sturgeon spawning runs. One of these historical spawning tributaries, the Michipicoten River, is in the Agawa/ Michipicoten-Magpie regional unit. The population status and population trajectory are both considered unknown (Golder Associates Ltd. 2011); however there is no recent evidence of lake sturgeon spawning in the Michipicoten River (Lake Superior Lake Sturgeon Work Group 2012, unpublished data).
- A Lake Sturgeon Rehabilitation Plan for Lake Superior (Auer 2003) identifies the Michipicoten River as one of the seventeen tributaries to Lake Superior in which there should be a focus on lake sturgeon rehabilitation. Habitat restoration is a priority in the Michipicoten River, due to barriers to migration and spawning (Auer 2003).
- The Michipicoten River mouth is a site for bald eagles during fall salmon spawning (OMNR 2004a).
- The rivers flowing through Lake Superior Provincial Park are noted to drop rapidly as they flow from the interior highlands to Lake Superior. This is noted to create rapids and waterfalls (OMNR 2006d).
- The Lake Superior Highlands Recommended Conservation Reserve is a 54,007 hectare area along the northeastern Lake Superior coast, in the White and Pic and Agawa/ Michipicoten-Magpie regional units. The boundary of this unit was extended to include a waterway which would connect the Lake Superior shoreline. This area also provides critical habitat for caribou (OMNR 2006h).
- The Nimoosh Provincial Park combines a number of rivers, parks, and life and earth science areas. The Nimoosh River links Obtonaga Provincial Park with the Lake Superior shore (OMNR 2006i). Areas of important spawning habitat for lake trout are located in the Dog River (OMNR 2006i).

Figure 2.1: Agawa/ Michipicoten – Magpie - Coastal and Watershed Features




TABLE 2.2: Agawa/ Michipicoten – Magpie CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	NA
Nearshore ¹	B (0.62)	Unknown
Embayments and Inshore ^{1,2}	B (0.74)	Unknown
Coastal Wetlands ^{2,3}	A- (0.825)	Unknown
Islands ⁴	A	Unknown
Coastal Terrestrial ³	A+ (0.995)	Unknown
Tributaries and Watersheds ²	A (0.86)	Unknown

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 2.2: Agawa/ Michipicoten – Magpie - Condition



Important Issues & Threats

- The rehabilitation of lean lake trout in nearshore waters of the east end of the lake has not progressed to the same extent as the remainder of the lake. The Ontario Ministry of Natural Resources (OMNR) continues to make efforts to negotiate an agreement with First Nation communities in order to reduce harvests. In the interim, rehabilitation efforts through fish stocking have been discontinued (S. Greenwood, pers. comm., May 27 2013).
- Twenty-one of the species identified in Michipicoten Post Provincial Park were non-native species (Thompson 1994 as cited in OMNR 2004a).
- The Michipicoten River and Magpie River were both reported to have a number of dams. Four power dams and four storage dams were reported on the Michipicoten River system. Three power dams and one storage dam were reported to be in operation on the Magpie River (OMNR 2004a).
- Recreational use of Michipicoten Post Provincial Park is generally compatible, however some activities are incompatible, and some areas are more sensitive to activities. The majority of park use occurs in and around Driftwood Beach, in the nearby coastal forest and in the waters of Lake Superior. Driftwood Beach is a sandy beach community that is both well-used and sensitive to recreational use. All-terrain and off-road vehicle use on the beach and in the forest is significant. Camping, relic hunting, and other human activities have led to some degradation of areas of Michipicoten Post Provincial Park (OMNR 2004a).
- Portions of the South Michipicoten River – Superior Shoreline Conservation Reserve are subject to mining claims and leases. There is low to high mineral potential in these sites; if the mining claims are surrendered the claim and lease areas will be added to the conservation reserve lands (OMNR 2006f).

Conservation In Action

Parks & Protected Areas

- Lake Superior Provincial Park
- Wenebagon River Provincial Park
- The Shoals Provincial Park
- Potholes Provincial Nature Reserve
- Michipicoten Post Provincial Park
- Nimoosh Provincial Park
- Obatanga Provincial Park
- Montreal River Provincial Nature Reserve
- South Michipicoten River – Superior Shoreline Conservation Reserve
- South Michipicoten River Forest Reserve
- Lake Superior Highlands Recommended Conservation Reserve (Recommended)

Existing Programs & Projects

- The Magpie and Michipicoten Rivers were the focus of several research projects examining the ecological effects of hydropower peaking (OMNR No date a). Hydropower peaking can be described as the variable flows in rivers resulting from storing waters in reservoirs during non-peak periods and the release of water from reservoirs to produce power during peak hours. The alterations to river flow did not necessarily result in a decrease in productive fish habitat, but some fishes and invertebrates are constrained by the altered flow regime (OMNR No date a). Implications for river management include treating the high and low flows in hydropower peaking rivers as two rivers, and acknowledging the longitudinal and latitudinal gradients in hydropower peaking rivers (OMNR No date a).


Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

- The Michipicoten River system is the focus of a water management plan. In the lower reach of the river, the plan will focus on increasing the minimum flow from the lowest hydro dam (OMNR 2004a).
- Fall aerial helicopter flight counts for spawning chinook salmon were conducted on the Michipicoten River from 1987 to 2000 (Greenwood 2000). Volunteers have continued to count spawning Chinook salmon, but the data for these counts are not known (S. Greenwood, pers. comm., April 22 2013).
- Peregrine falcons have been reintroduced to Ontario, including in the Lake Superior basin, following their extirpation as a breeding species in Ontario in the early 1960s (Ontario Peregrine Falcon Recovery Team 2010). The Ontario Ministry of Natural Resources (OMNR), Canadian Wildlife Service and many naturalist organizations and corporations have been involved in the re-establishment of peregrine falcons across Ontario (Ontario Peregrine Falcon Recovery Team 2010). Project Peregrine is a project of the Thunder Bay Field Naturalists, supported by the OMNR. Established in 1989, Project Peregrine now conducts an intensive monitoring program with volunteers and OMNR staff in a number of areas throughout the Ontario portion of the Lake Superior Basin (Thunder Bay Field Naturalists No Date, Ontario Peregrine Falcon Recovery Team 2010). In the 2010 Ontario Peregrine Falcon survey, 72 of the 119 identified territories in Ontario were located in the Lake Superior basin. In Ontario the Lake Superior basin is the location of the highest quality of cliff nesting sites and supports the highest increase in the number of territories, and the highest density of cliff nesting birds (Chikoski and Nyman 2011). Peregrine falcons born and banded in Ontario have also been observed nesting in Minnesota, Wisconsin and Michigan (Redig et al. 2010 as cited in Chikoski and Nyman 2011). Peregrine falcons are listed as Special Concern by the Committee on the Status of Species at Risk in Ontario (COSSARO) and the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) (OMNR 2013a, COSEWIC 2011a).

TABLE 2.3: Agawa/ Michipicoten – Magpie IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
ON-001	Site	Michipicoten Corridor	Habitat for rare plants (arctic disjuncts) and animals; rocky outcrops and shallow soils with acid sensitivity
ON-001	Area	Michipicoten Corridor	Habitat for rare plants (arctic disjuncts) and animals; rocky outcrops and shallow soils with acid sensitivity
ON-025	Site	Dog River System	Fish spawning area (Dog R. strain of lake trout is one of few river-spawning populations); fish habitat; waterfowl breeding/staging, rare plant habitat
ON-029	Site	Dove Bay	Fish spawning area
ON-031	Site	Eagle River	Fish habitat
ON-035	Site	Wawa Fume Kill	High biodiversity values
ON-042	Site	Gravel Beach	Fish spawning area
ON-052	Site	Leach Island	Rare animal habitat (unoccupied)
ON-060	Site	Megason Lake	Old growth pockets, roadless area; provincially significant wetlands; fish spawning area, headwaters for several rivers
ON-062	Site	Michipicoten Harbor	Fish spawning area
ON-065	Site	Michipicoten River	Fish spawning area/habitat; coastal wetland/estuary
ON-066	Site	Lower Michipicoten River	Fish spawning habitat
ON-068	Site	Montreal Island	Former populations of woodland caribou
ON-069	Site	Montreal River Nature Reserve	Excellent moose habitat
ON-069	Area	Montreal River Nature Reserve	Excellent moose habitat
ON-075	Site	Montreal River Mouth	Fish spawning habitat
ON-105	Site	Sandy Beach	Shorebird habitat, dune habitat
ON-108	Site	Montreal Shoreline North	Rare arctic plant habitat
ON-124	Site	The Flats	Rare animal habitat
ON-135	Site	Wolf-Achigan Lakes	Old growth forest, high biodiversity value (landscape level), fish habitat, roadless area
ON-136	Area	Gros Cap Corridor	Migratory fish habitat; commercial fishery; colonial water bird habitat; Gros Cap reef
ON-168	Area	Potholes Nature Reserve	

Figure 2.3: Agawa/ Michipicoten – Magpie - Important Habitat Sites and Areas



3. White and Pic

HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	B	COASTAL WETLANDS	A
EMBAYMENTS & INSHORE	A	COASTAL TERRESTRIAL	A+
TRIBUTARIES & WATERSHEDS	A	OVERALL A	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



Pukaskwa Depot Beach. Photo credit: Sue Greenwood/ Ontario Ministry of Natural Resources

Summary/ Description

The White and Pic regional unit is located on the northeastern shore of Lake Superior, and including the associated nearshore waters this regional unit is 13,098.83 km² in size. The regional boundaries for this unit are north of the Pic River (and just south of Marathon) in the east, to just west of Pilot Harbour. Pukaskwa National Park, an 187,800 hectare wilderness national park of boreal forest is located along the Lake Superior coast (Parks Canada 2013). Communities in this regional unit include Ojibways of the Pic River First Nation, Heron Bay, Mobert, Pic Mobert First Nation, Hillsport, Manitouwadge and White River. The White and Pic regional unit combines two tertiary watersheds, White and Pic, and contains 19 quaternary watersheds. The watersheds dominated by forests, including protected forests within Pukaskwa National Park. The coast is characterized by rocky shores and cliffs, with scattered nearshore islands. Coastal wetlands and sand beaches are very rare in this region. Pukaskwa National Park protects over 70% of the coast in this region.

TABLE 3.1: White and Pic BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
<i>Agriculture</i>	1.71	0.01	1,287.56	
<i>Developed</i>	1.31	0.01	351.99	
<i>Forest</i>	11,882.15	90.71	96,789.30	
<i>Associated Nearshore Waters</i>	207.00	1.58	15,951.22	
<i>Other</i>	578.26	4.41	7,452.78	
<i>Water (inland)</i>	428.42	3.27	8,283.49	
Total Area	13,098.83	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
<i>Coastline (km)</i>	371.1	NA	6.52	Based on SOLEC shoreline
<i>Sand Beaches (km)</i>	18.30	4.93	2.91*	*% of Lake Superior Total Sand Beaches
<i>Coastal Wetlands (km²)</i>	2.77	1.10*	0.28**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
<i>Natural Cover in Coastal Zone</i>	240.30	95.59*	4.35**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
<i>Number of Islands</i>	387	NA	14.6	
Condition	Region	Region %	% of Lake Superior Total	
<i>Population Density (persons/km²)</i>	0.14	NA		
<i>Road Density (km/km²)</i>	0.08	NA		
<i>Number of Dams and Barriers</i>	486	NA	2.1	
<i>Artificial Shoreline (km)</i>	0.13	0.06	0.06	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
<i>Private</i>	435.88	3.38	12,885.94	Regional area based on landmass
<i>Public/Crown</i>	10,536.4	81.77	12,885.94	
<i>Tribes/ First Nations</i>	4.06	0.03	12,885.94	
<i>Parks & Protected Areas (total)</i>	1,920.87	14.91	12,885.94	
<i>Parks & Protected Areas (coast)</i>	176.87	70.35*	251.39**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- The White/Pic regional unit contains a number of sites of Important Habitat for lake trout; these Important Habitat sites are found in many areas along the coast (Lake Superior Binational Program Habitat Committee 2006) (Figure 3.1).

Coastal Zone and Islands

- This regional unit contains Important Habitat Areas, including one large Important Habitat Area which is Pukaskwa National Park. Another Important Habitat Area along the coast extends from the Agawa/ Michipicoten – Magpie regional unit into the White Pic regional unit. Smaller Important

Habitat Areas are also located in this regional unit. Several Important Habitat Sites are also found in this region (Lake Superior Binational Program Habitat Committee 2006) (Table 3.3, Figure 3.3).

- Arctic-alpine disjunct species can be found in Pukaskwa National Park (Parks Canada 2013).
- Woodland caribou are still found within Pukaskwa National Park, an area they have inhabited since the retreat of the last glaciers (Parks Canada 2013).
- Craig's Pit Provincial Nature Reserve is an important migratory bird observation area, and contains bluffs and kettle holes. This area also contains examples of landform processes and themes which are environmental indicators from the Lake Minong stage and later (OMNR 2006l).

Tributaries and Watersheds

- Historically 21 tributaries in Lake Superior had lake sturgeon spawning runs (Lake Superior Lake Sturgeon Work Group 2012, unpublished data). Two of these historical spawning tributaries, the White River (Ontario) and the Pic River are in the White and Pic regional unit. The White River population status and population trajectory are both listed as unknown (Golder Associates Ltd. 2011); however, recent research in 2011 and 2012 suggests that the population status in the White River is extant (Ecclestone 2013). The population status is still currently described as unknown (Lake Superior Lake Sturgeon Work Group 2012, unpublished data). Ongoing research indicates lake sturgeon abundance, sex ratio, natural recruitment and year class structure are approaching criteria for self-sustaining classification, based on the criteria listed in the Lake Sturgeon Rehabilitation Plan for Lake Superior (Ecclestone 2013).
- The White River (Ontario) is one of ten Lake Superior tributaries where lake sturgeon have currently been documented spawning (as of 2012); this is the same number as 2005, however the specific tributaries have changed (Lake Superior Lake Sturgeon Work Group 2012, unpublished data). The White River (Ontario) and the St. Louis River (Minnesota) have recent evidence of natural reproduction, while there is not recent evidence from the Gravel and Michipicoten rivers. The White River (Wisconsin) had been removed as it is a tributary to the Bad River and is not a separate spawning population (Lake Superior Lake Sturgeon Work Group 2012, unpublished data).
- The Pic River population status is extant, while the population trajectory is unknown (Golder Associates Ltd. 2011). Although recent research has indicated that the Pic River lake sturgeon population abundance is relatively low and the sex ratio is unequal, a total of 24 year classes are present in the system and natural recruitment has been documented at Manitou and Kagiano Falls (Ecclestone et al. 2013).
- A Lake Sturgeon Rehabilitation Plan for Lake Superior (Auer 2003) identifies the White and Pic (identified as the Big Pic) Rivers as two of the seventeen tributaries to Lake Superior in which there should be a focus on lake sturgeon rehabilitation.
- The Lake Superior Highlands Recommended Conservation Reserve is a 54,007 hectare area along the northeastern Lake Superior coast, in the White and Pic and Agawa/ Michipicoten-Magpie regional units. The boundary of this unit was extended to include a waterway which would help allow gene flow to the Lake Superior shoreline. This area also provides critical habitat for caribou (OMNR 2006h).
- Pukaskwa River Provincial Park is linked to Pukaskwa National Park. It is described as free-flowing, and with spectacular scenery and geology (OMNR 2006j).
- The Pokei Lake / White River Wetlands Provincial Park area includes riparian wetlands in the floodplain of the White River. The area of the wetlands is described as extensive, and includes marsh, fen and various swamps. The site is also noted for backwater ponds along the river which are suitable for waterfowl, eagle nesting sites, and caribou have been viewed in the area (OMNR 2006k).

Figure 3.1: White and Pic - Coastal and Watershed Features




TABLE 3.2: White and Pic CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	NA
Nearshore ¹	B (0.73)	Unknown
Embayments and Inshore ^{1,2}	A (0.80)	Unknown
Coastal Wetlands ^{2,3}	A (0.863)	Unknown
Islands ⁴	A	Unknown
Coastal Terrestrial ³	A+ (1.000)	Unknown
Tributaries and Watersheds ²	A (0.86)	Unknown

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 3.2: White and Pic - Condition



Important Issues & Threats

- An EA has been submitted for a metals and copper mine project near the north shore of Lake Superior, approximately 10 kilometres north of the community of Marathon. The Marathon Project (Marathon Platinum Group Metals-Copper Project) is a project of Stillwater Canada Inc., a subsidiary of Stillwater Mining Company.
- The rehabilitation of lean lake trout in nearshore waters of the east end of the lake has not progressed to the same extent as the remainder of the lake. The Ontario Ministry of Natural Resources (OMNR) continues to make efforts to negotiate an agreement with First Nation communities in order to reduce harvests. In the interim, rehabilitation efforts through fish stocking have been discontinued (S. Greenwood, pers. comm., May 27 2013).
- Although there has been no logging in any part of Pukaskwa National Park since the early 1900s, logging on lands adjacent to Pukaskwa National Park has been shown to significantly influence populations of animals within the boundaries of the park (Parks Canada 2013).
- One of the largest gold mines in North America opened 20 kilometres north of Pukaskwa National Park, shortly after the park was officially established (Parks Canada 2012).

Conservation In Action

Parks & Protected Areas

- Pukaskwa National Park
- Pokei Lake White River Wetlands Provincial Park
- White Lake Provincial Park
- White Lake Peatlands Provincial Nature Reserve
- Pen Lake Fen Provincial Park
- Craig's Pit Provincial Nature Reserve (also in Little Pic regional unit)
- Lake Superior Shoreline Enhanced Management Area (also in Little Pic and Jack Pine/Nipigon regional units)


Existing Programs & Projects

- The Friends of Pukaskwa is a volunteer organization which supports the philosophy and purpose of Pukaskwa National Park (Friends of Pukaskwa National Park No date).
- The Ojibways of the Pic River First Nation Reserve is located near the mouth of the Pic River, and near the north end of Pukaskwa National Park. The Lands and Resources Department of the Pic River First Nation has recently participated in the Environmental Review Panel for the Stillwater Mine, in land-use mapping, and in ongoing meetings with Pukaskwa National Park for the Park Management Plan and Park Advisory Committee (Ojibways of the Pic River First Nation 2012).
- Peregrine falcons have been reintroduced to Ontario, including in the Lake Superior basin, following their extirpation as a breeding species in Ontario in the early 1960s (Ontario Peregrine Falcon Recovery Team 2010).

TABLE 3.3: White and Pic IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
ON-001	Area	Michipicoten Corridor	Habitat for rare plants (arctic disjuncts) and animals; rocky outcrops and shallow soils with acid sensitivity
ON-037	Site	Redsucker Cove	Fish spawning habitat
ON-055	Site	Maple, Hilltop, and Jackfish Lakes	Fish spawning habitat
ON-084	Site	North Skipper Lake	
ON-097	Site	Pipe River Watershed	Excellent moose habitat; fish spawning area
ON-099	Site	Jarvey Lake	Fish habitat
ON-115	Site	Ogilvy Point Islands	Colonial water bird habitat
ON-122	Site	Starr Island	Colonial water bird habitat
ON-172	Area	Pukaskwa National Park	Large intact protected area, woodland caribou population, sand dunes, rare plant habitat
ON-185	Area	White Lake Peatlands Nature Reserve	

Figure 3.3: White and Pic - Important Habitat Sites and Areas



4. Michipicoten Island

HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	A	COASTAL WETLANDS	A
EMBAYMENTS & INSHORE	A	COASTAL TERRESTRIAL	A+
TRIBUTARIES & WATERSHEDS	A	OVERALL A	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



Photo of Caribou Island Lighthouse in the distance taken from Caribou Island. Photo credit Bruce Tomlinson, Ontario Ministry of Natural Resources (OMNR). Photo supplied by Sue Greenwood, OMNR

Summary/ Description

The Michipicoten Island regional unit contains Michipicoten Island and Caribou Island, as well as other smaller islands and island complexes. This regional unit measures 783.66 km² in size, including the associated nearshore waters. Michipicoten Island is located in the northeastern portion of Lake Superior. It is the third largest island in the lake, and is approximately 26 by 10 km wide (from north to south) (OMNR 2004a). The topography of the island is variable, with close to 300 m in topographic relief from the Lake Superior shoreline (181 m above sea level) to the island's highest elevation (479 m above sea level) (OMNR 2004a). First nations of the Lake Superior area gave the island the name Missipacouatong, which means "land of the big bluffs" (Annells 1974 as cited in OMNR 2004a). The closest mainland to Michipicoten Island is approximately 16 kilometres to the north, and the community of Wawa is approximately 65 kilometres to the northeast (OMNR 2004a). No moose, deer, bear or wolves occur on Michipicoten Island, but beaver and woodland caribou are abundant (COSEWIC 2011b, OMNR 2013b, 2004a). Approximately 40 kilometres south of Michipicoten Island is the 6.09 km² Caribou Island. Caribou Island is the most isolated island in the Great Lakes, and it is also the most isolated freshwater island in the world (Henson et al. 2010). Michipicoten Island regional unit does not contain any tertiary or quaternary watershed units, but is comprised of islands and island complexes. While Michipicoten Island is characterized by high cliffs and cobble beaches, Caribou Island is generally flat and has large sand beaches. Much of Michipicoten Island is provincial park, except for some small private and federal parcels. Caribou Island is privately owned by a U.S. foundation.

TABLE 4.1: Michipicoten Island BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
<i>Agriculture</i>	0.08	0.01	1,287.56	
<i>Developed</i>	0.11	0.01	351.99	
<i>Forest</i>	170.11	21.71	96,789.30	
<i>Associated Nearshore Waters</i>	608.07	77.59	15,951.22	
<i>Other</i>	1.83	0.23	7,452.78	
<i>Water (inland)</i>	3.45	0.44	8,283.49	
Total Area	783.66	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
<i>Coastline (km)</i>	128.1	NA	2.25	Based on SOLEC shoreline
<i>Sand Beaches (km)</i>	11.20	8.74	1.78*	*% of Lake Superior Total Sand Beaches
<i>Coastal Wetlands (km²)</i>	0.79	0.67*	0.08**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
<i>Natural Cover in Coastal Zone</i>	111.25	93.95*	2.01**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
<i>Number of Islands</i>	83	NA	3.1	
Condition	Region	Region %	% of Lake Superior Total	
<i>Population Density (persons/km²)</i>	0.00	NA		
<i>Road Density (km/km²)</i>	0	NA		
<i>Number of Dams and Barriers</i>	0	NA	0	
<i>Artificial Shoreline (km)</i>	0.18	0.16	0.08	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
<i>Private</i>	6.13	3.51	174.43	Regional area based on landmass
<i>Public/Crown</i>	0.43	0.24	174.43	
<i>Tribes/ First Nations</i>	0.00	0.00	174.43	
<i>Parks & Protected Areas (total)</i>	168.30	96.49	174.43	
<i>Parks & Protected Areas (coast)</i>	112.28	94.83*	118.41**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- Spawning beds for lake trout are known to be present in the Lake Superior portion of Michipicoten Island Provincial Park (OMNR 2004a).
- The fish community in the waters of Michipicoten Island Provincial Park includes lake trout, coho salmon, chinook salmon, cisco and whitefish. These species support a sport fishing fishery, especially in areas along the southern and southwestern shore of Michipicoten Island (OMNR 2004a).
- The provincially rare pygmy whitefish is found in the eastern waters of Lake Superior, around Michipicoten Island (Petzold 2003 as cited in OMNR 2004a).

Coastal Zone and Islands

- The Michipicoten Island regional unit contains sites of Important Habitat for lake trout and whitefish (Figure 4.1). Lake trout spawning sites are noted around the Breeder Islands just south west of Michipicoten Island and in Quebec Harbour (Goodier 1982) located on the south shore. Lake Whitefish are also noted as spawning in Quebec Harbour. Lake trout spawning is noted on Caribou Island and commercial fishers report whitefish spawning along the southwest shore of Caribou Island prior to 1955 (Lake Superior Binational Program Habitat Committee 2006, Goodier 1982).
- Michipicoten Island is considered to be an Important Habitat Area (Lake Superior Binational Program Habitat Committee 2006) (Table 4.3, Figure 4.3).
- Much of Michipicoten Island's shoreline is bare bedrock knolls, exposed to wave action. The shoreline is also composed of beach deposits of sand, pebble and cobble (OMNR 2004a).
- Michipicoten Island's present shoreline was established approximately 2,000 years ago (OMNR 2004a).
- East Sand Bay and West Sand Bay are provincially rare sand beach communities, and are classified as provincially significant vegetation communities (White 2000 as cited in OMNR 2004).
- The Rocky Shoreline habitats of Michipicoten Island support provincially and regionally significant plants, and are classified as provincially significant vegetation communities (White 2000 as cited in OMNR 2004).
- The Quebec Harbour Wetland Complex is regionally significant, and is the most extensive and diverse wetland on Michipicoten Island, containing a variety of wetland communities (White 2000 as cited in OMNR 2004a).
- The Caribou Island shoreline has large sections of sandy beaches and dunes (Henson et al. 2010). Wetlands occur between the ridges of the dunes, and open bogs, treed muskegs and forested dunes are also present (Liebermann 1998 as cited in Henson et al. 2010).
- The topographic relief found on Michipicoten Island contributes to habitat diversity, and allows southern and northern plant species to be found in relatively close proximity to one another (OMNR 2004a). Many of these species are arctic-alpine disjuncts, or at the extent of their geographic range. Ten provincially significant and 53 regionally significant plant species have been identified (OMNR 2004a).

Tributaries and Watersheds

- The Interior Lake Shorelines of Michipicoten Island are classified as provincially significant vegetation communities (White 2000 as cited in OMNR 2004). They are described as "lakes of little disturbance on an isolated island exposed to the climatic influence of Lake Superior" (OMNR 2004a:12). There are over 20 inland lakes on Michipicoten Island (OMNR 2004a).
- The presence of an abundant beaver population has altered the natural hydrology of Michipicoten Island, with many areas flooded (OMNR 2004a).
- Approximately 87% of Michipicoten Island is forested (Noble 1984 as cited in OMNR 2004a); remaining areas are water, non-forested wetlands, beaches, talus slopes and rock outcrops (OMNR 2004a).

Figure 4.1: Michipicoten Island - Coastal and Watershed Features




TABLE 4.2: Michipicoten Island CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	NA
Nearshore ¹	A (0.90)	Unknown
Embayments and Inshore ^{1,2}	A (0.90)	Unknown
Coastal Wetlands ^{2,3}	A (0.933)	Unknown
Islands ⁴	A	Unknown
Coastal Terrestrial ³	A+ (0.998)	Unknown
Tributaries and Watersheds ²	A (0.90)	Unknown

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 4.2: Michipicoten Island - Condition



Important Issues & Threats

- Michipicoten Island Provincial Park's sand beaches are sought out by visitors to the island for recreational use, and they are especially sought out given the rarity of the beaches on the island (OMNR 2004a). These sand beach communities contain significant and sensitive species, and are also considered the most sensitive feature to recreation use (OMNR 2004a).
- The shorelines of Michipicoten Island Provincial Park are the focus of recreational activities. Many of these shorelines contain significant species; recreation and interpretation must be weighed against constraints to manage the areas (OMNR 2004a).
- 170 acres in the southeast portion of Michipicoten Island are listed for sale. The property includes over 9,300 feet of Lake Superior coastline, including pebble and boulder beaches, and two small islands (Apex Realty Services Ltd. 2013).

Conservation In Action

Parks & Protected Areas

- Michipicoten Island Provincial Park: The boundaries of Michipicoten Island Provincial Park extend 2.5 kilometres offshore of Michipicoten Island, and the park designation applies to the islands and shoals within that area (OMNR 2004a). Some private and federal lands within the park boundaries are excluded from the provincial park and related policies (OMNR 2004a).


Existing Programs & Projects

- Michipicoten Island is the source of one of two lake trout brood stock gametes (Michipicoten strain) used by the provincial fish culture system for lake trout rehabilitation stocking in Lake Superior and Lake Huron (S. Greenwood, pers. comm., March 11 2013).
- The caribou population found on Michipicoten Island today is the result of a successful 1982 translocation of eight animals from Slate Islands Provincial Park (Ontario Woodland Caribou Recovery Team 2008), in addition to the one adult male already present on the island (believed to be from the Pukaskwa herd) (Gogan and Cochrane 1994). The caribou population that was historically present on the island was extirpated in the late 1800s, when there were active copper mining operations on the island. The current population is estimated to be between 400 and 500 individuals (S. Lebel and G. Eason, pers. comm., June 18 2013).
- Peregrine falcons have been reintroduced to Ontario, including in the Lake Superior basin, following their extirpation as a breeding species in Ontario in the early 1960s (Ontario Peregrine Falcon Recovery Team 2010).
- Caribou Island has been on the market in the past and may be available for purchase in the future.

TABLE 4.3: Michipicoten Island IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
ON-162	Area	Michipicoten Island	Provincial Park; fish spawning

Figure 4.3: Michipicoten Island - Important Habitat Sites and Areas



5. Little Pic


HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	C	COASTAL WETLANDS	B
EMBAYMENTS & INSHORE	B	COASTAL TERRESTRIAL	A+
TRIBUTARIES & WATERSHEDS	B	OVERALL B	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



Woodland caribou inhabit the islands of Slate Island Provincial Park. Photo credit: Township of Schreiber & Township of Terrace Bay www.terracebayschreiber.ca

Summary/ Description

The Little Pic regional unit is located on the northern Lake Superior shore. This regional unit extends from near the community of Schreiber in the west, to between Marathon and the Pic River in the east. This regional unit also contains the Slate Islands, which are designated as a provincial park. The Little Pic regional unit is 4,811.5 km² in size, including the associated nearshore waters. Communities in this regional unit include Marathon, Terrace Bay, Schreiber, Pays Plat First Nation and Rosspoint. The Little Pic regional unit contains one tertiary watershed, Little Pic, and eight quaternary watersheds. The watersheds in this unit are characterized by forests, primarily within provincial Crown Lands. The coast is dominated by rocky shores and cliffs and cobble beaches with scattered sandy beaches in sheltered coves. Coastal wetlands are very rare. Almost one-third of the coast is in protected areas.

TABLE 5.1: Little Pic BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
Agriculture	3.93	0.08	1,287.56	
Developed	0.16	0.00	351.99	
Forest	4,182.45	86.93	96,789.30	
Associated Nearshore Waters	257.07	5.34	15,951.22	
Other	156.36	3.25	7,452.78	
Water (inland)	211.50	4.40	8,283.49	
Total Area	4,811.5	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
Coastline (km)	406.2	NA	7.14	Based on SOLEC shoreline
Sand Beaches (km)	28.90	7.11	4.59*	*% of Lake Superior Total Sand Beaches
Coastal Wetlands (km ²)	0.92	0.27*	0.09**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
Natural Cover in Coastal Zone	315.85	93.78*	5.72**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
Number of Islands	317	NA	12.0	
Condition	Region	Region %	% of Lake Superior Total	
Population Density (persons/km ²)	0.83	NA		
Road Density (km/km ²)	0.13	NA		
Number of Dams and Barriers	305	NA	1.3	
Artificial Shoreline (km)	3.03	0.96	1.37	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
Private	117.65	2.58	4,552.56	Regional area based on landmass
Public/Crown	3,962.43	87.04	4,552.56	
Tribes/ First Nations	1.90	0.04	4,552.56	
Parks & Protected Areas (total)	477.36	10.49	4,552.56	
Parks & Protected Areas (coast)	93.07	27.63*	336.79**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- The Little Pic regional unit contains a number of sites of Important Habitat for lake trout and whitefish; these Important Habitat Sites are found in many areas along the coast (Lake Superior Binational Program Habitat Committee 2006). The lake trout Important Habitat Sites are found in many areas along the coast, as well as around the Slate Islands (Lake Superior Binational Program Habitat Committee 2006) (Figure 5.1).

Coastal Zone and Islands

- Areas identified as Important Habitat Areas are found in this regional unit, including around the Slate Islands, Neys Provincial Park and Steel River Provincial Park. Several Important Habitat Sites are also found in the White Pic region (Lake Superior Binational Program Habitat Committee 2006) (Table 5.3, Figure 5.3).
- Craig's Pit Provincial Nature Reserve is an important migratory bird observation area, and contains bluffs and kettle holes. This area also contains examples of landform processes and themes which are environmental indicators from the Lake Minong stage and later (OMNR 2006I).

Tributaries and Watersheds

- Historically 21 tributaries in Lake Superior had lake sturgeon spawning runs (Lake Superior Lake Sturgeon Work Group 2012, unpublished data). One of these historical spawning tributaries, the Prairie River, is in the Little Pic regional unit. The population status in this river is considered extirpated (Golder Associates Ltd. 2011). However a 2011 lakewide juvenile lake sturgeon survey captured sturgeon off the Prairie River. The natal source of these fish is under investigation (Lake Superior Lake Sturgeon Work Group 2012, unpublished data).

Figure 5.1: Little Pic - Coastal and Watershed Features




TABLE 5.2: Little Pic CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	NA
Nearshore ¹	C (0.52)	Unknown
Embayments and Inshore ^{1,2}	B (0.62)	Unknown
Coastal Wetlands ^{2,3}	B (0.742)	Unknown
Islands ⁴	A	Unknown
Coastal Terrestrial ³	A+ (0.996)	Unknown
Tributaries and Watersheds ²	B (0.71)	Unknown

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 5.2: Little Pic - Condition



Important Issues & Threats

- The Peninsula Harbour and Jackfish Bay Areas of Concern (AOC) are both located in the Little Pic regional unit (Figure 5.2). Seven Beneficial Use Impairments (BUIs) were identified as impaired or requiring further assessment in the Peninsula Harbour AOC and eight BUIs were identified as impaired or requiring further assessment in the Jackfish Bay AOC (Peninsula Harbour Remedial Action Plan Team 1991; Jackfish Bay Remedial Action Plan Team 1998). Mercury contamination remains a severe problem in the Peninsula Harbour AOC, with biomagnification studies and sediment analyses confirming biotic uptake of mercury from sediments (Environment Canada 2010a). The effluent discharged into Blackbird Creek in the Jackfish Bay AOC meets various legislative requirements, but the creek is considered to be contaminated along its entire length due to historical discharges (Environment Canada 2010b).
- An Environmental Assessment has been submitted for a metals and copper mine project near the north shore of Lake Superior, approximately 10 kilometres north of the community of Marathon. The Marathon Project (Marathon Platinum Group Metals-Copper Project) is a project of Stillwater Canada Inc., a subsidiary of Stillwater Mining Company.
- The Coldwell Wind Energy Project is a 66 wind turbine project in development (Brookfield Renewable Energy Partners L.P. 2013) approximately 20 kilometres northwest of Marathon, Ontario, near Neys Provincial Park and Red Sucker Point Provincial Nature Reserve (Brookfield Renewable Power Inc. No date).
- The Lower Lake Hydroelectric Project is a 10 MW hydroelectric project in development near Terrace Bay (Brookfield Renewable Energy Partners L.P. 2013).
- There is currently approximately 4,500 acres for sale on the Lake Superior coast, near Terrace Bay.

Conservation In Action

Parks & Protected Areas

- The Lake Superior National Marine Conservation Area (LSNMCA) of Canada includes a significant portion of the nearshore and inshore waters associated with the Little Pic regional unit, as well as some coastal lands (Parks Canada 2009). By law, each national marine conservation area must contain at least one zone that fosters and encourages the ecologically sustainable use of aquatic resources and at least one zone that fully protects special features or sensitive elements of ecosystems. Currently, the Lake Superior NMCA has proposed two zones that offer some protection for the Gunilda shipwreck (Jack Pine/Nipigon regional unit), and Gapen's Pool Brook Trout Spawning Area (Jack Pine/Nipigon regional unit) (C. Vis, pers. comm., May 21 2013).
- The LSMCA extends beyond the nearshore water boundary associated with the Little Pic region, to the international boundary with the United States.
- Craig's Pit Provincial Nature Reserve (also in White/ Pic)
- Red Sucker Point Provincial Nature Reserve
- Neys Provincial Park
- Slate Islands Provincial Park
- Prairie River Mouth Provincial Nature Reserve
- Steel River Provincial Park
- Rainbow Falls Provincial Park
- Schreiber Channel Provincial Nature Reserve
- Lake Superior Shoreline Enhanced Management Area (also in White and Pic and Jack Pine/Nipigon regional units)


Existing Programs & Projects

- The Slate Islands are the source of one of two lake trout brood stock gametes (Slate Island strain) used by the provincial fish culture system for lake trout rehabilitation stocking in Lake Superior (S. Greenwood, pers. comm., March 11 2013).
- The Ontario MNR Upper Great lakes Management Unit (UGLMU) has established the Fish Community Index Netting (FCIN) program on Lake Superior. Started in 2009, the FCIN program represents an ecosystem-based fish community approach, which provides trend-through-time information on the fish community. Emphasis is on the commercially important species of lake trout and lake whitefish, but the shift is away from a single species approach, to monitoring of fish population dynamics (Thunder Bay RAP 2013). The FCIN program is underway in the Thunder Bay, Peninsula Harbour and Jackfish Bay AOCs, as well as in Nipigon Bay (no commercial fisheries) (M. Chase, pers. comm., June 3 2013).
- Peregrine falcons have been reintroduced to Ontario, including in the Lake Superior basin, following their extirpation as a breeding species in Ontario in the early 1960s (Ontario Peregrine Falcon Recovery Team 2010).

TABLE 5.3: Little Pic IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
ON-008	Site	Beatty and Sturdee Coves	Colonial water bird nesting area
ON-009	Site	Big Duck Creek	Fish spawning habitat in Big Duck Creek; fish habitat
ON-014	Site	Blackbird Creek	Fish habitat (brook trout)
ON-018	Site	Nicholl Island Causeway	Spring and fall staging area for migratory birds
ON-039	Site	Golfcourse Creek	Fish spawning habitat
ON-045	Site	Hawkins Island	Colonial water bird habitat
ON-047	Site	Jackfish Lake	Former fish spawning areas
ON-067	Site	Mink Creek	Former fish spawning area
ON-087	Site	Schreiber Mine	Rare animal habitat (overwintering area for bats)
ON-109	Site	Sturdee Cove Shore	Fish spawning area
ON-125	Site	Tunnel Bay	Former fish spawning habitat
ON-137	Area	Peninsula Harbor	Former fish spawning area; colonial nesting bird habitat
ON-143	Area	Craig's Pit Nature Reserve	Environmentally Sensitive Area; hawk watch site; broken end moraine
ON-164	Area	Neys	Remnant woodland caribou population, dune and beach communities; Provincial Park
ON-170	Area	Prairie River Mouth Nature Reserve	Migratory fish habitat; sandbar, beach ridges, and dunes
ON-174	Area	Red Sucker Point Nature Reserve	
ON-177	Area	Slate Islands	Globally significant woodland caribou population, rare plant habitat
ON-179	Area	Steel River	Environmentally Sensitive Area; migratory waterfowl site; raptor and wading bird habitat; migratory fish habitat

Figure 5.3: Little Pic - Important Habitat Sites and Areas



6. Jack Pine/ Nipigon

HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	C	COASTAL WETLANDS	B
EMBAYMENTS & INSHORE	B	COASTAL TERRESTRIAL	A+
TRIBUTARIES & WATERSHEDS	B	OVERALL A-	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



Cypress Bay, in the Jack Pine/ Nipigon regional unit.
Photo credit: Ontario Ministry of Natural Resources

Summary/ Description

The Jack Pine / Nipigon regional unit is located along the northern shore of Lake Superior, and extends the furthest north of any of the regional units. The regional unit is 25,558 km² in size, including the associated nearshore waters, and is the largest of any of the regional units in this study. The regional unit extends inland considerably, and includes Lake Nipigon and the surrounding area, as well as Nipigon Bay. The relatively short shoreline of this regional unit starts just west of St. Ignace Island, and extends to east of Wilson Island, near the community of Schreiber. Communities in this regional unit include Nipigon, MacDiarmid, Beardmore, Armstrong, Whitesand First Nation, Kiashke Zaaging Anishinaabek (Gull Bay) First Nation, Biinjitiwabik Zaaging Anishinaabek (Rocky Bay) First Nation, Bingwi Neyaashi Anishinaabek (Sand Point) First Nation, Namaygoosisagagun First Nation, and Red Rock Indian Band (Lake Helen First Nation). A number of provincial parks and nature reserves are located in this region, along with an enhanced management area. Several large islands, including St. Ignace Island, Simpson Island and Wilson Island are located in this area. This regional unit contains the largest remnant population of brook trout in Lake Superior (M. Chase, pers. comm., June 3 2013). The Jack Pine/ Nipigon regional unit combines two tertiary watershed units, Jackpine and Nipigon, and contains 32 quaternary watersheds. The watersheds are almost completely dominated by forests. The coasts are characterized by rocky shores and cliffs, with scattered sand beaches and coastal wetlands.

TABLE 6.1: Jack Pine/ Nipigon BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
<i>Agriculture</i>	5.90	0.02	1,287.56	
<i>Developed</i>	1.28	0.01	351.99	
<i>Forest</i>	17,520.65	68.55	96,789.30	
<i>Associated Nearshore Waters</i>	725.96	2.84	15,951.22	
<i>Other</i>	1,774.27	6.94	7,452.78	
<i>Water (inland)</i>	5,529.76	21.64	8,283.49	
Total Area	25,557.82	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
<i>Coastline (km)</i>	522.2	NA	9.18	Based on SOLEC shoreline
<i>Sand Beaches (km)</i>	27.20	5.21	4.32*	*% of Lake Superior Total Sand Beaches
<i>Coastal Wetlands (km²)</i>	4.24	0.97*	0.43**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
<i>Natural Cover in Coastal Zone</i>	393.91	89.98*	7.13**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
<i>Number of Islands</i>	212	NA	8.0	
Condition	Region	Region %	% of Lake Superior Total	
<i>Population Density (persons/km²)</i>	0.07	NA		
<i>Road Density (km/km²)</i>	0.18	NA		
<i>Number of Dams and Barriers</i>	1188	NA	5.0	
<i>Artificial Shoreline (km)</i>	6.85	1.74	3.09	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
<i>Private</i>	521.46	2.10	24,843.18	Regional area based on landmass
<i>Public/Crown</i>	20,620.15	83.00	24,843.18	
<i>Tribes/ First Nations</i>	61.41	0.25	24,843.18	
<i>Parks & Protected Areas (total)</i>	3,662.21	14.74	24,843.18	
<i>Parks & Protected Areas (coast)</i>	56.02	12.80*	437.79**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- Nipigon Bay is noted as a Lake Superior embayment important for lake sturgeon (Auer 2003). In the Jack Pine/ Nipigon regional unit this embayment and the nearshore zone, which provides corridors for movement, are identified as critical management areas for lake sturgeon in the Lake Superior basin (Auer 2003).
- Nipigon Bay and the nearshore waters are critical management areas for brook trout in the Lake Superior basin (M. Chase, pers. comm., June 3 2013).

Coastal Zone and Islands

- The Jack Pine/ Nipigon regional unit provides several sites of Important Habitat for lake trout, including many sites around St. Ignace Island, Simpson Island, Vein Island, Wilson Island, La Grange Island and Vert Island. Additional Important Habitat sites for lake trout are found in areas along the Lake Superior coast (Lake Superior Binational Program Habitat Committee 2006) (Figure 6.1).
- One area identified as an Important Habitat Area is found around St. Ignace Island and Simpson Island, while other Important Habitat Areas are found throughout the regional unit (Lake Superior Binational Program Habitat Committee 2006) (Table 6.3, Figure 6.3).
- A number of Important Habitat Sites are clustered around the shore and some of the eastern islands (Lake Superior Binational Program Habitat Committee 2006) (Table 6.3, Figure 6.3).

Tributaries and Watersheds

- Few wild populations of brook trout in the Lake Superior basin are both sufficiently large to serve as a source population for brood stock and exhibit the migratory or lake-dwelling life history. The populations which fit these criteria include populations from Nipigon Bay and Lake Nipigon. In 2002 there were three strains of brook trout brood stock available for stocking, including the Lake Nipigon strain from Lake Nipigon, Ontario (Newman et al. 2003). The trigger that causes the development of the coaster brook trout life-history variant is unknown, and the right conditions must be present for this trait to develop. Decisions concerning the rehabilitation of brook trout must consider a number of factors (M. Chase, pers. comm., June 3 2013).
- Coaster brook trout in the Nipigon River have shown some ability to co-exist with other salmonid species, although competition may still be a factor in certain habitats and at certain densities (Newman et al. 2003).
- Historically 21 tributaries in Lake Superior had lake sturgeon spawning runs. Two of these historical spawning tributaries, the Gravel River and the Nipigon River are in the Jack Pine/ Nipigon regional unit. The Gravel River population status and population trajectory are both unknown (Golder Associates Ltd. 2011); however there is not recent evidence of natural reproduction in the Gravel River (Lake Superior Lake Sturgeon Work Group 2012, unpublished data). The Nipigon River population status is extant, while the population trajectory is unknown (Golder Associates Ltd. 2011).
- A sub-population of lake sturgeon which is physically isolated from other populations is found within Lake Nipigon. The Lake Nipigon population status is extant, while the population trajectory is stable (Golder Associates Ltd. 2011).
- A Lake Sturgeon Rehabilitation Plan for Lake Superior (Auer 2003) identifies the Gravel River and Nipigon River as two of the seventeen tributaries to Lake Superior in which there should be a focus on lake sturgeon rehabilitation.
- Several areas around Lake Nipigon are identified as Important Habitat Areas, including Kabitotikwia River Provincial Nature Reserve, West Bay Provincial Nature Reserve, Kopka River Provincial Park Addition, Livingstone Point Provincial Park and Lake Nipigon Provincial Park. A large Important Habitat area which lies mostly north of the Jack Pine/ Nipigon regional unit partially extends into the northern portion of the unit (Lake Superior Binational Program Habitat Committee 2006) (Table 6.3, Figure 6.3).

Figure 6.1: Jack Pine/ Nipigon - Coastal and Watershed Features




TABLE 6.2: Jack Pine/ Nipigon CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	NA
Nearshore ¹	C (0.59)	Unknown
Embayments and Inshore ^{1,2}	B (0.68)	Unknown
Coastal Wetlands ^{2,3}	B (0.782)	Unknown
Islands ⁴	A	Unknown
Coastal Terrestrial ³	A+ (0.996)	Unknown
Tributaries and Watersheds ²	B (0.76)	Unknown

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 6.2: Jack Pine/ Nipigon - Condition



Important Issues & Threats

- The Nipigon Bay Area of Concern (AOC) is located in the Jack Pine/ Nipigon regional unit (Figure 6.2). Eight Beneficial Use Impairments (BUIs) were identified as impaired or requiring further assessment in the 1991 Stage 1 Status. As of the 2010 Beneficial Use Impairments Status Report the number of BUIs which remain classified as impaired or requiring further assessment has decreased to four (Nipigon Bay Remedial Action Plan Team 2011).

Conservation In Action

Parks & Protected Areas

- The Lake Superior National Marine Conservation Area (LSNMCA) of Canada includes a significant portion of the nearshore and inshore waters associated with the Jack Pine/ Nipigon regional unit, as well as some coastal lands (Parks Canada 2009). By law, each national marine conservation area must contain at least one zone that fosters and encourages the ecologically sustainable use of aquatic resources and at least one zone that fully protects special features or sensitive elements of ecosystems. Currently, the Lake Superior NMCA has proposed two zones that offer some protection for the Gunilda shipwreck (Jack Pine/Nipigon regional unit), and Gapen's Pool Brook Trout Spawning Area (Jack Pine/Nipigon regional unit) (C. Vis, pers. comm., May 21 2013).
- The LSNMCA extends beyond the nearshore water boundary associated with the Little Pic region, to the international boundary with the United States.
- Gravel River Provincial Nature Reserve
- Ruby Lake Provincial Park
- Lake Nipigon Provincial Park
- Livingstone Point Provincial Park
- Wabakimi Provincial Park (portion)
- Whitesand Provincial Park
- Windigo Bay Provincial Park
- Kopka River Provincial Park
- Kopka River Provincial Park Addition
- Wilson Island
- West Bay Provincial Nature Reserve
- Pantagrue Creek Provincial Nature Reserve
- Gull River Provincial Park
- Kabitotkwia River Provincial Nature Reserve
- Kaiashk Provincial Nature Reserve
- Black Sturgeon River Provincial Park (portion)
- Lake Superior Shoreline Enhanced Management Area (also in White and Pic and Little Pic regional units)

Existing Programs & Projects

- Existing populations of native brook trout in the Nipigon area seem to be responding well to measures taken for their conservation, including restrictions on harvests and the protection of spawning habitat (Newman et al. 2003). Seasonal refuge areas have been established on the Nipigon River by the Ontario Ministry of Natural Resources, in order to protect spawning adults and habitat from destruction by fishermen wading through the area (Newman et al. 2003).


Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

- Both the Ontario Ministry of Natural Resources Dorion Fish Culture Station and the Red Cliff Tribal Fish Hatchery rear captive brood stock and production fish of the Lake Nipigon strain of brook trout (Newman et al. 2003). The OMNR Dorion Fish Culture Station is located in the Black-Sturgeon region, while the Red Cliff Tribal Fish Hatchery is located in the Beartrap-Nemadji region. The offspring from the Dorion Fish Culture Station are not stocked into Lake Superior (M. Chase, pers. comm., June 3 2013).
- The Ontario MNR Upper Great Lakes Management Unit (UGLMU) has established the Fish Community Index Netting (FCIN) program on Lake Superior. Started in 2009, the FCIN program represents an ecosystem-based fish community approach, which provides trend-through-time information on the fish community. Emphasis is on the commercially important species of lake trout and lake whitefish, but the shift is away from a single species approach, to monitoring of fish population dynamics (Thunder Bay RAP 2013). The FCIN program is underway in the Thunder Bay, Peninsula Harbour and Jackfish Bay AOCs, as well as in Nipigon Bay (no commercial fisheries) (M. Chase, pers. comm., June 3 2013).
- Peregrine falcons have been reintroduced to Ontario, including in the Lake Superior basin, following their extirpation as a breeding species in Ontario in the early 1960s (Ontario Peregrine Falcon Recovery Team 2010).
- Wilson Island is owned by the Nature Conservancy of Canada, and managed through an agreement with the Pays Plat First Nations.

TABLE 6.3: Jack Pine/ Nipigon IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
ON-017	Site	Cat Islands	Nesting site for colonial water birds
ON-022	Site	Cobinosh Island	Historic rare animal habitat
ON-023	Site	Cypress River	Fish spawning area
ON-030	Site	Dublin Creek	Suspected fish spawning area
ON-046	Site	Inside Islands of Nipigon Bay	Fish habitat; raptor habitat
ON-048	Site	Kama Bay West	Fish spawning area
ON-051	Site	Lake Helen	Former fish spawning area
ON-058	Site	McInnes Lake and Creek	Fish spawning area
ON-074	Area	Gravel River Nature Reserve	Important staging area for migration
ON-077	Site	Kabitotikuia River Mouth	Coastal wetland, rare animal habitat; Provincial Nature Reserve
ON-078	Site	Nipigon River and Bay	Significant coastal wetland; waterfowl nesting/staging area; high biodiversity value
ON-078	Area	Nipigon River and Bay	Significant coastal wetland; waterfowl nesting/staging area; high biodiversity value
ON-085	Site	Northeast Wilson Island	Rare plant habitat
ON-088	Site	Onaman River Mouth	Large wetland; high biodiversity value
ON-091	Site	Outan Island	Rare animal habitat
ON-093	Site	Parmachene Bridge at Polly Lake	
ON-119	Site	Speckle Islands	Colonial water bird habitat
ON-121	Area	St. Ignace and Simpson Islands	Rare animal habitat (woodland caribou unoccupied)
ON-123	Site	Steamboat Bay	Former fish spawning area
ON-133	Site	Windikokan Lake	Fish spawning area
ON-149	Area	Kabitotikuia River Nature Reserve	
ON-152	Area	Kama Hill Nature Reserve	Kama Cliffs; representative landform and vegetation types (sparse forests and conifer on broken bedrock)
ON-156	Area	Lake Nipigon	Woodland caribou calving islands, long undeveloped lakeshore
ON-157	Area	Windigo Bay Nature Reserve	Woodland caribou habitat and migration route, sand dune communities
ON-161	Area	Livingstone Point Nature Reserve	
ON-182	Area	Wabakimi Wilderness Area	Woodland caribou habitat, large pristine protected area
ON-183	Area	West Bay Nature Reserve	Fish spawning habitat

Figure 6.3: Jack Pine/ Nipigon - Important Habitat Sites and Areas



7. Black Sturgeon

HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	C	COASTAL WETLANDS	A-
EMBAYMENTS & INSHORE	B	COASTAL TERRESTRIAL	A+
TRIBUTARIES & WATERSHEDS	A	OVERALL	A-

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



The Black Sturgeon River, looking towards the river mouth. Photo credit: Ontario Ministry of Natural Resources

Summary/ Description

The Black Sturgeon regional unit is located on the northern shore of Lake Superior. This regional unit is 6,333 km² in size, including the associated nearshore waters. The Black Sturgeon regional unit extends from near Thunder Bay in the west, to near the communities of Red Rock and Nipigon in the east. Communities in this regional unit include Red Rock, Red Rock Indian Band (Lake Helen First Nation), the Township of Shuniah, and the Township of Dorion. This regional unit includes Black Bay, a portion of Thunder Bay, and the Sibley and Black Bay Peninsulas. The A number of provincial parks, nature reserves and conservation areas are located in this regional unit. The Lakehead Region Conservation Authority's area of jurisdiction covers eight municipalities, including municipalities in the Black Sturgeon and Dog/Arrow regional units. The Black Sturgeon regional unit contains one tertiary watershed, Black Sturgeon, and 12 quaternary watersheds. The watersheds are characterized by forests. The coast is characterized by rocky shores, cliffs, cobble beaches, and includes some of the most extensive coastal wetlands on the Ontario portion of Lake Superior.

TABLE 7.1: Black Sturgeon BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
Agriculture	13.24	0.21	1,287.56	
Developed	0.96	0.02	351.99	
Forest	4,431.78	69.97	96,789.30	
Associated Nearshore Waters	1,407.41	22.22	15,951.22	
Other	256.68	4.05	7,452.78	
Water (inland)	223.40	3.53	8,283.49	
Total Area	6,333.46	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
Coastline (km)	840.8	NA	14.78	Based on SOLEC shoreline
Sand Beaches (km)	46.70	5.55	7.42*	*% of Lake Superior Total Sand Beaches
Coastal Wetlands (km ²)	46.16	6.57*	4.69**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
Natural Cover in Coastal Zone	641.21	91.32*	11.61**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
Number of Islands	426	NA	16.1	
Condition	Region	Region %	% of Lake Superior Total	
Population Density (persons/km ²)	0.74	NA		
Road Density (km/km ²)	0.32	NA		
Number of Dams and Barriers	504	NA	2.1	
Artificial Shoreline (km)	8.17	1.27	3.68	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
Private	883.99	17.85	4,953.68	Regional area based on landmass
Public/Crown	3,648.67	73.66	4,953.68	
Tribes/ First Nations	0.00	0.00	4,953.68	
Parks & Protected Areas (total)	472.10	9.53	4,953.68	
Parks & Protected Areas (coast)	135.87	19.35*	702.19**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- Black Bay is noted as a Lake Superior embayment important for lake sturgeon (Auer 2003). In the Black Sturgeon regional unit this embayment and the nearshore zone, which provides corridors for movement, are identified as critical management areas for lake sturgeon in the Lake Superior basin (Auer 2003).
- Cisco are an important feature in western Lake Superior. Large spawning aggregations of cisco are found in Black Bay and Thunder Bay (S. Greenwood, pers. comm., March 11 2013).

Coastal Zone and Islands

- The Black Sturgeon regional unit provides several sites of Important Habitat for lake trout, including many sites around the southern tip of the Black Bay Peninsula, Edward Island and Porphyry Island, and the southern tip of the Sibley Peninsula. One Important Habitat area for lake whitefish is noted in Black Bay. Additional Important Habitat sites for lake trout are found in areas along the Lake Superior coast in Thunder Bay (Lake Superior Binational Program Habitat Committee 2006) (Figure 7.1).
- Several Important Habitat Areas are located in the Black Sturgeon regional unit, including Black Bay, the Black Bay Peninsula, much of the Sibley Peninsula, and several smaller Important Habitat Areas on Edward Island and Porphyry Island. Additional Important Habitat Areas are found in other areas of the regional unit, along with several Important Habitat Sites (Lake Superior Binational Program Habitat Committee 2006) (Table 7.3, Figure 7.3).

Tributaries and Watersheds

- The Black Sturgeon River contains spawning habitat for native migratory fishes, but the Camp 43 dam restricts fish movement beyond the dam. The Ontario Ministry of Natural Resources (OMNR) is currently undertaking an Environmental Assessment to determine if they will decommission the Camp 43 dam (OMNR 2012). (See Important Issues and Threats).
- Historically 21 tributaries in Lake Superior had lake sturgeon spawning runs (Lake Superior Lake Sturgeon Work Group 2012, unpublished data). Two of these historical spawning tributaries, the Black Sturgeon River and the Wolf River are in the Black Sturgeon regional unit. The Black Sturgeon River population status is extant and the population trajectory is unknown. The Wolf River population status is extirpated (Golder Associates Ltd. 2011).
- A Lake Sturgeon Rehabilitation Plan for Lake Superior (Auer 2003) identifies the Black Sturgeon River and Wolf River as two of the seventeen tributaries to Lake Superior in which there should be a focus on lake sturgeon rehabilitation. Habitat restoration is a priority in the Black Sturgeon River and Wolf River, due to barriers to migration and spawning (Auer 2003).

Figure 7.1: Black Sturgeon - Coastal and Watershed Features




TABLE 7.2: Black Sturgeon CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	NA
Nearshore ¹	C (0.59)	Unknown
Embayments and Inshore ^{1,2}	B (0.72)	Unknown
Coastal Wetlands ^{2,3}	A- (0.808)	Unknown
Islands ⁴	A	Unknown
Coastal Terrestrial ³	A+ (0.994)	Unknown
Tributaries and Watersheds ²	A (0.84)	Unknown

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 7.2: Black Sturgeon - Condition



Important Issues & Threats

- The Ontario Ministry of Natural Resources (OMNR) is currently undertaking an Environmental Assessment process to examine the recommendations of the Fishery Management Zone 9 Advisory Council: to decommission the Camp 43 dam and construct a multi-purpose sea lamprey barrier at Eskwanonwatin Lake (OMNR 2012). One of the largest populations of walleye in Lake Superior previously inhabited Black Bay. The Camp 43 dam, 17 kilometres upstream on the Black Sturgeon River, cut off access to spawning habitat for native migratory fishes. The removal of the dam and the reinstatement of access to naturally occurring spawning habitat would be an essential step in the large scale rehabilitation of the Black Bay and Black Sturgeon River native fish community (OMNR 2012). However, the removal of the dam requires the construction of a new sea lamprey barrier further upstream. If the proposed project is completed an additional 50 kilometres of river habitat will be available to native migratory fish species, including walleye, lake sturgeon and coaster brook trout (OMNR 2012). Additional possible benefits stemming from the decommissioning of the dam would provide brook trout with access to 40 small tributaries, along with an un-quantified amount of spawning and nursery habitat; the potential reestablishment of walleye and cisco as the top predator and prey species in Black Bay, with the reduction of the rainbow smelt population; and avoiding catastrophic failure of the dam before it is at the end of its lifespan. Potential effects and mitigation measures are also described in the project summary (OMNR 2012).
- The preferred option for the decommissioning of the Camp 43 dam also identifies the construction of a new barrier to sea lamprey, 50 kilometres upstream. The construction of this new barrier is described as a way to open up the spawning area in the Black Sturgeon River, while still minimizing the production of new sea lamprey in the river (OMNR 2012). This preferred option will result in the opening of 50 kilometres of the Black Sturgeon River to sea lamprey, and is stated to result in “an incremental increase in parasitic sea lamprey in Lake Superior” (OMNR 2012:1). However some experts note that the cost of additional lampricide treatment is high relative to the Lake Superior control budget for sea lamprey, and that there are additional risks to lake trout and other large fish species by allowing sea lamprey additional river access (S Greenwood, pers. comm., March 11 2012).

Conservation In Action

Parks & Protected Areas

- The Lake Superior National Marine Conservation Area (LSNMCA) of Canada includes all of the nearshore and inshore waters in the Black Sturgeon regional unit, as well as some coastal lands (Parks Canada 2009). By law, each national marine conservation area must contain at least one zone that fosters and encourages the ecologically sustainable use of aquatic resources and at least one zone that fully protects special features or sensitive elements of ecosystems. Currently, the Lake Superior NMCA has proposed two zones that offer some protection for the Gunilda shipwreck (Jack Pine/Nipigon regional unit), and Gapen’s Pool Brook Trout Spawning Area (Jack Pine/Nipigon regional unit) (C. Vis, pers. comm., May 21 2013).
- The LSMCA extends beyond the nearshore water boundary associated with the Little Pic region, to the international boundary with the United States.
- Black Sturgeon River Provincial Park (portion)
- Albert Lake Mesa Provincial Nature Reserve
- Cavern Lake Provincial Nature Reserve
- Ouimet Canyon Provincial Park
- Hurkett Cove Conservation Area
- Granite Point Conservation Area

- Shesheeb Bay Provincial Nature Reserve
- Sleeping Giant Provincial Park
- Edward Island Provincial Park
- Porphyry Island Provincial Park
- Puff Island Provincial Nature Reserve
- Silver Harbour Conservation Area


Existing Programs & Projects

- The Ontario Ministry of Natural Resources Dorion Fish Culture Station is the site of captive rearing of brood stock and production fish for the Lake Nipigon strain of brook trout (Newman et al. 2003). This facility is located in the Black-Sturgeon region of the Lake Superior basin. The offspring from the Dorion Fish Culture Station are not stocked into Lake Superior (M. Chase, pers. comm., June 3 2013).
- The Ontario Ministry of Natural Resources undertook tracking of adult lake sturgeon on the Black Sturgeon River, to monitor seasonal distribution and identify important habitat (OMNR 2013c). This work ended in the fall of 2009 (M. Chase, pers. comm., June 3 2013).
- The Lakehead Region Conservation Authority (LRCA) undertakes a number of programs and services related to the Black Sturgeon regional unit. These include water control structures such as dams and channels, forest management and tree planting, and erosion control, among other programs (LRCA No date).
- Peregrine falcons have been reintroduced to Ontario, including in the Lake Superior basin, following their extirpation as a breeding species in Ontario in the early 1960s (Ontario Peregrine Falcon Recovery Team 2010).

TABLE 7.3: Black Sturgeon IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
ON-011	Site	Black Bay Peninsula	High biodiversity value, relatively undisturbed habitat, significant wetland
ON-011	Area	Black Bay Peninsula	High biodiversity value, relatively undisturbed habitat, significant wetland
ON-013	Site	Black Sturgeon River, Split Rapids	Spawning area and previous spawning area for fish
ON-020	Site	Clark Island	Historic rare animal habitat
ON-021	Site	Nipigon Bay Clay Banks	Historic fish spawning habitat
ON-034	Site	Fluor Island	High biodiversity values
ON-083	Site	Nonwatin Lake and Black Sturgeon River	Fish spawning area
ON-106	Site	Shangoina Island	Rare animal habitat (unoccupied by peregrine falcon), colonial water bird habitat
ON-107	Site	Shillibeer Lake	Wetland; staging area for waterfowl
ON-138	Area	Black Bay	Extensive coastal wetlands, diversity of aquatic habitats; provincially significant bog community at east end of bay
ON-140	Area	Albert Lake Mesa Nature Reserve	Rare plant habitat
ON-142	Area	Cavern Lake Nature Reserve	Arctic-alpine plant community, rare plants, bat hibernaculum
ON-146	Area	Edward Island Nature Reserve	Fish spawning habitat; unique landform vegetation; part of Lake Superior Archipelago
ON-166	Area	Ouimet Canyon Nature Reserve	Arctic-alpine plant community, rare plant habitat
ON-171	Area	Puff Island Nature Reserve	Environmentally Sensitive Area (unusual geological features, bog area)
ON-175	Area	Shesheeb Bay Nature Reserve	Fish spawning habitat
ON-178	Area	Sleeping Giant	Old growth red and white pine stands, rare plant habitat, cliff habitats, coastal wetlands

Figure 7.3: Black Sturgeon - Important Habitat Sites and Areas



8. Dog/Arrow

HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	D	COASTAL WETLANDS	C
EMBAYMENTS & INSHORE	C	COASTAL TERRESTRIAL	A
TRIBUTARIES & WATERSHEDS	C	OVERALL B-	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



The mouth of the Current River, near Thunder Bay, Ontario. Photo credit: Ontario Ministry of Natural Resources

Summary/ Description

The Dog/ Arrow regional unit is located on the northwestern shore of Lake Superior, and is 10,111.61 km² in size, including the associated nearshore waters. This regional unit extends from the international boundary in the west to just past the community of Thunder Bay in the east. Communities in this regional unit include the City of Thunder Bay, Fort William First Nation, Lac Des Mille Lacs First Nation, the Municipality of Oliver-Paipoonge, the Municipality of Neebing, the Township of O'Connor, the Township of Conmee, and the Township of Gillies. Several provincial parks, nature reserves and conservation areas are located in this regional unit. The Lakehead Region Conservation Authority's area of jurisdiction covers eight municipalities, including municipalities in the Dog/Arrow and Black Sturgeon regional units. The Dog/ Arrow regional unit combines two tertiary watersheds, Dog and Arrow, and 17 quaternary watersheds. The watersheds are characterized by forests. The coast is dominated by cobble beaches with rocky shores and scattered coastal wetlands.

TABLE 8.1: Dog/ Arrow BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
Agriculture	70.13	0.69	1,287.56	
Developed	40.39	0.40	351.99	
Forest	8,485.26	83.92	96,789.30	
Associated Nearshore Waters	437.89	4.33	15,951.22	
Other	474.62	4.69	7,452.78	
Water (inland)	603.32	5.97	8,283.49	
Total Area	10,111.61	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
Coastline (km)	302.1	NA	5.31	Based on SOLEC shoreline
Sand Beaches (km)	1.20	0.40	0.19*	*% of Lake Superior Total Sand Beaches
Coastal Wetlands (km ²)	15.52	7.34*	1.58**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
Natural Cover in Coastal Zone	173.94	82.30*	3.15**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
Number of Islands	109	NA	4.1	
Condition	Region	Region %	% of Lake Superior Total	
Population Density (persons/km ²)	6.40	NA		
Road Density (km/km ²)	0.84	NA		
Number of Dams and Barriers	2942	NA	12.4	
Artificial Shoreline (km)	36.83	21.17	16.59	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
Private	2,798.91	28.92	9,676.59	Regional area based on landmass
Public/Crown	6,567.32	67.87	9,676.59	
Tribes/ First Nations	56.95	0.59	9,676.59	
Parks & Protected Areas (total)	254.02	2.63	9,676.59	
Parks & Protected Areas (coast)	15.69	7.42*	211.35**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- Thunder Bay is noted as a Lake Superior embayment important for lake sturgeon (Auer 2003). In the Dog/ Arrow regional unit this embayment and the nearshore zone, which provides corridors for movement, are identified as critical management areas for lake sturgeon in the Lake Superior basin (Auer 2003).
- Cisco are an important feature in western Lake Superior. Large spawning aggregations of cisco are found in Thunder Bay (S. Greenwood, pers. comm., March 11 2013).

Coastal Zone and Islands

- The Dog/ Arrow regional unit provides several sites of Important Habitat for lake trout, including many in Thunder Bay and around Pie Island and Flatland Island. One location of Important Habitat for lake whitefish is noted along the shore, near the international border (Lake Superior Binational Program Habitat Committee 2006) (Figure 8.1).

Tributaries and Watersheds

- Historically 21 tributaries in Lake Superior had lake sturgeon spawning runs. Two of these historical spawning tributaries, the Kaministiquia River and the Pigeon River are in the Dog/Arrow regional unit (the Pigeon River is on the shared boundary between the Dog/Arrow and Baptism-Brule regional units). The Kaministiquia River population status is extant and the population trajectory is stable. The Pigeon River population status is extant, while the population trajectory is unknown (Golder Associates Ltd. 2011).
- A Lake Sturgeon Rehabilitation Plan for Lake Superior (Auer 2003) identifies the Kaministiquia and Pigeon Rivers as two of the seventeen tributaries to Lake Superior in which there should be a focus on lake sturgeon rehabilitation.
- Several Important Habitat Areas are located throughout the Dog/Arrow regional unit, including Castle Creek Provincial Nature Reserve and Fraleigh Lake Provincial Nature Reserve and Kashabowie Provincial Park. Important Habitat Sites are also located throughout the Dog/Arrow regional unit (Lake Superior Binational Program Habitat Committee 2006) (Table 8.3, Figure 8.3).

Figure 8.1: Dog/ Arrow - Coastal and Watershed Features




TABLE 8.2: Dog/ Arrow CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	NA
Nearshore ¹	D (0.25)	Unknown
Embayments and Inshore ^{1,2}	C (0.40)	Unknown
Coastal Wetlands ^{2,3}	C (0.578)	Unknown
Islands ⁴	A	Unknown
Coastal Terrestrial ³	A (0.945)	Unknown
Tributaries and Watersheds ²	C (0.54)	Unknown

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 8.2: Dog/ Arrow - Condition



Important Issues & Threats

- The Thunder Bay Area of Concern (AOC) is located in the Dog/Arrow regional unit (Figure 8.2). The boundaries run along 28 km of shoreline and 9 km off shore, and the Welcome Islands are included in the boundaries. The issues associated with this AOC include Impairments to Fish Health, Impairments to Sediment Quality, Impairments to Water Quality and Impairments to Wildlife. Approximately \$165.6 million (Canadian dollars) has been spent to complete a number of remedial projects. The Public Advisory Committee (PAC) was re-established in 2009, following a six year hiatus after the Stage 2 report. The PAC is organized into five subcommittees to focus on the remaining beneficial use impairments (BUIs) (Thunder Bay RAP 2013).

Conservation In Action

Parks & Protected Areas

- Centennial Park
- Cascades Conservation Area
- Wishart Conservation Area
- Hazelwood Lake Conservation Area
- Silver Falls Provincial Park
- Kashabowie Provincial Park
- Little Greenwater Lake Provincial Nature Reserve
- Matawin River Provincial Nature Reserve
- Arrow Lake Provincial Park
- Castle Creek Provincial Nature Reserve
- Divide Ridge Provincial Nature Reserve
- Fraleigh Lake Provincial Nature Reserve
- Kakabeka Falls Provincial Park
- Cedar Falls Conservation Area
- Le Pate Provincial Nature Reserve
- Thomson Island Provincial Nature Reserve
- Pigeon River Provincial Park
- Devon Road Mesa Provincial Nature Reserve
- Le Verendrye Provincial Park

Existing Programs & Projects

- The Ontario Ministry of Natural Resources (MNR) conducted an eight-year study to track adult lake sturgeon and monitor larval lake sturgeon in the Kaministiquia River below Kakabeka Falls (Thunder Bay RAP 2013). The fieldwork was completed in 2012 (M. Chase, pers. comm., June 3 2013). The purpose was to determine the impacts of water flow and water level variation on lake sturgeon spawning success. This research will help to protect spawning habitat and rehabilitate lake sturgeon populations which use the Kaministiquia River, as well as to guide future hydro-electric development and water management planning around Lake Superior. This work is being completed in partnership with Ontario Power Generation (OMNR 2013c).
- The Ontario MNR Upper Great lakes Management Unit (UGLMU) has established the Fish Community Index Netting (FCIN) program on Lake Superior. Started in 2009, the FCIN program represents an ecosystem-based fish community approach, which provides trend-through-time information on the fish community. Emphasis is on the commercially important species of lake trout


and lake whitefish, but the shift is away from a single species approach, to monitoring of fish population dynamics (Thunder Bay RAP 2013). The FCIN program is underway in the Thunder Bay, Peninsula Harbour and Jackfish Bay AOCs, as well as in Nipigon Bay (no commercial fisheries) (M. Chase, pers. comm., June 3 2013).

- The Kaministiquia River Walleye Radio Telemetry Project is a project which uses radio telemetry to record the movement patterns and seasonal distribution of adult walleye in the Kaministiquia River. The project will help to identify critical habitat and migration routes, as well as to monitor improvements made to alleviate water quality barriers. The multi-year project started in 2009 (Thunder Bay RAP 2013).
- The Current River Walleye Assessment involves the periodic assessment of the spawning population of walleye at the mouth of the Current River, within Thunder Bay harbour. Artificial spawning habitat was created in 1991 to replace historical spawning habitat lost due to construction and development. A mark-recapture protocol has been used during three phases (1991-1993, 1999-2000 and 2010-2012) to conduct spawning assessments (Thunder Bay RAP 2013).
- The Lakehead Region Conservation Authority (LRCA) undertakes a number of programs and services related to the Dog/Arrow regional unit. These include water control structures such as dams and channels, forest management and tree planting, and erosion control, among other programs (LRCA No date).
- Peregrine falcons have been reintroduced to Ontario, including in the Lake Superior basin, following their extirpation as a breeding species in Ontario in the early 1960s (Ontario Peregrine Falcon Recovery Team 2010).
- The Thunder Bay Field Naturists have an active land securement program and has recently purchased important coastal wetlands at Pine Bay.

TABLE 8.3: Dog/ Arrow IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
ON-016	Site	Caldwell Lake	Wetland; rare plant and animal habitat
ON-049	Site	Keifer Terminal Floodway	Coastal wetland; waterfowl staging area and brood rearing
ON-057	Site	Matawin River Nature Reserve	System of productive wetlands; fish spawning habitat waterfowl staging area, rare plant habitat
ON-057	Area	Matawin River Nature Reserve	System of productive wetlands; fish spawning habitat waterfowl staging area, rare plant habitat
ON-059	Area	Cloud Bay	Coastal wetland, waterfowl migration habitat
ON-086	Site	Northwood Bog	Rare plant habitat
ON-098	Site	Prince and Jarvis	Undisturbed, diverse habitat; raised cobble beaches and Norwester Chain Mountains
ON-114	Site	Slate River Drainage	Waterfowl staging and brood habitat
ON-116	Site	South Fowl Lake	Rare plant habitat (provincially and locally rare plants); Rare animal habitat, excellent waterfowl habitat
ON-129	Site	West Whitefish Lake	Large inland wetland, waterfowl breeding and staging area, rare animal habitat
ON-141	Area	Castle Creek Nature Reserve	Cliff habitat, wetlands, rare plants
ON-144	Area	Devon Road Mesa Nature Reserve	Rare plant habitat
ON-145	Area	Divide Ridge Nature Reserve	Cliff habitat
ON-147	Area	Fraleigh Lake Nature Reserve	
ON-151	Area	Kakabeka Falls	Fish spawning habitat
ON-153	Area	Kashabowie	Fish spawning habitat
ON-155	Area	La Verendrye	Rare plant habitat, cliff communities, wild rice marshes
ON-158	Area	Le Pate Nature Reserve	Unique landform feature (mesa)
ON-160	Area	Little Greenwater Lake Nature Reserve	
ON-163	Area	Middle Falls	Fish spawning habitat
ON-181	Area	Thompson Island Nature Reserve	

Figure 8.3: Dog/ Arrow - Important Habitat Sites and Areas




9. Baptism-Brule

HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	C	COASTAL WETLANDS	B
EMBAYMENTS & INSHORE	C	COASTAL TERRESTRIAL	A+
TRIBUTARIES & WATERSHEDS	C	OVERALL B	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



Susie Island is the largest of 13 small, rocky islands jutting out of Lake Superior at the Pigeon River outlet. The island has been protected by The Nature Conservancy. Photo credit: The Nature Conservancy.

Summary/ Description

The Baptism-Brule region is located in the western portion of the Lake Superior basin, from the Ontario-Minnesota international boundary to just north of Silver Bay (near Illgen City), Minnesota. Including the nearshore waters associated with this regional unit, it is 3,912 km² in size. This hydrologic region is referred to as HUC 04010101 and is part of the larger Subregion 0401, Western Lake Superior. The region is located within the Northern Lakes and Forest ecoregion of Minnesota (USDA NRCS No date a), and is also referred to as the Lake Superior North Watershed by the Minnesota Pollution Control Agency (Minnesota PCA 2012a). Most of the land-base of the regional unit is in Cook County, with a smaller portion in Lake County (USDA NRCS No date a). The largest land ownership type in the watershed is federal ownership. State ownership is the second largest ownership type, followed by private. The remaining land is owned by tribal, private major, county or conservancy agencies (USDA NRCS No date a). Communities in the area include: Finland, Schroeder, Tofte, Lutsen, Grand Marais, Hovland (Minnesota PCA 2012a). The Grand Portage Band of Lake Superior Chippewa community is located at Grand Portage, and is one of the oldest Ojibwa settlements in Minnesota (Grand Portage No date). The Sawtooth Mountains are found along the shoreline (USDA NRCS No date a, No date b). The Baptism-Brule regional unit contains one tertiary (HUC 8) watershed, Baptism-Brule, and 11 quaternary (HUC 10) watersheds. The watersheds are almost completely forested. The coasts are dominated by exposed rocky shores and cliffs. Coastal wetlands are very rare in this region.

TABLE 9.1: Baptism-Brule BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
Agriculture	0.49	0.01	1,287.56	
Developed	0.72	0.02	351.99	
Forest	3,454.99	88.31	96,789.30	
Associated Nearshore Waters	242.06	6.19	15,951.22	
Other	68.23	1.74	7,452.78	
Water (inland)	145.95	3.73	8,283.49	
Total Area	3,912.44	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
Coastline (km)	211.5	NA	3.72	Based on SOLEC shoreline
Sand Beaches (km)	2.00	0.95	0.32*	*% of Lake Superior Total Sand Beaches
Coastal Wetlands (km ²)	11.23	3.80*	1.14**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
Natural Cover in Coastal Zone	280.75	94.94*	5.08**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
Number of Islands	46	NA	1.7	
Condition	Region	Region %	% of Lake Superior Total	
Population Density (persons/km ²)	1.41	NA		
Road Density (km/km ²)	0.38	NA		
Number of Dams and Barriers	816	NA	3.5	
Artificial Shoreline (km)	3.98	1.42	1.79	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
Private	107.7	2.93	3,680.89	Regional area based on landmass
Public/Crown	3,267.96	88.78	3,680.89	
Tribes/ First Nations	171.51	4.66	3,680.89	
Parks & Protected Areas (total)	136.23	3.7	3,680.89	
Parks & Protected Areas (coast)	57.58	19.47*	295.72**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- Grand Portage Bay, Clark's Bay and Wausaugoning Bay are noted as Lake Superior embayments which are important for lake sturgeon (Auer 2003). In the Baptism-Brule regional unit these embayments and the nearshore zone, which provides corridors for movement, are identified as critical management areas for lake sturgeon in the Lake Superior basin (Auer 2003). Pigeon Bay is also noted to be used by lake sturgeon, in fact it is the bay most used by lake sturgeon in Grand Portage waters (S. Moore, pers. comm., May 14 2013).
- Areas identified as important habitat for lake trout are found in areas along the coast, and areas identified as important habitat for lake whitefish are found near Grand Portage, at the northern end

of the Baptism-Brule regional unit (Lake Superior Binational Program Habitat Committee 2006) (Figure 9.1).

- The Baptism-Brule regional unit has several areas identified as biologically important by the Lake Superior Binational Program Habitat Committee (2006). A number of Important Habitat Sites and Important Habitat Areas are located within the region and along the shoreline (Table 9.3, Figure 9.3).

Coastal Zone and Islands

- A few small State Important Bird Areas are found along the coast in the Baptism-Brule regional unit (National Audubon Society 2013, 2012). These sites are some of the nine locations along Lake Superior that make up the 125 acre North Shore Peregrine Falcon Eyries IBA. These nine cliff areas are geographically separate and under different ownerships, but combined represent 70% of the recorded natural nest sites for Peregrine falcons in Minnesota (Minnesota DNR 2013b).
- Susie Island is a Minnesota Biological Survey Site of Statewide Biodiversity Significance. Susie Island is noted to be ecologically significant due to unique flora and the presence of arctic-alpine disjunct species. Nine rare plant species are known to occur on the island (Minnesota DNR 1984).
- Arctic disjunct plant species occur in several locations along the coast (B. Carlson, pers. comm., March 20 2013)

Tributaries and Watersheds

- Historically 21 tributaries in Lake Superior had lake sturgeon spawning runs. One of these historical spawning tributaries, the Pigeon River is on the shared boundary between the Dog/Arrow and Baptism-Brule regional units. The Pigeon River population status is extant, while the population trajectory is unknown (Golder Associates Ltd. 2011).
- A Lake Sturgeon Rehabilitation Plan for Lake Superior (Auer 2003) identifies the Pigeon River as one of the seventeen tributaries to Lake Superior in which there should be a focus on lake sturgeon rehabilitation.
- Despite impairment issues in some areas, many areas of the watershed are described as of exceptional water quality (Minnesota PCA 2012a).
- The Art Lake Hardwood Ridges area is a Minnesota Biological Survey Site of Outstanding Biodiversity Significance. It is noted as a large natural area (4,670 acres) which is unfragmented in nature. The site contains high-quality native plant communities, including communities ranked as S2 (Imperiled) and S3 (Vulnerable to Extirpation) by the Natural Heritage and Nongame Wildlife Research Program. Large patches of old-growth upland forests and lowland forests are found within this area, as are rare plants and a rare bird species (defined as rare by Minnesota statutes) (Minnesota DNR 2008).
- The Baptism-Brule watershed is highly forested; other land covers include open water, shrub or scrubland and wetlands (USDA NRCS No date a). Agriculture accounts for a very small portion of land use; much of the land in this region is not well-suited for agriculture (USDA NRCS No date a).

Figure 9.1: Baptism-Brule - Coastal and Watershed Features




TABLE 9.2: Baptism-Brule CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	
Nearshore ¹	C (0.57)	
Embayments and Inshore ^{1,2}	C (0.57)	
Coastal Wetlands ^{2,3}	B (0.706)	
Islands ⁴	A	
Coastal Terrestrial ³	A+ (0.979)	Some local experts feel a grade of A may accurately reflect local conditions in the Coastal Terrestrial target. This is due to the combined effects of recent housing development fragmenting the forest, and the forest lacking much of its natural conifer component. Significant changes to forest cover and forest disturbance over the past 100 years have impacted the Coastal Terrestrial target (E. Perry, pers. comm., February 26 2013).
Tributaries and Watersheds ²	C (0.57)	

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 9.2: Baptism-Brule - Condition



Important Issues & Threats

- The Rapid Watershed Assessment completed by the United States Department of Agriculture Natural Resources Conservation Service (USDA NRCS No date a) identifies several watershed concerns in the Baptism-Brule region. These include erosion (sheet and rill, streambank, lakeshore and roadside), groundwater and surfacewater quality and quantity, and management of timberlands, shoreline and wetlands (USDA NRCS No date a). The transport of sediments and pollutants to surfacewater due to erosion and stormwater are also identified (USDA NRCS No date a).
- Development pressures are stated to be moderate in this region (USDA NRCS No date a), however, development along the Lake Superior shoreline is noted as significant (Minnesota PCA 2012a). Other areas of the Baptism-Brule region are noted to be facing increased growth and development pressures, including the along the shorelines of the lower reaches of the Poplar and Flute Reed rivers (Minnesota PCA 2012a). This development is noted to be a contributing factor to pollution problems (Minnesota PCA 2012a).
- Estimates indicate 22 farm operations are located in the watershed region; more than 80% of these farms are less than 180 acres in size (USDA NRCS No date a).
- Some streams and lakes in the Baptism-Brule region are classified as impaired due to identified impairments, such as mercury or PCB in fish tissue, mercury in the water column, or turbidity. In the affected waterbodies, these impairments lead to designated uses being affected (Minnesota PCA 2012a).
- The USGS lists a total of 14 records for Nonindigenous Aquatic Species in the Baptism-Brule region. Of these, 4 are classified as exotic, 9 as native, and 1 as native hybrid (USGS 2012a).
- An Emergency Prevention and Response Plan for Viral Hemorrhagic Septicemia has been developed for Isle Royale National Park, Pictured Rocks National Lakeshore, Apostle Islands National Lakeshore and the Grand Portage Band of the Lake Superior Chippewa Reservation (within which is the Grand Portage National Monument) (NPS 2013a).
- Forest fragmentation as a result of housing development is an emerging concern in Minnesota. The forest that is present lacks much of its natural conifer component (E. Perry, pers. comm., February 26 2013).

Conservation In Action

Parks & Protected Areas

- Temperance State Park
- Cascade State Park
- Judge CR Magney State Park
- Boundary Waters Canoe Area Wilderness (within Superior National Forest)
- Superior National Forest

Existing Programs & Projects

- Brook trout restocking efforts in Grand Portage, Minnesota, using fertilized eggs or fry of the Nipigon-strain of brook trout have been successful. The stocked brook trout migrated to Lake Superior and as adults they returned to the streams where they were stocked and successfully reproduced (Newman et al. 2003). The success of this restocking effort may be due to a combination of factors, including the strain of brook trout used, the early life stage at which the brook trout were stocked, and the protection from overharvest provided by Grand Portage (Newman et al. 2003).

- There are a number of Minnesota Biological Survey (MBS) Sites delineated in the Baptism-Brule regional units within Minnesota, some of which have been ranked with Outstanding or High Biodiversity Significance, based on statewide ranking criteria. The Minnesota Department of Natural Resource's MBS systematically collects, interprets, and delivers data on the distribution and ecology of native plants, animals, native plant communities, and functional landscapes throughout the state. MBS conducts landscape assessments, field surveys and monitoring activities, and provides data and tools to guide conservation and management within *MBS Sites of Statewide Biodiversity Significance* (MBS Sites). Biodiversity information includes the location and biodiversity significance rank of MBS Sites, the location and status of rare species populations, the type and condition of native plant communities, and, for selected sites, *MBS Ecological Evaluation* reports (Minnesota DNR 2013e, B. Carlson, pers. comm., March 20 2013). The MBS Sites located within the Baptism-Brule regional unit are Art Lake Hardwood Ridges, Deronda Bay, George Crosby Manitou State Park & Caribou Falls State Wayside, Horseshow Bay Shore, Hovland lookout Tower, Hovland Woods, Hovland Woods SNA (Swamp River W), Icelandite Coastal Fen, Iona's Beach, Lake Agnes Northern Hardwoods, Lutsen Natural Area, Myhr Creek Ridge, Ninemile Lakes and Ridges, Susie Island and Thomsonite Beach (L. Gerdes, pers. comm., March 18 2013).
- The State of Minnesota specifies a policy goal of non-degradation for all waters, maintaining them in a natural and unpolluted state. There are three levels of protection for surface waters. The highest level of protection applies to Outstanding Resource Value Waters (ORVWs). Additionally, all surface waters in the Lake Superior basin are Outstanding International Resource Waters (OIRW) (MPCA 2012e).
- The Grand Portage Band of Lake Superior Chippewa uses the designation of Outstanding Tribal Water Resources (OTRWs) as part of an anti-degradation policy to maintain and protect high quality waters. All waters within the boundaries of the Grand Portage Reservation are OTWRs, assigned to one of two subcategories. Each subcategory has specific implementation procedures (Grand Portage Band of Lake Superior Chippewa 2006).
- The Natural Resources Conservation Service (NRCS) Performance Results System (PRS) provides support for reporting the development and delivery of conservation programs (USDA NRCS No date d). From 1999 through 2007 a total of 46,953 acres were planned for conservation use, through the Total Conservation Systems. During this same period, a total of 3,945 acres of the planned conservation systems were applied (USDA NRCS No date a). Some of the conservation practices implemented included tree and shrub establishment (amounting to 1,032 total acres), total wildlife habitat (977 total acres), total wetlands created, restored or enhanced (60 total acres) and erosion control total soil saved (amounting to 722 tons per year) (USDA NRCS No date a).
- A number of projects, plans and monitoring programs are underway in this region. The 10 year rotation for intensive watershed monitoring for Minnesota's major watersheds will take place in the Baptism-Brule region in 2013; further studies and plans may be developed depending on the results of the monitoring program (Minnesota PCA 2012a). Monitoring of the Flute Reed River is undertaken by a partnership including a citizen's organization, the Minnesota Pollution Control Agency and the county Soil and Water Conservation District (Minnesota PCA 2012a). Lake associations are also monitoring lakes and working to develop lake management plans (Minnesota PCA 2012a).
- The Manitou Collaborative is a partnership which includes the United States Forest Service, the Minnesota Forest Resources Council, The Nature Conservancy, the Minnesota Department of Natural Resources, Wolf Ridge Environmental Learning Center and Lake County. The partnership of

public and private landowners began in 2000, and collaboratively the partners manage 100,000 acres in northeastern Minnesota. One fifth of the Manitou Landscape area is classified as Outstanding for statewide biodiversity significance, and 200 miles of high quality streams are located within this area. Mutually agreed upon management objectives for the vegetation include mimicking the range of natural variability to restore diverse and multi-aged forests and promoting diverse forests of multiple growth stages, while supporting the local economy (The Manitou Collaborative No date, USDA Forest Service No date a). The Art Lake Hardwood Ridges Minnesota Biological Survey Site of Outstanding Biodiversity Significance is located within the Manitou Collaborative area of focus (Minnesota DNR 2008).

- The North Shore Forest Collaborative is a combined effort of local, state and federal groups, along with public and private groups and individuals. Concentrated on the ecosystems along the North Shore of Lake Superior, the Collaborative agencies work together to restore and maintain native trees and forest communities for a healthy forest environment (North Shore Forest Collaborative No date).
- The North Shore Stewardship Association works to promote the protection and restoration of the North Shore of Lake Superior (Sugarloaf: The North Shore Stewardship Association No date).
- Six Citizen-based Groups are noted to do work in the Baptism-Brule (U.S. EPA 2013b). Additional projects, plans, conservation districts, organizations and partners related to the Baptism-Brule regional unit are noted in the Rapid Watershed Assessment (USDA NRCS No date a).
- Minnesota Biological Survey (MBS) Sites of Biodiversity Significance ranked High and Outstanding (B. Carlson, pers. comm., March 20 2013)


TABLE 9.3: Baptism-Brule IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
MN-004	Site	Amenda Creek	Northern Hardwood Forest, Upland White Cedar Forest
MN-006	Site	Grand Marais Point	Arctic disjunct plant community, rare plant habitat, geologic features
MN-011	Site	Big Bay	Geologic Feature
MN-013	Area	Boundary Waters Canoe Area	Rare plant and animal habitat, large representative ecosystems, geologic features
MN-015	Area	Butterwort Cliffs SNA	Rock shore community, aspen-birch forest, rare plant habitat, colonial waterbird habitat
MN-017	Site	Cannonball Bay	Arctic disjunct plant community, rare plant habitat
MN-018	Site	Caribou Falls WMA	Anadromous fish habitat, deer concentration area
MN-019	Area	Cascade River State Park	Arctic disjunct plant community, rare plant habitat
MN-026	Site	Deronda Bay and Red Rock	Rare plant habitat, geologic feature
MN-028	Site	Devil Track Lake	Rare animal habitat
MN-031	Site	Five Mile Rock	Colonial waterbird nesting habitat, geologic feature
MN-034	Area	George H. Crosby Manitou State Park	Northern hardwood forest, upland white cedar forest, rare animal habitat, rare plant habitat, anadromous fish habitat
MN-035	Site	Good Harbor Bay	Arctic disjunct plant community, rare plant habitat, geologic feature
MN-037	Site	Grand Marais Fen	Poor fen, sedge subtype
MN-038	Site	Grand Portage 4	Rare plant community
MN-039	Area	Grand Portage State Park	Rare plant habitat
MN-040	Area	Hat Point Area	Representative forest ecosystems, coastal shore communities, important natural/cultural resource
MN-041	Site	Heartbreak Creek	Northern hardwood-conifer forest, yellow birch-white cedar subtype, upland white cedar forest
MN-042	Site	Hollow Rock	Geomorphic feature (sea arch)
MN-045	Site	Horseshoe Bay	Geomorphic feature (raised beach)
MN-046	Area	Hovland Woods SNA	Large old growth forest complex with bogs, swamps, lake in the landscape, rare plant and animal habitat
MN-050	Area	Judge C. R. Magney State Park	Old growth white pine forest, rare plant habitat, geomorphic features, anadromous fish habitat
MN-051	Site	Kadunce Creek	Rare plant community
MN-052	Site	Kennedy Creek	Rare animal habitat
MN-055	Site	Lake Agnes Hardwoods	Northern hardwood forest, rare plant habitat
MN-056	Area	Lake Superior Highlands	Extensive natural communities and high biodiversity, rare plant and animal habitat
MN-058	Site	LeVeaux WMA	Representative natural plant communities, rare animal habitat
MN-060	Site	Little Marais	Rare animal habitat, colonial waterbird nesting habitat
MN-062	Area	Lutsen SNA	Old growth Northern Hardwood forest and upland white cedar forest
MN-065	Site	Manitou River	Fish spawning habitat, rare plant habitat
MN-068	Site	Mineral Center Maple Ridge	Northern Hardwood forest, rare plant habitat
MN-071	Site	Moose Fence Cedars	Upland white cedar forest
MN-077	Site	Oberg Mountain Hardwoods	Northern hardwood forest, rare plant habitat
MN-078	Site	Onion River Hardwoods	Northern hardwood forest, rare plant habitat
MN-079	Area	Paradise Beach	Coastal wetland, tamarack swamp, colonial waterbird nesting habitat, waterbird concentrations, geologic formations

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

Code	Site/ Area	Important Habitat Site/Area Name	Key Features
MN-080	Site	Pearl Beach Hardwoods	Northern hardwood forest
MN-082	Site	Poplar River	Fish spawning habitat
MN-088	Site	Schroeder RNA	Northern hardwood forest, rare plant habitat
MN-090	Site	South Fowl Lake	Rare plant habitat
MN-091	Site	South Lutsen	Rare plant and animal habitat
MN-093	Area	Spring Beauty Hardwoods SNA	Old growth northern hardwood forest, rare plant habitat
MN-095	Area	Sugar Loaf Point SNA	Coastal wetland restoration project, rare plant habitat, geologic features
MN-097	Area	Susie Islands	Arctic-disjunct plant community, rare animal habitat
MN-098	Area	Swamp River Bog	Rare plant communities, old growth forest, rare plant habitat, rare animal habitat, waterbird concentrations
MN-099	Area	Temperance River State Park	Rare plant habitat, arctic disjunct plant populations, unusual geomorphic feature
MN-100	Area	Tettegouche State Park	Lake Superior pebble and bedrock beaches, exposed cliffs, Northern Hardwood-Conifer Forest, Northern Oak Forest, Upland White Cedar Forest
MN-102	Site	Tofte Town Park	Arctic and alpine disjunct plant habitat
MN-105	Site	Wolf Ridge	Rare animal habitat
MN-106	Site	Wringer Lake Hardwoods	Northern hardwood forest, rare plant habitat
MN-107	Site	Yellow Birch	Northern hardwood forest, rare plant habitat
MN-109	Site	Cross River State Park	Northern hardwoods forest, rare plant habitat
MN-110	Area	Devils Track Falls State Park	Rare plant habitat
MN-112	Area	Kadunce River State Park	Rare plant habitat
ON-155	Area	La Verendrye	Rare plant habitat, cliff communities, wild rice marshes

Figure 9.3: Baptism-Brule - Important Habitat Sites and Areas



10. Isle Royale

HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	B	COASTAL WETLANDS	A-
EMBAYMENTS & INSHORE	B	COASTAL TERRESTRIAL	A+
TRIBUTARIES & WATERSHEDS	A	OVERALL A	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



Summary/ Description

Isle Royale is located in the northwestern part of Lake Superior, the closest mainland is approximately 14 miles (22.5 kilometres) away (NPS 2013c). The island of Isle Royale is a national park, along with the more than 450 smaller islands located around it (NPS 2013c). This regional unit is 950.37 km² in size, including the associated nearshore waters. The waters of Isle Royale are said to contain the most productive native fishery in Lake Superior, as well as the most genetically diverse lake trout populations in the entire lake (NPS 2013c). Only about half of the mammal species found on the mainland are found in the park, and for some of these species long-term research has been studying ecological interactions (NPS 2013c, UNESCO 2005). A number of other research projects are carried out in the park, due to its remote and relatively undisturbed ecosystem; research topics include vegetation studies and acid rain (UNESCO 2005). Several western disjunct plant species are found in the Isle Royale regional unit (NPS 2013c). The Isle Royale regional unit is composed solely of islands and island complexes. There are no tertiary (HUC 8) or quaternary (HUC 10) watersheds identified for this regional unit. The watersheds are completely forested, and are some of the most intact in the Lake Superior basin. The coasts are dominated by exposed rocky shores and cliffs. Wetlands are common within the coastal area.

TABLE 10.1: Isle Royale BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
<i>Agriculture</i>	0.0	0.0	1,287.56	
<i>Developed</i>	0.08	0.01	351.99	
<i>Forest</i>	451.82	47.54	96,789.30	
<i>Associated Nearshore Waters</i>	446.70	47.00	15,951.22	
<i>Other</i>	26.06	2.74	7,452.78	
<i>Water (inland)</i>	25.71	2.71	8,283.49	
Total Area	950.37	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
<i>Coastline (km)</i>	470.0	NA	8.26	Based on SOLEC shoreline
<i>Sand Beaches (km)</i>	0.00	0.00	0.00*	*% of Lake Superior Total Sand Beaches
<i>Coastal Wetlands (km²)</i>	94.05	28.92*	9.55**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
<i>Natural Cover in Coastal Zone</i>	290.82	89.42*	5.27**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
<i>Number of Islands</i>	433	NA	16.3	
Condition	Region	Region %	% of Lake Superior Total	
<i>Population Density (persons/km²)</i>	0.00	NA		
<i>Road Density (km/km²)</i>	0	NA		
<i>Number of Dams and Barriers</i>	0	NA	0	
<i>Artificial Shoreline (km)</i>	0.89	0.31	0.40	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
<i>Private</i>	2.87	0.56	512.55	Regional area based on landmass
<i>Public/Crown</i>	0.00	0.00	512.55	
<i>Tribes/ First Nations</i>	0.00	0.00	512.55	
<i>Parks & Protected Areas (total)</i>	511.72	99.84	512.55	
<i>Parks & Protected Areas (coast)</i>	322.36	99.12*	325.23**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- Important habitat for both lake whitefish and lake trout is found in many areas of Isle Royale (Lake Superior Binational Program Habitat Committee 2006) (Figure 10.1).
- The fisheries around Isle Royale may contain unique morphotypes of lake trout (NPS 2013a).
- Sampling of the nearshore and inshore fish community over approximately 300 kilometres of Isle Royale shoreline in and around major embayments resulted in 17 fish species sampled. Twelve of these species were common in the Lake Superior fish community (Gorman et al. 2008).

Coastal Zone and Islands

- The coastal zone of Isle Royal is one of the most intact in the entire Great Lakes
- The entire Isle Royale regional unit is identified as an Important Habitat Area by the Lake Superior Binational Program Habitat Committee (2006). Several Important Habitat Sites are also located on the island (Table 10.3, Figure 10.3).
- Isle Royale National Park is noted as a State Important Bird Area (National Audubon Society 2013, 2012).
- The islands of Isle Royale National Park are home to 18 species of mammal, in comparison to the 40 or more mammal species found on the mainland (NPS 2013c). The isolation of the island makes colonisation by new species more difficult and some species which were previously present (e.g. caribou and coyote) are no longer part of the Isle Royale ecosystem (NPS 2013c).
- The shoreline of Isle Royale is described as heavily forested (NPS 2013c).

Tributaries and Watersheds

- Several of the streams support coaster brook trout
- As with the coastal areas, the watersheds of the island are some of the last undisturbed watersheds in the Great Lakes. Many types of wetlands are supported on Isle Royale (NPS 2013c).
- Some inland lakes support high abundances of clams, while other inland lakes contain no clams at all (NPS 2013c).

Figure 10.1: Isle Royale - Coastal and Watershed Features




TABLE 10.2: Isle Royale CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	
Nearshore ¹	B (0.61)	
Embayments and Inshore ^{1,2}	B (0.76)	
Coastal Wetlands ^{2,3}	A- (0.837)	
Islands ⁴	A	
Coastal Terrestrial ³	A+ (1.000)	
Tributaries and Watersheds ²	A (0.90)	

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 10.2: Isle Royale - Condition



Important Issues & Threats

- An Emergency Prevention and Response Plan for Viral Hemorrhagic Septicemia has been developed for Isle Royale National Park, Pictured Rocks National Lakeshore, Apostle Islands National Lakeshore and the Grand Portage Band of the Lake Superior Chippewa Reservation (within which is the Grand Portage National Monument) (NPS 2013a).
- The presence of Emerald Ash Borer (EAB) has not yet been detected on Isle Royale; however, the island is now under Federal EAB quarantine and Michigan state quarantine (Cooperative Emerald Ash Borer Project 2013).
- Some harvesting of white pine for timber has occurred in the past (UNESCO 2005).
- Acid rain has been studied in Isle Royale (UNESCO 2005).
- High levels of PCBs are noted in the lakes of Isle Royale (UNESCO 2005).
- Invasive gypsy moths (*Lymantria dispar*) have been trapped on Isle Royale since 2000, but no reproducing populations are known to exist on the island (NPS 2013c).
- Invasive spiny water fleas (*Bythotrephes cederstroemi*) have been found in the Lake Superior waters of Isle Royale, but they have not yet been discovered in the inland lakes of Isle Royale (NPS 2013c).

Conservation In Action

Parks & Protected Areas

- Isle Royale National Park
- Isle Royale Biosphere Reserve

Existing Programs & Projects

- Few wild populations of brook trout are both sufficiently large to serve as a source population for brood stock, and exhibit the migratory or lake-dwelling life history. In 2002 three strains in the Lake Superior basin met both of these criteria. Two of these strains were from the Isle Royale area, including the Tobin Harbor and Siskiwit Bay strains (Newman et al. 2003). The Siskiwit Bay strain comes from a migratory population which spawns in the Big Siskiwit River and the Little Siskiwit River (Newman et al. 2003). The gametes that contribute to the captive stock were collected in 1995 and 1999 (Newman et al. 2003).
- Captive brood stock and production fish from the Tobin Harbor and Siskiwit Bay strains of brook trout are reared at the U.S. Fish and Wildlife Service Iron River National Fish Hatchery (in the Beartrap-Nemadji region) (Newman et al. 2003).
- Under the Michigan Water Quality Standards (WQS), all waterbodies within the designated boundary of Isle Royale National Park have been designated as Outstanding State Resource Waters. Under the above designation, additional anti-degradation controls are applied for new or increased pollutant loadings (Michigan DEQ 2013).
- The ongoing ecological study of wolves on Isle Royale has been underway for 55 years (Vucetich and Peterson 2013); it is the longest continuous predator/prey study in the world (Vucetich 2012, Vucetich and Peterson 2013). The predator/prey interaction study between wolves and moose is part of the ongoing research in Isle Royale National Park (UNESCO 2005). The wolf population in January 2013 was 8 individuals, the lowest the population has ever been during the history of the study (Vucetich and Peterson 2013). This is also the first year in the history of the project that no reproduction was documented (Vucetich and Peterson 2013).
- A 13 year breeding bird survey conducted on Isle Royale revealed significant increases in abundance for ten species, and significant decreases in abundance for eight species; these trends were


consistent with studies from other areas (Egan 2009). 85 species were detected over the 13 years, with an average of 57 species detected each year (Egan 2009).

- The Isle Royale & Keweenaw Parks Association is a non-profit educational organization that works with the National Park Service to promote public understanding of Isle Royale National Park and Keweenaw National Historical Park. The group also works to fund research projects (Isle Royale & Keweenaw Parks Association 2010).

TABLE 10.3: Isle Royale IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
MI-008	Site	Caribou Island	Rare plant habitat
MI-011	Site	Davidson Island	Rare plant and animal habitat
MI-012	Site	Edwards Island	Rare plant habitat
MI-015	Site	Hat Island	Rare plant and animal habitats
MI-043	Area	Isle Royale	Rare plant and animal habitat, fish spawning habitat, colonial waterbird habitats

Figure 10.3: Isle Royale - Important Habitat Sites and Areas




11. Beaver-Lester


HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	D	COASTAL WETLANDS	C
EMBAYMENTS & INSHORE	D	COASTAL TERRESTRIAL	A
TRIBUTARIES & WATERSHEDS	C	OVERALL C	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



Cliffs and rocky shores at Palisade Head and Shovel Point, Tettegouche State Park. Photo credit: [wikipedia.org/wiki/Portal: Minnesota/ Scenic](https://wikipedia.org/wiki/Portal:Minnesota/Scenic)

Summary/ Description

The Beaver-Lester region extends from just north of Silver Bay, Minnesota, to Duluth, Minnesota. The Beaver-Lester regional unit is 1,686.67km² in size, including the associated nearshore waters. This regional unit spans two Minnesota counties, with approximately 35% of the land-base of the regional unit in St. Louis County and approximately 65% of the regional unit land-base in Lake County (USDA NRCS No date b). The Beaver-Lester region is located in the Northern Lakes and Forest ecoregion, and the watershed is described as largely forested (USDA NRCS No date b). This hydrologic region is referred to as HUC 04010102. This region is part of Subregion 0401 - Western Lake Superior. Long ridges slope towards Lake Superior, and the Sawtooth Mountains are found along the shoreline (USDA NRCS No date a, No date b). Communities in the area include Silver Bay, Two Harbors, Arnold, Knife River and Beaver Bay. The Beaver- Lester regional unit contains one tertiary (HUC 8) watershed, Beaver-Lester, and four quaternary (HUC 10) watersheds. Most of the region is forested. The coast is characterized by rocky shores and cliffs. Coastal wetlands are very rare, and concentrated in the Knife River area.

TABLE 11.1: Beaver-Lester BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
<i>Agriculture</i>	6.97	0.41	1,287.56	
<i>Developed</i>	21.21	1.26	351.99	
<i>Forest</i>	1,354.35	80.30	96,789.30	
<i>Associated Nearshore Waters</i>	249.88	14.81	15,951.22	
<i>Other</i>	49.06	2.91	7,452.78	
<i>Water (inland)</i>	5.20	0.31	8,283.49	
Total Area	1,686.67	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
<i>Coastline (km)</i>	114.5	NA	2.01	Based on SOLEC shoreline
<i>Sand Beaches (km)</i>	0.90	0.79	0.14*	*% of Lake Superior Total Sand Beaches
<i>Coastal Wetlands (km²)</i>	6.53	3.57*	0.66**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
<i>Natural Cover in Coastal Zone</i>	160.25	87.46*	2.90**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
<i>Number of Islands</i>	12	NA	0.5	
Condition	Region	Region %	% of Lake Superior Total	
<i>Population Density (persons/km²)</i>	44.24	NA		
<i>Road Density (km/km²)</i>	0.91	NA		
<i>Number of Dams and Barriers</i>	933	NA	3.9	
<i>Artificial Shoreline (km)</i>	10.81	6.75	4.87	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
<i>Private</i>	1,158.35	80.29	1,442.78	Regional area based on landmass
<i>Public/Crown</i>	244.05	16.92	1,442.78	
<i>Tribes/ First Nations</i>	0.00	0.00	1,442.78	
<i>Parks & Protected Areas (total)</i>	40.57	2.81	1,442.78	
<i>Parks & Protected Areas (coast)</i>	20.92	11.42*	183.22**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- Important habitat areas for lake trout are found in several locations along the coast, and one important habitat area for lake whitefish is noted between Duluth and Two Harbors (Lake Superior Binational Program Habitat Committee 2006) (Figure 11.1).

Coastal Zone and Islands

- The Beaver-Lester regional unit contains a number of areas identified as biologically important by the Lake Superior Binational Program Habitat Committee (2006). A number of Important Habitat Sites are located along the Lake Superior shore, as well as in some inland areas. Much of the coastal area is an Important Habitat Area, as is Tettegouche State Park, a portion of which lies in both the Beaver-Lester and the Baptism-Brule regional units (Table 11.3, Figure 11.3).

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

- Several small State Important Bird Areas (IBAs) are found along the coast in the Beaver-Lester regional unit (National Audubon Society 2013, 2012). These sites are some of the nine locations along Lake Superior that make up the 125 acre North Shore Peregrine Falcon Eyries IBA. These nine cliff areas are geographically separate and under different ownerships, but combined represent 70% of the recorded natural nest sites for Peregrine falcons in Minnesota (Minnesota DNR 2013b). The Hawk Ridge Nature Reserve IBA is another State IBA located in the Beaver-Lester regional unit (National Audubon Society 2013, 2012). This IBA is noted as one of the best locations in North America for autumn hawk watching (Minnesota DNR 2013c).
- Arctic disjunct plant species, native plant communities occur along the coast (B. Carlson, pers. comm., March 20 2013).
- Minnesota Point is a freshwater sandbar and associated rare species, rare native plant communities (B. Carlson, pers. comm., March 20 2013).
- Knife Island is an important island for colonial nesting waterbirds (T. Kaspar, pers. comm., March 14 2013)

Tributaries and Watersheds

- Although the watershed are characterized for forests, this region is highly fragmented by roads.

Figure 11.1: Beaver-Lester - Coastal and Watershed Features




TABLE 11.2: Beaver-Lester CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	
Nearshore ¹	D (0.20)	
Embayments and Inshore ^{1,2}	D (0.31)	
Coastal Wetlands ^{2,3}	C (0.519)	
Islands ⁴	A	
Coastal Terrestrial ³	A (0.938)	Local experts feel a condition grade of B (Good) may accurately reflect local conditions in the Coastal Terrestrial target. The rationale for this condition is that the forest of this area, though extensive, is heavily dominated by relatively young forest of aspen and birch (with notable exceptions) and missing conifers and older age classes. It has also been fragmented by development near the shore (E. Perry, pers. comm., February 26 2013).
Tributaries and Watersheds ²	C (0.42)	

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 11.2: Beaver-Lester - Condition



Important Issues & Threats

- The ports of Two Harbors, Minnesota, Superior, Wisconsin-Duluth, Minnesota and Presque Isle-Marquette, Michigan have been identified as sites at high risk for invasion by aquatic invasive species, due to ballast water from laker traffic (Rup et al. 2010 as cited in International Joint Commission (IJC) Work Group on Aquatic Invasive Species Rapid Response 2011).
- The St. Louis River Area of Concern (AOC) is located in three regional units, including portions of the Beaver-Lester regional unit. The St. Louis River has faced issues of habitat loss and degradation, and pollution and contamination, especially in its lower reaches. The lower 39 miles (63 kilometres) of the St. Louis River are the main focus of the St. Louis River Remedial Action Plan (RAP) (U.S. EPA 2013c) and are designated as a Lake Superior Area of Concern (AOC). Nine beneficial use impairments were identified in the St. Louis River AOC (U.S. EPA 2013c).
- Thirty-three farms are noted to operate in this regional unit. Approximately two-thirds of those farms are less than 180 acres, while one-third of the farms are 180 to 1,000 acres (USDA NRCS No date b). Land within this regional unit is generally considered as poorly suited for agriculture (USDA NRCS No date b).
- Erosion, water quality and quantity, and management of shorelines, woodlands, stormwater and wetlands are noted in the Rapid Watershed Assessment as the main resource concerns in this regional unit (USDA NRCS No date b).
- Development pressure is noted to be moderate in this region (USDA NRCS No date b). The lakeshore and timberlands are two areas noted to be parceled out for new recreation, lake or country homes (USDA NRCS No date b).
- A number of waterbodies within the Beaver-Lester regional unit are listed as impaired. Reasons for impairment include mercury, pH, turbidity, low dissolved oxygen, absence of trout due to temperature and PCBs. Affected uses include aquatic consumption and aquatic life (USDA NRCS No date b).
- Over 50 species were listed as endangered or threatened (USDA NRCS No date b).
- The USGS lists a total of 13 records for Nonindigenous Aquatic Species in the Beaver-Lester region; 4 are classified as exotic, 8 as native, and 1 as native hybrid (USGS 2012b).
- Forest fragmentation as a result of housing development is an emerging concern in Minnesota. The forest that is present lacks much of its natural conifer component (E. Perry, pers. comm., February 26 2013).

Conservation In Action

Parks & Protected Areas

- Tettegouche State Park

Existing Programs & Projects

- There are a number of Minnesota Biological Survey (MBS) Sites delineated in the Beaver-Lester regional unit, some of which have been ranked with Outstanding or High Biodiversity Significance, based on statewide ranking criteria. The Minnesota Department of Natural Resource's MBS systematically collects, interprets, and delivers data on the distribution and ecology of native plants, animals, native plant communities, and functional landscapes throughout the state. MBS conducts landscape assessments, field surveys and monitoring activities, and provides data and tools to guide conservation and management within *MBS Sites of Statewide Biodiversity Significance* (MBS Sites). Biodiversity information includes the location and biodiversity significance rank of MBS Sites, the location and status of rare


species populations, the type and condition of native plant communities, and, for selected sites, *MBS Ecological Evaluation* reports (Minnesota DNR 2013e, B. Carlson, pers. comm., March 20 2013). The MBS Sites located within the Beaver-Lester regional unit are Encampment Forest Area, Fault Line Ridges, Magney-Snively, Minnesota Point Pine Forest, Moose Mountain, Nopeming Unconformity and Tettegouche State Park (L. Gerdes, pers. comm., March 18 2013).

- The State of Minnesota specifies a policy goal of nondegradation for all waters, maintaining them in a natural and unpolluted state. There are three levels of protection for surface waters. The highest level of protection applies to Outstanding Resource Value Waters (ORVWs). Additionally, all surface waters in the Lake Superior basin are Outstanding International Resource Waters (OIRW) (MPCA 2012e).
- The Manitou Collaborative is a partnership which includes the United States Forest Service, the Minnesota Forest Resources Council, The Nature Conservancy, the Minnesota Department of Natural Resources, Wolf Ridge Environmental Learning Center and Lake County. The partnership of public and private landowners began in 2000, and collaboratively the partners manage 100,000 acres in northeastern Minnesota. One fifth of the Manitou Landscape area is classified as Outstanding for statewide biodiversity significance, and 200 miles of high quality streams are located within this area. Mutually agreed upon management objectives for the vegetation include mimicking the range of natural variability to restore diverse and multi-aged forests and promoting diverse forests of multiple growth stages, while supporting the local economy (The Manitou Collaborative No date, USDA Forest Service No date a).
- The North Shore Forest Collaborative is a combined effort of local, state and federal groups, along with public and private groups and individuals. Concentrated on the ecosystems along the North Shore of Lake Superior, the Collaborative agencies work together to restore and maintain native trees and forest communities for a healthy forest environment (North Shore Forest Collaborative No date).
- Sugarloaf: The North Shore Stewardship Association works to promote the protection and restoration of the North Shore of Lake Superior (Sugarloaf: The North Shore Stewardship Association No date).
- 12 Citizen-based Groups are noted to do work in the Beaver-Lester regional unit (U.S. EPA 2013d). Additional projects, plans, conservation districts, organizations and partners related to the Beaver-Lester regional unit are noted in the Rapid Watershed Assessment (USDA NRCS No date b).
- The Natural Resources Conservation Service (NRCS) Performance Results System (PRS) provides support for reporting the development and delivery of conservation programs (USDA NRCS No date d). From 1999 to 2007 plans were made for a total of 1,454 acres of Total Conservation Systems. From 1999 to 2007 the Total Conservation Systems Applied amounted to 656 acres. The activities which contributed the largest amount to the Total Conservation Systems Applied were Total Wildlife Habitat (350 acres), Erosion Control Total Soil Saved (99 tons/year), Tree and Shrub Establishment (250 acres) and Riparian Forest Buffers (48 acres). Additional activities involved prescribed grazing and wetlands (created, restored or enhanced) (USDA NRCS No date b).
- Shoreline Management of the North Shore Management Zone is described as local districts regulating and managing development density on and adjacent to shorelines, while giving priority to environmental protection and orderly growth (USDA NRCS No date b).

TABLE 11.3: Beaver-Lester IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
MN-001	Site	Agate Bay	Colonial waterbird nesting, waterfowl concentrations
MN-007	Site	Beaver Bay	Waterbird concentrations, Upland White Cedar Forest
MN-008	Site	Beaver Island	Arctic disjunct plant community, rare plant habitat, colonial waterbird nesting habitat
MN-014	Site	Burlington Bay	Colonial Waterbirds, waterfowl concentrations
MN-020	Site	Cathedral Grove	Great Lakes pine forest, old growth forest
MN-025	Site	Crow Creek Bluff	Rare animal habitat
MN-029	Site	Duluth 8	Rare animal habitat
MN-030	Site	Flood Bay	Colonial waterbird nesting habitat, shorebird migratory habitat, geomorphic feature
MN-036	Area	Gooseberry Falls State Park	Conifer, aspen and birch forests, fish spawning habitat, rare plant and animal habitat
MN-053	Site	Knife River	Rare plant habitat, anadromous fish habitat, geomorphic features
MN-054	Site	Lafayette Bluff and Encampment Island	Colonial waterbird nesting habitat
MN-056	Area	Lake Superior Highlands	Extensive natural communities and high biodiversity, rare plant and animal habitat
MN-057	Site	Lester Park Waterworks	Rare plant habitat
MN-059	Site	Lighthouse Point	Rare plant habitat
MN-067	Site	Marble Lake Lookout Tower	Northern Hardwoods forest, rare plant habitat
MN-072	Area	Moose Mountain SNA	Old growth Northern Hardwoods forest, rare plant habitat
MN-076	Site	Normanna 18	Rare animal habitat
MN-086j	Area	St. Louis Estuary	Great Lakes freshwater estuary, rare plant and animal habitat, colonial waterbird nesting habitat
MN-092	Area	Split Rock Lighthouse State Park	Rare plant and animal habitats, colonial waterbird nesting habitat, geomorphic feature, waterbird concentrations
MN-094	Site	Stony Point	Rare plant habitat, arctic disjunct plant community, waterbird concentrations
MN-100	Area	Tettegouche State Park	Lake Superior pebble and bedrock beaches, exposed cliffs, Northern Hardwood-Conifer Forest, Northern Oak Forest, Upland White Cedar Forest
MN-110	Area	Devils Track Falls State Park	Rare plant habitat

Figure 11.3: Beaver-Lester - Important Habitat Sites and Areas




12. St. Louis/ Cloquet

HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	D	COASTAL WETLANDS	D
EMBAYMENTS & INSHORE	D	COASTAL TERRESTRIAL	C
TRIBUTARIES & WATERSHEDS	D	OVERALL C	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



Clough Island is located in the St. Louis River Estuary – one of the largest estuaries in the Great Lakes. Clough Island was protected by The Nature Conservancy, the Wisconsin Department of Natural Resources (WDNR) and the U.S. Fish and Wildlife Service in 2010. As of 2011 the WDNR owns and manages Clough Island.

Summary/ Description

The St. Louis/ Cloquet region includes the westernmost portion of the Lake Superior watershed. The St. Louis/ Cloquet regional unit is 8,677 km² in size, including the associated nearshore waters. This region encompasses only a small portion of the Lake Superior shoreline, the Duluth-Superior harbour from Duluth, Minnesota to Superior, Wisconsin. However, the region extends inland to the north and to the west, into Minnesota, and includes the communities of Hoyt Lakes, Virginia, Hibbing, Floodwood and Cloquet. The Fond du Lac Indian Reservation is located in this regional unit and the Fond du Lac Band of Lake Superior Chippewa is a significant landholder in the regional unit.

The St. Louis River is the largest U.S. tributary and the second largest tributary to Lake Superior (Minnesota PCA 2012c). The St. Louis River is a very biologically productive area and becomes an estuary in its lower reaches. The Cloquet and Whiteface rivers are the main tributaries. This regional unit contains both highly developed areas, as well as high quality areas. The St. Louis/ Cloquet regional unit combines two tertiary (HUC 8) watersheds, St. Louis and Cloquet, and contains 22 quaternary (HUC 10) watersheds. The watersheds of this region are dominated by forests, but the coastal areas have the highest amount of developed land and artificial shoreline in Lake Superior.

TABLE 12.1: St. Louis/ Cloquet BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
<i>Agriculture</i>	160.64	1.85	1,287.56	
<i>Developed</i>	214.44	2.47	351.99	
<i>Forest</i>	7,179.87	82.75	96,789.30	
<i>Associated Nearshore Waters</i>	71.06	0.82	15,951.22	
<i>Other</i>	881.22	10.16	7,452.78	
<i>Water (inland)</i>	169.67	1.96	8,283.49	
Total Area	8,676.90	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
<i>Coastline (km)</i>	85.9	NA	1.51	Based on SOLEC shoreline
<i>Sand Beaches (km)</i>	8.59	10.00	1.36*	*% of Lake Superior Total Sand Beaches
<i>Coastal Wetlands (km²)</i>	27.58	20.21*	2.80**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
<i>Natural Cover in Coastal Zone</i>	65.06	47.68*	1.18**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
<i>Number of Islands</i>	2*	NA	0	*Based on (Henson et al. 2010). Additional islands may be present in the regional unit
Condition	Region	Region %	% of Lake Superior Total	
<i>Population Density (persons/km²)</i>	17.74	NA		
<i>Road Density (km/km²)</i>	0.75	NA		
<i>Number of Dams and Barriers</i>	2478	NA	10.5	
<i>Artificial Shoreline (km)</i>	45.82	70.43	20.64	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
<i>Private</i>	5,354.95	62.28	8,597.74	Regional area based on landmass
<i>Public/Crown</i>	2782.4	32.36	8,597.74	
<i>Tribes/ First Nations</i>	344.67	4.01	8,597.74	
<i>Parks & Protected Areas (total)</i>	115.85	1.35	8,597.74	
<i>Parks & Protected Areas (coast)</i>	4.24	3.11*	136.43**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

IMPORTANT BIODIVERSITY FEATURES

Nearshore and Inshore Waters

- St. Louis Bay is noted as a Lake Superior embayment important for lake sturgeon (Auer 2003).
- North Bay is a proposed site for the Duluth Natural Areas Program (DNAP). North Bay is also located within a Minnesota Biological Survey Site of Outstanding Biodiversity Significance (B. Carlson, pers. comm., March 20 2013).
- Rask Bay is within a Minnesota Biological Survey Site of Outstanding Biodiversity Significance (B. Carlson, pers. comm., March 20 2013).

Coastal Zone and Islands

- The 12,000 acre St. Louis River Estuary wetland complex is one of the largest estuarine wetland complexes in the Lake Superior Basin. Areas described as wilderness-like are still found in the upper estuary (U.S. EPA 2013c), and the estuary shoreline is a mix of forest, wetlands and industry (Minnesota DNR 2012b).
- The estuary provides critical habitat for a number of migrating and breeding birds; up to 230 different species have been observed in the area (Allen 2013). Included in the 115 known species of breeding birds are common terns, which are listed as endangered in Wisconsin and threatened in Minnesota (Allen 2013). Nearly two-thirds of the entire Lake Superior basin population of common terns have nested in the estuary in recent years (Allen 2013).
- In total, 45 species of native fish have been documented in the estuary, and it is an important area for migratory fish species that spawn upstream of Lake Superior (Allen 2013). A number of other species, including wolf, bear, bobcat, mink, beaver and river otter all use the estuary habitat (TNC No date a).
- Spirit Island and Clough Island are key island for biodiversity.
- The Wisconsin Department of Natural Resources (WDNR) has identified primary coastal wetlands deemed to be ecologically significant coastal wetlands. Several of these Lake Superior ecologically significant coastal wetlands are located in the St. Louis/ Cloquet regional unit, including S-01 Red River Breaks – St. Louis River Marshes, S-02 Oliver Marsh, S-03 Superior Municipal Forest and S-04 Pokegama Carnegie Wetlands (WDNR 2012c). A complete list of the Lake Superior ecologically significant wetlands and their specific site attributes is available on the WDNR website (WDNR 2012c, 2012d).

Tributaries and Watersheds

- The St. Louis River is 195 miles (314 kilometres) long and its watershed drains nearly 3,600 square miles (9,300 square kilometres) (Minnesota PCA 2012b). The upper St. Louis River watershed is largely remote and forested, with bluffs and wooded hills (Minnesota DNR 2012a). The Fond du Lac Dam is 21 miles (34 kilometres) upstream of Lake Superior; at the dam there is a marked difference in hydrologic variables between upstream and downstream segments of the river. For this reason agencies like the Minnesota Pollution Control Agency (Minnesota PCA) address the two segments separately (Minnesota PCA 2012b).
- Historically 21 tributaries in Lake Superior had lake sturgeon spawning runs. One of these historical spawning tributaries, the St. Louis River, is in the St. Louis regional unit. The lake sturgeon population status in the St. Louis River is reintroduced and the population trajectory is unknown (Golder Associates Ltd. 2011). Natural reproduction of lake sturgeon in the St. Louis River following 28 years of stocking was first documented in 2011 (W. Blust, pers. comm., March 6 2013). The St. Louis River is one of ten Lake Superior tributaries with recent evidence of natural reproduction (as of

2012). This is the same number of rivers as in 2005, however the specific tributaries have changed (Lake Superior Lake Sturgeon Work Group 2012, unpublished data).

- A Lake Sturgeon Rehabilitation Plan for Lake Superior (Auer 2003) identifies the St. Louis River as one of the seventeen tributaries to Lake Superior in which there should be a focus on lake sturgeon rehabilitation.
- The Cloquet River is a main tributary of the St. Louis River. It flows approximately 100 miles (161 kilometres) from Cloquet Lake to the St. Louis River, and it is relatively undeveloped, especially in the upper reaches (Minnesota PCA 2012b). Above Island Lake are areas classified as both primitive and as natural areas in the St. Louis River Management Plan. This plan also notes the river's exceptional recreational and scenic value (Minnesota PCA 2011a). Farmland, recreational homes and hydro power uses are found in the lower part of the river, but there are no large towns or cities along the river (Minnesota PCA 2011a, Minnesota DNR 2013a). A number of common game fish are found in the Cloquet River, and a small number of brook trout are found in the far upper reaches. Wildlife species found in this area include timber wolves, bobcats, lynx, moose, black bears and bald eagles (Minnesota DNR 2013a). Lakes are prominent features of the landscape in the Cloquet portion of the region, unlike most areas of the Lake Superior basin, which tend to be poorly drained or consist of rivers draining elevated terrain (Minnesota PCA 2011a).
- The Whiteface River is another main tributary of the St. Louis River. It flows 80 miles (129 kilometres) from the Whiteface Reservoir to the St. Louis River (Minnesota PCA 2012b). The federal government owns much of the land in the headwaters of the St. Louis, Cloquet and Whiteface rivers, while land ownership varies in other sections of each river (Minnesota PCA 2012b).
- The St. Louis/Red River Breaks Streambank Protection Area includes 6,500 acres of land owned by the Wisconsin Department of Natural Resources. This area is characterized by steep, branching ravines which flow into the Red River and St. Louis River, and is protected to protect water quality (WDNR 2012b, R. O'Connor, pers. comm., March 15 2013).
- The red clay plain near Superior, Wisconsin is the location of more than a dozen rare plant species, including some that are found nowhere else in Wisconsin (R. O'Connor, pers. comm., March 15 2013).

Figure 12.1: St. Louis/ Cloquet - Coastal and Watershed Features




TABLE 12.2: St. Louis/ Cloquet CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	
Nearshore ¹	D (0.03)	
Embayments and Inshore ^{1,2}	D (0.20)	
Coastal Wetlands ^{2,3}	D (0.300)	Local experts believe a condition score of C may be accurate for coastal wetlands in this regional unit. Several high quality wetlands help to provide some balance for those that are severely degraded (C. Hagen et al., pers. comm., March 20 2013). An estimated 7,700 acres of wetland and open water habitat have been lost from the estuary (U.S. EPA 2013c).
Islands ⁴	A	Local experts believe a condition score of B may be accurate for islands in this regional unit. Many islands have invasive species; if left unmanaged these invasive species will degrade island biodiversity (C. Hagen et al., pers. comm., March 20 2013).
Coastal Terrestrial ³	C (0.500)	
Tributaries and Watersheds ²	D (0.37)	Natural reproduction of lake sturgeon in the St. Louis River following 28 years of stocking was first documented in 2011. The lake sturgeon population had been extirpated due to pollution and overfishing (W. Blust, pers. comm., March 6 2013).

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 12.2: St. Louis/ Cloquet - Condition



IMPORTANT ISSUES & THREATS

- The St. Louis River Area of Concern (AOC) is located in three regional units, including the St. Louis/Cloquet regional unit. The St. Louis River has faced issues of habitat loss and degradation, and pollution and contamination, especially in its lower reaches. The lower 39 miles (63 kilometres) of the St. Louis River are the main focus of the St. Louis River Remedial Action Plan (RAP) (U.S. EPA 2013c) and are designated as a Lake Superior Area of Concern (AOC). The St. Louis River Expanded Study Area encompasses the entire St. Louis/Cloquet regional unit. Nine beneficial use impairments were identified in the St. Louis River AOC (U.S. EPA 2013c).
- The lower St. Louis River has long been used for transportation. Native Americans and early European explorers used the St. Louis River as a linkage between the Great Lakes and the Mississippi River system, by way of a 6 mile portage between the East Savannah River (near Floodwood, Minnesota) and the West Savannah River and the Mississippi River system (Minnesota DNR 2012b). Transportation continues to be a major part of the St. Louis River today, and Great Lakes freighters use the lower reaches as a shipping channel. The Duluth-Superior Harbor is at the end of the estuary, and is one of the busiest ports on the Great Lakes (Minnesota PCA 2012c).
- Major cities and industries are part of the landscape of the lower estuary, with urban development, an industrial harbour and a major port within this area (U.S. EPA 2013c). Dredging restrictions and containment actions in parts of the St. Louis River AOC, in place due to sediments containing elevated levels of toxic, bio-accumulative contaminants, have negative economic consequences, in addition to the negative environmental impacts of the contaminated sediments. Minnesota and Wisconsin are conducting sediment characterization and assessment projects in the St. Louis River AOC, to determine prioritization for areas requiring remediation and restoration (LSBP 2012).
- The ports of Two Harbors, Minnesota, Superior, Wisconsin-Duluth, Minnesota and Presque Isle-Marquette, Michigan have been identified as sites at high risk for invasion by aquatic invasive species, due to ballast water from laker traffic (Rup et al. 2010 as cited in International Joint Commission (IJC) Work Group on Aquatic Invasive Species Rapid Response 2011).
- Pollution and habitat degradation have caused significant impairments, specifically to the St. Louis AOC region. An estimated 7,700 acres of wetland and open water habitat have been lost from the estuary (U.S. EPA 2013c). The continued loss of habitat remains an issue for the St. Louis River (U.S. EPA No date). The loss of habitat in the St. Louis estuary over 100 years ago contributed to the extirpation of the stock of lake whitefish that spawned in this area (Horns et al. 2003). Work by the Great Lakes Environmental Indicators Project has found contamination from polycyclic aromatic hydrocarbons (PAHs) in the St. Louis River, and in other areas of industrial activity (LSBP 2008).
- Levels of mercury in fish tissue remain a cause for impairment in this region. Fish consumption advisories based on mercury and polychlorinated biphenyls (PCBs) are issued by both Minnesota and Wisconsin for Lake Superior and the St. Louis River (U.S. EPA No date). The levels of mercury and PCBs in fish tissues also exceed the levels established for the protection of aquatic life and fish eating birds (U.S. EPA No date).
- The Cloquet River Valley was extensively forested before widespread logging. Estimates put the amount of pine in the valley at eight billion board feet; by 1925 nearly all of the pine stands had been logged. Some small areas of mature white and red pine are still scattered amongst second-growth forests (Minnesota DNR 2013a). During times of extensive logging, loggers blasted away rock formations, curves, and logjams in the St. Louis River, to allow log drives to move freely downriver (Minnesota DNR 2012b). Logging and forest fragmentation continue to be primary stressors of the upper Cloquet River. Forest cover has been converted from the historic conifer forest to the present aspen-dominated forest cover. Efforts to restore the forest to its original composition are underway (C. Hagen et al., pers. comm., March 20 2013).

- Forest fragmentation as a result of housing development is an emerging concern in Minnesota. The forest that is present lacks much of its natural conifer component (E. Perry, pers. comm., February 26 2013).
- Invasive species including but not limited to zebra mussels, round goby, rusty crayfish and purple loosestrife are found in the lower St. Louis River, including on Clough Island (Minnesota DNR 2012b, TNC No date a). Competition from invasive species is a threat to the region (U.S. EPA 2013e). Purple loosestrife in wetlands in the lower estuary in Minnesota and Wisconsin is being subjected to bio-control as a management measure (U.S. EPA 2013c).
- Due to the proximity of the headwaters to the Mesabi Iron range, there is intensive mining (e.g. US Steel-Minntac Mine. In addition to mines in current operation, there are additional proposed mines (e.g. PolyMet-NorthMet Mine) (See Figure 3.4 Mining in Lake Superior Basin in Volume 1: Lakewide Assessment). Future threats to the region include those related to infrastructure transportation changes, including changes to roads, rail and shipping (C. Hagen et al., pers. comm., March 20 2013).
- The Arrowhead Refinery Co. Superfund Site is a 10 acre site located in Hermantown, Minnesota, in St. Louis County. The site is located within a white cedar wetland. The Arrowhead Refinery Co. reclaimed waste oil from 1945 to 1977. Waste sludge amounting to 7,000 cubic yards was disposed of on the property in an unlined 2 acre lagoon. The State of Minnesota found PCBs, phenols, cyanide, lead, barium, arsenic, cadmium, chromium and selenium contaminating the ground and surface water near the site (U.S. EPA 2012a ; U.S. EPA 2013f). The site has now been fully remediated and the MPCA and EPA will begin the process to delist the site from the National Priorities List (NPL) (U.S. EPA 2013f).
- The St. Louis River Superfund Site is two Superfund Sites, the St. Louis River/Interlake/Duluth Tar site (SLRIDT) and the U.S. Steel site. The SLRIDT site is 255 acres of land, river embayments, wetlands and shipping slips located on the St. Louis River, four miles upstream of Lake Superior. The U.S. Steel site is eight miles upstream from Lake Superior, on the St. Louis River. The U.S. Steel site consists of 500 acres of land and 200 acres of sediment. Both sites have been contaminated by past industrial operations and disposal activities, resulting in contamination of soil, groundwater and sediment. The U.S. Steel site 2008 Five Year Review noted that the migration of contaminants to the St. Louis River is occurring (U.S. EPA 2013g)

CONSERVATION IN ACTION

PARKS & PROTECTED AREAS

- Jay Cooke State Park
- Cloquet Valley State Forest
- Superior National Forest
- Boundary Waters Canoe Area Wilderness (within Superior National Forest)
- Savanna Portage State Park
- The Superior Municipal Forest is 4,400 acres and the third largest forest within a city in the United States (City of Superior No date a).
- Wisconsin Point is a 3 mile 229 acre area that combined with Minnesota Point (7 miles long) is the largest freshwater sandbar in the world (City of Superior No date b, City of Duluth 2013a).
- The St. Louis River Estuary has been named a National Estuarine Research Reserve (NERR). The Lake Superior National Estuarine Research Reserve includes estuarine wetlands and red clay bluffs which are steep and highly erodible, as well as Wisconsin Point, the largest freshwater bay mouth sand bar in the world (NERRS 2010).
- Two Wisconsin State Natural Areas and 11 Wisconsin Priority Wetlands are located in the boundaries of the Lake Superior NERR (NERRS 2010).

- The Duluth Natural Areas Program permanently conserves the most ecologically significant lands owned by the City of Duluth and private landowners who have volunteered their lands for consideration (City of Duluth 2013b).
- Magney-Snively Forest (B. Carlson, pers. comm., March 20 2013).
- City of Duluth Mission Creek Forest (B. Carlson, pers. comm., March 20 2013).

EXISTING PROGRAMS & PROJECTS

- The U.S. EPA has developed a state of the art early detection design for invasive fish species, using the St. Louis River Estuary as a case study (G. Czipinski, pers. comm., March 20 2013).
- The U.S. Fish and Wildlife Service is undertaking early detection monitoring for new invasive fish species in the St. Louis River Estuary using the design developed by the U.S. EPA (G. Czipinski, pers. comm., March 20 2013).
- The 1854 Treaty Authority is conducting forage fish community monitoring in the St. Louis River Estuary and propagates and monitors wild rice in the treaty area, including the Cloquet River watershed. (G. Czipinski, pers. comm., March 20 2013).
- The Fond du Lac Band of Lake Superior Chippewa is a significant landholder in this regional unit (C. Hagen et al., pers. comm., March 20 2013).
- The St. Louis/ Cloquet regional unit contains a number of areas identified as biologically important by the Lake Superior Binational Program Habitat Committee (2006) (Table 12.3, Figure 12.3).
- There are a number of Minnesota Biological Survey (MBS) Sites delineated in the Minnesota portion of the St. Louis/ Cloquet regional unit, some of which have been ranked with Outstanding or High Biodiversity Significance, based on statewide ranking criteria. The Minnesota Department of Natural Resource's MBS systematically collects, interprets, and delivers data on the distribution and ecology of native plants, animals, native plant communities, and functional landscapes throughout the state. MBS conducts landscape assessments, field surveys and monitoring activities, and provides data and tools to guide conservation and management within *MBS Sites of Statewide Biodiversity Significance* (MBS Sites). Biodiversity information includes the location and biodiversity significance rank of MBS Sites, the location and status of rare species populations, the type and condition of native plant communities, and, for selected sites, *MBS Ecological Evaluation* reports (Minnesota DNR 2013e, B. Carlson, pers. comm., March 20 2013). The MBS Sites located within the St. Louis/ Cloquet regional unit are Cloquet, Cloquet River, Headwaters and Mission Creek (L. Gerdes, pers. comm., March 18 2013).
- The WDNR has a Designated Waters designation for waterbodies with permit requirements. Designated Waters include Areas of Special Natural Resource Interest (ASNRI), Public Rights Features (PRF) and Priority Navigable Waters (PNW) (WDNR No date). These designations offer protection for various important waters, including Wild Rice Waters and Outstanding and Exceptional Resource Waters (C. Hagen et al., pers. comm., March 20 2013).
- The headwaters of the Stoney Brook watershed are located in the Fond du Lac Band reservation, and they are designated as Outstanding Reservation Reserve Waters (ORRWs) by the Fond du Lac Band (LSBP 2008, N. Schudlt, pers. comm., March 19 2013).
- The Outstanding Reservation Reserve Waters designated by the Fond du Lac Band include the Band's wild rice waters. The designation comes from the Fond du Lac Band's promulgation of water quality standards and antidegradation procedures (N. Schudlt, pers. comm., March 19 2013).
- The St. Louis River Alliance started as a Citizens Advisory Committee, and helped develop the Remedial Action Plan (RAP) for the St. Louis River AOC. The group contributes volunteer hours towards river clean-up, macroinvertebrate monitoring, and invasive species removal. The Lower St.

Louis River Habitat Plan was created by the St. Louis River Alliance, along with city, county, state and federal groups. The plan is used to protect, enhance and restore the river (U.S. EPA 2013c).

- The Duluth Natural Areas Program (DNAP) identifies and conserves ecologically significant city-owned lands, as well as private lands which are voluntarily identified with the support and consent of the landowner. Nominated properties are reviewed by city government and the eligibility of the property is determined by scientific criteria and the development of a management plan (City of Duluth 2013).
- A number of projects involving multiple agencies have resulted in the purchase and protection of lands in this region. The Wisconsin Department of Natural Resources' St. Louis River Streambank Protection Project purchased 6,900 acres along the St. Louis River, the Red River, and the main tributaries of the Red River. Another 22,000 acres were acquired along the St. Louis, Cloquet and Whiteface rivers by the St. Louis River Board, and were then transferred to the Minnesota Department of Natural Resources (U.S. EPA 2013c).
- In 2010 The Nature Conservancy acquired the 358 acre Clough Island, the largest island in the St. Louis River Estuary, in order to preserve critical habitat for migratory and breeding birds and fish (Allen 2013; TNC 2010). The acquisition of Clough Island was aided by WDNR Stewardship Funding, and the US Fish and Wildlife Service. The WDNR became the owner of Clough Island in 2011 (C. Hagen et al., pers. comm., March 20 2013). A recent grant to initiate restoration work on Clough Island included funding for invasive species control and the planting of thousands of conifer trees (R. O'Connor, pers. comm., March 15 2013).
- Reintroduction of lake sturgeon to the estuary has been undertaken by the Minnesota Department of Natural Resources and the Wisconsin Department of Natural Resources (TNC 2010).
- The University of Wisconsin-Extension has been designated by the State of Wisconsin as the lead state agency for the Lake Superior NERR, and has led the development of a five year management plan (University of Wisconsin-Extension 2010). Wisconsin's Wildlife Action Plan identified the St. Louis River Freshwater Estuary complex and the associated wetland and boreal forests as areas of continental significance (NERRS 2010).
- Wisconsin's Wildlife Action Plan identified a number of Conservation Opportunity Areas for Wildlife Species of Greatest Conservation Need. In the Superior Coastal Plain Ecological Landscape several areas of State, Continental and Global Significance were identified, including some in the St. Louis/Cloquet regional unit (WDNR 2008a, 2008b, 2008c).
- Large tracts of land in the Cloquet River Watershed portion of this region are owned by Minnesota Power. The Boulder Lake Environmental Learning Center is sponsored by Minnesota Power and provides programming for recreational users of the 18,000 acre Boulder Lake Management Area and Cloquet River (Minnesota PCA 2011a). Many other organizations work within the Cloquet River Watershed, including the Friends of the Cloquet Valley State Forest (Minnesota PCA 2011a).
- The Wisconsin Wetlands Association has identified a set of representative high quality wetlands in different regions of Wisconsin. These are referred to as Wetland Gems, and were identified by building on existing conservation planning efforts (Wisconsin Wetlands Association No date a). Several Wetland Gems are in the Superior Region, including some in the St. Louis/ Cloquet regional unit (Wisconsin Wetlands Association No date b).
- The Sax-Zim Bog is a 98,657 hectare (243,786 acre) State Important Bird Area (IBA) located within the St. Louis/ Cloquet regional unit (National Audubon Society 2013, 2012). This IBA is a known wintering area for hawks and owls, and also provides habitat for over 240 species of migrant and breeding birds (Minnesota DNR 2013d).

- Wisconsin Point is another State Important Bird Area in the St. Louis/ Cloquet regional unit; this site is 77% open water (National Audubon Society 2013, 2012). The City of Superior and partners have developed a plan for Wisconsin Point (C. Hagen et al., pers. comm., March 20 2013).
- Many other diverse programs and projects focus on parts of the St. Louis/Cloquet regional unit and 34 Citizen-based Groups are noted to do work in the St. Louis- Cloquet regional unit (U.S. EPA 2013h, 2013i). A number of other projects, undertaken by the Minnesota Pollution Control Agency and partner organizations are completed, in progress, or scheduled to start; these include the St. Louis River Citizens River Watch, the Cloquet River Major Watershed WRAP Strategy and the Mercury Pollutant Reduction Plan. The St. Louis River Trail Association promotes recreational opportunities based on good environmental stewardship, and non-motorized hiking along a natural surface trail they aim to build and maintain along the St. Louis River in Northern Minnesota (SLRT 2012). The City of Duluth and the University of Minnesota – Duluth collaborated on an urban stream awareness project, designed to increase public understanding of and demonstrate the importance of aquatic ecosystems (Minnesota PCA 2012c). Other initiatives include the Sand Lake-Seven Beavers Forest Collaborative and St. Louis County Cooperative Weed Management Area (B. Carlson, pers. comm., March 20 2013).

TABLE 12.3: St. Louis/ Cloquet IMPORTANT HABITAT SITES AND AREAS


<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
MN-002	Site	Alborn Fen	Poor Fen, sedge subtype, rare plant habitat
MN-005	Site	Arlberg Bog	Rare plant habitat
MN-010	Site	Blackfoot Lake Peatland	Poor Fen, Sedge subtype, rare plant habitat
MN-024	Site	Cloquet River Macrosite	Rare animal habitat
MN-044	Site	Hornby Lake	Rare plant habitat
MN-048	Area	Jay Cooke State Park	Rare plant habitat, northern hardwood forest
MN-064	Site	Magney Hardwoods Forest	Old growth mixed hardwoods/conifer forest and northern hardwoods forest; rare plant habitat
MN-069	Site	Minnesota Point	Rare plant habitat, unique geomorphic formation, old growth pine forest
MN-086a	Site	Interstate Island	Rare animal habitat, colonial waterbird nesting habitat
MN-086b	Site	Hearding Island	Open dunes, representative natural plant community
MN-086d	Site	Spirit Lake Point	Great Lakes marsh, migratory wildlife habitat
MN-086e	Site	Mud Lake	Great Lakes marsh, migratory wildlife habitat
MN-086f	Site	Fond Du Lac	Great Lakes marsh, migratory wildlife habitat
MN-086g	Site	Grassy Point	Great Lakes wetland complex
MN-086h	Site	Bong Bridge	Rare animal habitat
MN-086i	Site	Swamp Lake	Rare animal habitat
MN-086j	Area	St. Louis Estuary	Great Lakes freshwater estuary, rare plant and animal habitat, colonial waterbird nesting habitat
MN-087	Area	Sand Lake Peatland	Large patterned peatland, significant bog and fen features, rare plant habitat, rare animal habitat, geological processes
MN-101	Site	Tikander	Rare animal habitat, rare plant habitat, colonial waterbird nesting habitat
MN-111	Area	Hemlock Ravine SNA	Rare plant habitat, old growth northern hardwoods, white pine, and eastern hemlock forest
MN-114	Area	Savanna Portage State	Northern hardwood forest

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

		Park	
MN-115	Area	Tettegouche State Park	Northern hardwood forest, Northern hardwood-conifer forest, old growth white cedar forest, complex of forested highlands, wetlands, streams and lakes with high biodiversity, Shoreline cliffs, rare animal habitat
WI-048	Site	Superior Municipal Forest	Great Lakes pine forest, rare plant and animal habitat, Great Lakes freshwater estuary
WI-049	Site	Oliver Wetlands	Great Lakes freshwater estuary, coastal marsh, migratory bird habitat
WI-050	Site	Red River	Rare plant habitat, eroding red clay bluffs, fish spawning habitat

In the St. Louis/Cloquet Unit, there are potentially many additional priority areas for conservation. One is the region around the headwaters of the St. Louis River, including but not limited to the Sand Lake Peatland Scientific and Natural Area. It is a large landscape of high ecological quality storing lots of water for the river, and only parts have been protected.

Figure 12.3: St. Louis/ Cloquet - Important Habitat Sites and Areas




*The St. Louis Red River Streambank Protection Area is not a State Park, but was classified with this category due to the state ownership of the site.

13. Beartrap-Nemadji


HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	C	COASTAL WETLANDS	B
EMBAYMENTS & INSHORE	C	COASTAL TERRESTRIAL	A
TRIBUTARIES & WATERSHEDS	C	OVERALL B	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



The sandstone in the Apostle Islands have eroded into unique cliff formations, including sea caves. Photo credit: National Park Service

Summary/ Description

The Beartrap-Nemadji regional unit is located along the southern Lake Superior coast. It is 6736 km² in size, including the associated nearshore waters. The regional unit boundary extends along the Lake Superior shoreline from near Superior, Wisconsin to past Ashland, Wisconsin. In this Biodiversity Conservation Assessment the Beartrap Creek subwatershed is considered part of the Lower Bad River subwatershed and the Bad-Montreal regional unit. This regional unit delineation reflects local management boundaries as adopted by the Bad River Band of Lake Superior Tribe of Chippewa Indians and the Wisconsin Department of Natural Resources (N. Tillison, pers. comm., April 26 2013). Inland, most of this regional unit is located in Wisconsin; however the western portion of this regional unit extends into Minnesota. Approximately 15% of the regional unit land base is in two Minnesota counties (Carlton and Pine counties) and 85% of the regional unit land base is in Wisconsin (in Bayfield, Ashland and Douglas counties) (USDA NRCS No date c). The reservation of the Red Cliff Band of Lake Superior Chippewa is located in this regional unit. A portion of the reservation of the Bad River Band of Lake Superior Tribe of Chippewa Indians is also located in this regional unit (A. McCammon Soltis, pers. comm., March 19 2013). The eastern portion of this region includes the Bayfield Peninsula, Chequamegon Bay and the Apostle Islands National Lakeshore. This regional unit is referred to as HUC 04010301, and it is part of Subregion 0401 – Western Lake Superior. The Beartrap-Nemadji regional unit contains one tertiary (HUC 8) watershed, Beartrap-Nemadji, and 11 quaternary (HUC 10) watersheds. This region contains more agricultural land than most other regions, and just over 50% of the region is forested. The coast is dominated by rocky shores and cliffs, and many coastal wetlands.

TABLE 13.1: Beartrap-Nemadji BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
Agriculture	472.47	7.01	1,287.56	
Developed	18.35	0.27	351.99	
Forest	3,544.92	52.63	96,789.30	
Associated Nearshore Waters	2,393.41	35.53	15,951.22	
Other	282.26	4.19	7,452.78	
Water (inland)	24.41	0.36	8,283.49	
Total Area	6,735.82	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
Coastline (km)	476.1	NA	8.37	Based on SOLEC shoreline
Sand Beaches (km)	33.99	7.14	5.40*	*% of Lake Superior Total Sand Beaches
Coastal Wetlands (km ²)	85.58	14.09*	8.69**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
Natural Cover in Coastal Zone	537.32	88.47*	9.73**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
Number of Islands	63	NA	2.4	
Condition	Region	Region %	% of Lake Superior Total	
Population Density (persons/km ²)	7.96	NA		
Road Density (km/km ²)	1.03	NA		
Number of Dams and Barriers	2,681	NA	11.3	
Artificial Shoreline (km)	29.41	5.47	13.25	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
Private	2,522.25	58.08	4,342.42	Regional area based on landmass
Public/Crown	1,522.12	35.05	4,342.42	
Tribes/ First Nations	53.98	1.24	4,342.42	
Parks & Protected Areas (total)	339.18	7.81	4,342.42	
Parks & Protected Areas (coast)	176.51	29.06*	607.38*	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- The Beartrap-Nemadji regional unit contains a number of areas identified as Important Habitat for lake whitefish and lake trout, especially in and around the Bayfield Peninsula and the Apostle Islands (Lake Superior Binational Program Habitat Committee 2006) (Figure 13.1). The shoals near the Apostle Islands National Lakeshore's one-quarter mile lakeward boundary are especially important for lake trout and whitefish, as they provide critical spawning areas (NPS 2013b).
- Chequamegon Bay is noted as a Lake Superior embayment important for lake sturgeon (Auer 2003). In the Beartrap-Nemadji regional unit this embayment and the nearshore zone, which provides corridors for movement, are identified as critical management areas for lake sturgeon in the Lake Superior basin (Auer 2003).

- The waters around the Apostle Islands are described as relatively shallow, with a diverse and complex fish community for Lake Superior (NPS 2013b).

Coastal Zone and Islands

- The Wisconsin Department of Natural Resources (WDNR) has identified primary coastal wetlands deemed to be ecologically significant coastal wetlands. This assessment has been completed for both the Lake Superior and Lake Michigan basins. Several of these Lake Superior ecologically significant coastal wetlands are located in the Beartrap-Nemadji regional unit, including S-06 Nemadji River Bottoms, S-08 Allouez Bay-Wisconsin Point, S-09 Port Wing, S-11 Lost Creek, S-17 Big Bay Wetlands and S-22 Mouth of the Brule River (WDNR 2012c, 2012d). A number of the ecologically significant coastal wetlands are located on the islands of the Apostle Islands National Lakeshore (WDNR 2012d). A complete list of the Lake Superior ecologically significant wetlands and their specific site attributes is available on the WDNR website (WDNR 2012c, 2012d).
- A number of Important Habitat Sites are located along the Lake Superior shore, as well as a number of additional sites inland. Several Important Habitat Areas are also located in the Beartrap-Nemadji region; the group of Apostle Islands are one of these Important Habitat Areas (Lake Superior Binational Program Habitat Committee 2006) (Table 13.3, Figure 13.3).
- Resident breeding birds and neotropical migrant birds use the islands within the Apostle Islands National Lakeshore for important habitat. Important migratory bird concentration points are also located in the lakeshore (NPS 2013b).
- Chequamegon Bay is described as among the best Wisconsin birding locations. More than 300 bird species have been observed in the area (a 40-mile radius around Ashland, Wisconsin) since 1972 (NPS 2013b).
- Stockton Islands, part of the Apostle Islands National Lakeshore, has one of the highest concentrations of black bears in North America. Many other mammals, including whitetail deer, snowshoe hare, red fox and beaver are found on the islands. Many common mainland species, such as raccoon, skunk, porcupines and gray squirrels are not found on the islands (NPS 2013b).
- Over 800 plant species are known to occur within the Apostle Islands National Lakeshore. Some islands contain old-growth remnant forests, as they were never commercially logged. Wetland, Precambrian sandstone ledges and bluffs, and many dunal features provide habitat for a number of rare plant species (NPS 2013b).

Tributaries and Watersheds

- There are native, self-sustaining brook trout in many of the tributaries of this regional unit (M. Jennings, pers. comm., March 13 2013).
- The deeply cut streams characteristic of this watershed have influenced land use in the regional unit. The valleys remain largely forested, while flat upland areas tend to be used for agriculture and recreation (USDA NRCS No date c).
- The Beartrap-Nemadji regional unit is in the Northern Lakes and Forest Ecoregion. Portions of the regional unit are located within the Lake Superior Lacustrine Clay Plain and the Minnesota/Wisconsin Upland Till Plain (USDA NRCS No date c).
- The Nemadji River Floodplain Forest contains a floodplain forest which is unusual in its composition, and may be unique to Wisconsin. An unusual mix of tree species and a rich herb layer are found. Floodplain forests are very rare along Lake Superior tributaries. The Nemadji River Floodplain Forest is located along the Nemadji River, on terraces 15 feet above normal water levels. It was designated a State Natural Area in 1997, and is owned by Douglas County (WDNR 2013a).

Figure 13.1: Beartrap-Nemadji - Coastal and Watershed Features




TABLE 13.2: Beartrap-Nemadji CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	
Nearshore ¹	C (0.57)	
Embayments and Inshore ^{1,2}	C (0.51)	
Coastal Wetlands ^{2,3}	B (0.655)	Local experts believe a grade of B may be low for the coastal wetlands target. Although the Nemadji and other smaller wetlands are described as compromised, the Beartrap-Nemadji unit also contains a dozen or more exemplary sites, including Bark Bay, Raspberry Bay, Lost Creek, and others (R. O'Connor, pers. comm., March 15 2013).
Islands ⁴	A	Local experts believe a condition score of B may be accurate for islands in this regional unit. The islands are under continual management to maintain their current good condition. Without continued management the islands' biodiversity would suffer degradation (C. Hagen et al., pers. comm., March 20 2013).
Coastal Terrestrial ³	A (0.956)	Local experts believe a condition score of B may be accurate for the coastal terrestrial portion of this regional unit. Although there is still quite a bit of public land in this area, significant areas of private land have fragmented the riparian corridor (C. Hagen et al., pers. comm., March 20 2013). In addition, the loss of conifers and prevalence of aspen, which is still being heavily promoted by state agencies, are provided as additional reasons that a condition score of A may be too high. Research has demonstrated that from a water quality standpoint, young aspen (0-15 years) is no different than open land (R. O'Connor, pers. comm., March 15 2013).
Tributaries and Watersheds ²	C (0.44)	

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 13.2: Beartrap-Nemadji - Condition



Important Issues & Threats

- The St. Louis River Area of Concern (AOC) is located in three regional units, including portions of the Beartrap-Nemadji regional unit. The St. Louis River has faced issues of habitat loss and degradation, and pollution and contamination, especially in its lower reaches. The lower 39 miles (63 kilometres) of the St. Louis River are the main focus of the St. Louis River Remedial Action Plan (RAP) (U.S. EPA 2013c) and are designated as a Lake Superior Area of Concern (AOC). Nine beneficial use impairments were identified in the St. Louis River AOC (U.S. EPA 2013c).
- Chequamegon Bay has two highly contaminated sites that were not given AOC designations. These are both Superfund sites. One is located on the Ashland waterfront and upland areas and is called the Ashland/Northern States Power Lakefront Site (U.S. EPA 2013j). The second site is a mostly upland area on the west side of Chequamegon Bay (at Barksdale, Wisconsin) called the DuPont Barksdale Explosives Plant Site (U.S. EPA 2011). These are both priority sites for improving the health of Chequamegon Bay (M. Hudson, pers. comm., March 20 2013).
- The populations of native, self-sustaining brook trout in many of the tributaries of this regional unit face a number of threats, since many of the streams are thermally marginal for coldwater fish. Climate change has the potential to make these streams unsuitable for brook trout, as well as non-native but self-sustaining trout and salmon. Maintenance of shaded riparian zones is important to conservation in these streams. Conversion of forest cover to cover types that increase runoff would threaten these fish, as would major changes in availability of groundwater, if future demand increases withdrawals (M. Jennings, pers. comm., March 13 2013).
- The Beartrap-Nemadji watershed is the largest single source of sediment to Lake Superior (USDA NRCS No date c). Because the mouth of the Nemadji River is located in Superior Bay, the U.S. Army Corps of Engineers must dredge Superior Bay annually, to maintain the necessary depth for shipping traffic (USDA NRCS No date c). Estimates completed 15 years ago by the Minnesota Natural Resources Conservation Service (NRCS) indicate that the annual amount of sediment transported by the Nemadji River is on average 120,000 tons, with 33,000 tons of sediment dredged annually (USDA NRCS No date c, Baird & Associates 2000).
- The Nemadji River Basin Project determined that the erosion of valley walls accounts for 98% of the sediment yield from the Nemadji Basin. The amount of sediment eroded and transported along the tributaries of the Nemadji Basin to the mouth of the river (the sediment delivery ratio SDR) was nearly 98% (Baird & Associates 2000).
- A top resource concern noted in Rapid Watershed Assessment for this regional unit is Bankfull Flow and Channel Downcutting. Problems with erosion, sedimentation and turbidity in this regional unit can be linked to the conversion of land use from old growth coniferous forests to forests of poplar. The amount of water yielded by the land increased with this cover change. Incompatible forestry (e.g. clear-cut logging) and agriculture may create the same hydrologic system response as open lands² on the clay plains of the region (USDA NRCS No date c, C. Hagen et al., pers. comm., March 20 2013).
- This regional unit contains clay plains and sand barrens. Management of certain areas of the regional unit as open lands is desired. For example in the Moquah Barrens open grassland is the management goal, and there is not a detrimental effect to the watershed (C Hagen et al., pers. comm., March 20 2013).

² Open lands are defined by Baird & Associates (2000) as meadows, pasture or timber growth aged 0 to 15 years. The definition for open growth is not given in the Rapid Watershed Assessment (USDA No date c)

- The conversion of land cover to coniferous forests in red-clay soil areas is one management opportunity noted for woodland management which would also address erosion concerns (USDA NRCS No date c).
- Development pressure for this regional unit is described as moderate. Development may occur on farms, timberland, or lakeshore areas (USDA NRCS No date c).
- Agricultural land use accounts for approximately eleven percent of the land-base for this regional unit. There are estimated to be approximately 1,617 farms in this regional unit. Of these farms, approximately sixty percent are less than 180 acres in size, thirty-seven percent are between 180 and 1,000 acres in size, and three percent are larger than 1,000 acres (USDA NRCS No date c).
- A large portion of the soils in this regional unit are classified as having several limitations or very severe limitations through Land Capability Classification. These classifications reflect how the soils would fare with typical field crops, the possibility of damage if they were used for field crops, and how they respond to management (USDA NRCS No date c).
- A number of waterbodies within the Beartrap-Nemadji regional unit are listed as impaired. Reasons for impairment include excess nutrients, mercury, degraded habitat, e-coli, contaminated sediments, turbidity, and others. Affected uses include aquatic consumption, aquatic life and aquatic recreation (USDA NRCS No date c).
- An Emergency Prevention and Response Plan for Viral Hemorrhagic Septicemia has been developed for Isle Royale National Park, Pictured Rocks National Lakeshore, Apostle Islands National Lakeshore and the Grand Portage Band of the Lake Superior Chippewa Reservation (within which is the Grand Portage National Monument) (NPS 2013a).
- Concerns identified through the Rapid Watershed Assessment are mainly related to the large amounts of sediments transported in the waters, due to the highly erodible clay soil and high riverbanks (USDA NRCS No date c). The sediment transported through the watershed eventually is deposited into Western Lake Superior (USDA NRCS No date c).

Conservation In Action

Parks & Protected Areas

- Apostle Islands National Lakeshore is located in the Beartrap-Nemadji regional unit. The park is a combination of 21 islands located off the Bayfield Peninsula, and 12 miles of mainland (NPS 2013b).
- Although Long Island is included in the Apostle Islands, it is an extension of the Chequamegon Point barrier spit, and it is the only Apostle Island which does not have a bedrock core.
- Frog Bay acquisition by the Bayfield Nature Conservancy and the Red Cliff Tribe of Lake Superior Chippewa (C. Hagen et al., pers. comm., March 20 2013). The recently designated Frog Bay Tribal National Park by the Red Cliff Tribe is the first national park ever designated by a federally recognized tribe (R. O'Connor, pers. comm., March 15 2013).
- Chequamegon National Forest

Existing Programs & Projects

- The U.S. Fish and Wildlife Service Iron River National Fish Hatchery is located in the Beartrap-Nemadji region. The captive brood stock and production fish from the brook trout of the Isle Royale region (the Tobin Harbor and Siskiwit Bay strains) are reared at this facility (Newman et al. 2003). The Red Cliff Tribal Fish Hatchery is also located in this region. The Red Cliff Tribal Fish Hatchery rears captive brood stock and production fish from the Lake Nipigon strain of brook trout (Newman et al. 2003).
- Bayfield State Fish Hatchery

- The U.S. Fish and Wildlife Service is undertaking early detection monitoring for new aquatic invasive species in Chequamegon Bay (G. Cypinski, pers. comm., March 20 2013).
- Bog Lake on Madeline Islands has been designated as a high quality waterbody through the Bad River Band of Lake Superior Tribe of Chippewa Indians Outstanding Tribal Resource Waters (OTRWs) designation (Bad River Band of Lake Superior Tribe of Chippewa Indians 2011).
- The WDNR has a Designated Waters designation for waterbodies with permit requirements. Designated Waters include Areas of Special Natural Resource Interest (ASNRI), Public Rights Features (PRF) and Priority Navigable Waters (PNW) (WDNR No date). These designations offer protection for various important waters, including Wild Rice Waters and Outstanding and Exceptional Resource Waters (C. Hagen et al., pers. comm., March 20 2013).
- Wisconsin's Wildlife Action Plan identified a number of Conservation Opportunity Areas for Wildlife Species of Greatest Conservation Need. In the Superior Coastal Plain Ecological Landscape several areas of State, Continental and Global Significance were identified, including some in the Beartrap-Nemadji regional unit (WDNR 2008a, 2008b, 2008c).
- The Chequamegon Bay Area Partnership (CBAP) recently published a Strategic Priorities document that outlines important issues and threats to the Chequamegon Bay area (CBAP 2013).
- There are watershed partnership efforts underway to implement the Fish Creek Watershed Restoration and Management Plan and the Marengo River Watershed Action Plan. Fish Creek and the Marengo River quaternary watersheds are two of the largest sediment contributors to the Chequamegon Bay area (M. Hudson, pers. comm., March 20 2013).
- The Bayfield Regional Conservancy, a land trust working within the Beartrap-Nemadji and the Bad-Montreal regional units, has developed strategic conservation plans for areas in these regional units (M. Hudson, pers. comm., March 20 2013).
- The Wisconsin Wetlands Association has identified a set of representative high quality wetlands in different regions of Wisconsin. These are referred to as Wetland Gems, and were identified by building on existing conservation planning efforts (Wisconsin Wetlands Association No date a). Several Wetland Gems are in the Superior Region, including some in the Beartrap-Nemadji regional unit (Wisconsin Wetlands Association No date b).
- The Natural Resources Conservation Service (NRCS) Performance Results System (PRS) provides support for reporting the development and delivery of conservation programs (USDA NRCS No date d). From 1999 to 2007 plans were made for a total of 21,798 acres of Total Conservation Systems. From 1999 to 2007 the Total Conservation Systems Applied amounted to 22,439 acres. The activities which contributed the largest amount to the Total Conservation Systems Applied were Total Wildlife Habitat (10,866 acres), Erosion Control Total Soil Saved (7,912 tons/year), Total Nutrient Management (5,589 acres) and Riparian Forest Buffers (4,198 acres). Additional activities involved pest management systems, prescribed grazing, tree and shrub establishment and wetlands (created, restored or enhanced) (USDA NRCS No date c).
- A number of State Important Bird Areas (IBAs) are located in the Beartrap-Nemadji regional unit. These IBAs are Apostle Islands National Lakeshore IBA, Brule Glacial Spillway IBA, Lower Chequamegon Bay IBA, Moose Junction Peatlands IBA, Wisconsin Point IBA, Moquah Barrens IBA and South Shore Wetlands IBA (National Audubon Society 2013, 2012).
- 27 Citizen-based Groups are noted to do work in the Beartrap-Nemadji regional unit (U.S. EPA 2013r). Additional projects, plans, conservation districts, organizations and partners related to the Beartrap-Nemadji regional unit are noted in the Rapid Watershed Assessment (USDA NRCS No date c). The Chequamegon Bay Area Partnership is a strong partnership in the area (C. Hagen et al., pers. comm., March 20 2013).


TABLE 13.3: Beartrap-Nemadji IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
MN-009	Area	Black Lake Bog SNA	Black ash swamp, fen, forested bog, and open bog plant communities, rare plant habitat.
MN-043	Site	Holyoke	Northern hardwood forest
MN-086j	Area	St. Louis Estuary	Great Lakes freshwater estuary, rare plant and animal habitat, colonial waterbird nesting habitat
MN-089	Site	Soo Line	Northern hardwood forest, rare plant habitat
WI-010	Area	Apostle Islands	Extensive and diverse natural plant and animal communities. Rare plant and animal habitat
WI-012	Site	La Pointe Marina	Coastal wetlands, fish spawning habitat
WI-013	Site	Grant's Point	Coastal wetland and beach
WI-014	Site	Bog Lake and Amnicon Point	Unprotected sand beach, open-water coastal wetland
WI-015	Site	North Fish Creek Watershed	Old growth forest, fish spawning habitat, rare plant habitat, migratory wildlife areas
WI-016	Site	Whittlesey Creek Mouth	Coastal wetlands, fish spawning habitat, groundwater fed river system
WI-018	Site	Sioux River Wetland Estuary	Extensive coastal and riparian wetlands, sand beach and sandstone cliffs, rare plant habitat, shoreline bog
WI-019	Site	Onion River	Small coastal estuary, fish spawning habitat
WI-020	Site	Pikes Creek Slough	Coastal estuary wetland community
WI-021	Site	Schooner Bay/Red Cliff Bay	Small coastal estuary with extensive wetlands.
WI-022	Site	Frog Bay	Small coastal wetland estuary complex
WI-023	Site	Raspberry Bay and River	Great Lakes sand, gravel, and rock shoreline, riparian wetlands, fish spawning habitat
WI-024	Site	Little Sand Bay	Coastal wetlands, sand beach, hemlock and cedar forest
WI-025	Site	Big Sand Bay /Sand River	Small coastal freshwater estuary, good fish habitat
WI-026	Site	Mawikwe Bay Tributaries	Three small tributaries form small coastal wetlands at their mouths, sand beach, lake trout spawning area off point
WI-027	Site	Siskiwit Bay and River	Bay with spawning area for lake whitefish. Shorebirds use the sand beaches of Siskiwit Bay. Riverine fish spawning habitat.
WI-028	Site	Lost Creek Natural Area	Extensive coastal wetland, rare plant habitat, shorebird and fish habitats
WI-029	Site	Bark Bay and Point	Beach dunes landscape, extensive coastal wetlands and bog. The bay supports submergent vegetation.
WI-030	Site	Bark River	Extensive riparian wetlands, cedar and white pine forest, fish habitat
WI-031	Site	Cranberry River State Fishery Area	Great Lakes coastal lagoon and coastal wetlands, diverse fish community
WI-032	Site	Cranberry River Headwaters	Ground water source for fish habitat
WI-033	Site	Flag River Fishery Area & Port Wing Natural Area	Extensive coastal wetland area, old beach ridges, rare plant habitat
WI-034a	Site	Mud Lake	Riparian wetland habitat, bog
WI-034b	Site	Millpond Lake	Riparian wetland habitat
WI-034c	Site	Sand Barrens	Rare plant habitat, representative natural community, ecological processes

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

Code	Site/ Area	Important Habitat Site/Area Name	Key Features
WI-034d	Site	Iron River Mouth	Great lakes coastal wetlands, fish spawning habitat
WI-034e	Area	Iron River Watershed	Fish spawning habitat
WI-035	Site	Reefer Creek	Small freshwater estuary, fish spawning habitat
WI-036	Site	Fish Creek	Small estuary with coastal wetlands, eroding scarps in unconsolidated sediments
WI-037	Site	Martinson's Landing	Freshwater estuary and coastal wetlands, sand beach, erodible clay banks
WI-038	Site	Brule River Watershed	Great Lakes coastal wetlands, fish spawning habitat, old growth forest, riparian wetlands
WI-038	Area	Brule River Watershed	Great Lakes coastal wetlands, fish spawning habitat, old growth forest, riparian wetlands
WI-039	Site	Smith Creek Estuary	Coastal wetlands
WI-040	Site	Pearson Creek Estuary	Coastal wetlands, gravel beach, eroding red clay bluffs
WI-041	Site	Poplar River Estuary	Coastal freshwater wetland estuary, fish spawning habitat
WI-042	Site	Middle River Estuary	Lake Superior freshwater estuary with coastal wetlands, fish spawning habitat, eroding red clay deposits.
WI-043	Site	Amnicon River Estuary	Freshwater estuary with coastal wetlands and sheltered vegetated banks, fish spawning habitat
WI-044	Site	Small Estuaries	Several small streams flow through red clay soils and form small estuaries where they enter Lake Superior, alder thickets, shrub carr wetlands
WI-045	Site	Wisconsin Point & Allouez Bay	Bay mouth bar geological feature, longest freshwater sand spit, sand dune ecosystem, colonial waterbird nesting area, migratory wildlife habitat
WI-046	Site	Nemadji River Mouth	Extensive riparian wetlands, great lakes coastal marsh
WI-047	Site	Hog Island	Shallow open water and wetlands, breeding and migrating waterfowl habitat
WI-051	Site	Chequamegon Bay	Fish spawning habitat, coastal wetlands
WI-052	Site	Ashland Tern Island	Colonial waterbird nesting habitat, rare animal habitat
WI-053	Site	NSP Tern Island	Colonial waterbird nesting habitat, rare animal habitat
WI-055	Area	Big Bay State Park	Coastal lagoon and wetlands complex, floating bog, sand beaches, and unique plant communities.
WI-059	Site	Ashland County, WI (island)	Piping Plover critical habitat site
WI-060	Site	Gull Island Refuge	Critical nesting areas for gulls and colonial waterbirds (gulls, double-crested cormorants)
WI-061	Site	Devil's Island	The Devils Island Formation, between the sandstones, represents deposition across sand flats that were intermittently covered by shallow ponded water

Figure 13.3: Beartrap-Nemadji - Important Habitat Sites and Areas




14. Bad-Montreal

HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	C	COASTAL WETLANDS	B
EMBAYMENTS & INSHORE	C	COASTAL TERRESTRIAL	A+
TRIBUTARIES & WATERSHEDS	C	OVERALL B	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



The Bad River Kakagon Slough complex. Photo supplied by Ryan O'Connor, Wisconsin Department of Natural Resources (WDNR). Photo taken by Christina Isenring, WDNR.

Summary/ Description

The Bad-Montreal region is located along the southern shore of Lake Superior, and is 3,764 km² in size, including the associated nearshore waters. The regional unit extends along the shore from just east of Ashland to the mouth of the Montreal River, and the state boundary line between Wisconsin and Michigan. In this Biodiversity Conservation Assessment the Beartrap Creek subwatershed is considered part of the Lower Bad River subwatershed and the Bad-Montreal regional unit. This regional unit delineation reflects local management boundaries as adopted and managed by the Bad River Band of the Lake Superior Tribe of Chippewa Indians and the Wisconsin Department of Natural Resources (N. Tillison, pers. comm., April 26 2013). These locally adopted management boundaries place the Chequamegon Point barrier spit and Long Island in the Bad-Montreal regional unit. Most of the reservation of the Bad River Band of Lake Superior Tribe of Chippewa Indians is located in this regional unit. Inland, the western and central portions of the regional unit is comprised of three Wisconsin counties (Bayfield, Ashland and Iron, and the easternmost portion extends into one Michigan county (Gogebic County) (USDA NRCS No date d). The 25 mile Penoque-Gogebic Range is found in this regional unit, in Iron and Ashland counties (TNC No date b). This unit is referred to as HUC 04010302, and it is the easternmost region of Subregion 0401 – Western Lake Superior. The Bad and Montreal Rivers are the main tributaries and both drain to Lake Superior. The Montreal River forms the Wisconsin / Michigan state border for thirty miles upstream of Lake Superior (USDA NRCS No date d). Public and tribal ownership account for a large portion of ownership in the regional unit, which is described as forested; wetlands and agriculture also account for some land use (USDA NRCS No date d). The Bad-Montreal regional unit contains one tertiary (HUC 8) watershed, Bad-Montreal, and 7 quaternary (HUC 10) watersheds. The watershed is primarily forest, with some agricultural lands. The shoreline includes extensive coastal wetlands and sand beaches.

TABLE 14.1: Bad-Montreal BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
Agriculture	226.48	6.02	1,287.56	
Developed	6.01	0.16	351.99	
Forest	2,665.95	70.82	96,789.30	
Associated Nearshore Waters	668.61	17.76	15,951.22	
Other	171.78	4.56	7,452.78	
Water (inland)	25.38	0.67	8,283.49	
Total Area	3,764.22	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
Coastline (km)	75.0	NA	1.32	Based on SOLEC shoreline
Sand Beaches (km)	19.44	25.92	3.09*	*% of Lake Superior Total Sand Beaches
Coastal Wetlands (km ²)	42.56	45.01*	4.32**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
Natural Cover in Coastal Zone	90.84	96.07*	1.64**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
Number of Islands	13	NA	0.5	
Condition	Region	Region %	% of Lake Superior Total	
Population Density (persons/km ²)	5.70	NA		
Road Density (km/km ²)	0.96	NA		
Number of Dams and Barriers	1,516	NA	6.4	
Artificial Shoreline (km)	1.13	1.24	0.51	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
Private	1,397.23	45.14	3,095.61	Regional area based on landmass
Public/Crown	1,155.36	37.32	3,095.61	
Tribes/ First Nations	444.64	14.36	3,095.61	
Parks & Protected Areas (total)	111.82	3.61	3,095.61	
Parks & Protected Areas (coast)	1.11	1.17*	94.56**	*% of Regional Coastal Area **Regional Coastal Area (km ²)


Important Biodiversity Features

Nearshore and Inshore Waters

- The nearshore and inshore waters of this regional unit provide areas of Important Habitat for lake whitefish and for lake trout (Lake Superior Binational Program Habitat Committee 2006) (Figure 14.1).
- Chequamegon Bay is noted as a Lake Superior embayment important for lake sturgeon (Auer 2003). In the Bad-Montreal regional unit this embayment and the nearshore zone, which provides corridors for movement, are identified as critical management areas for lake sturgeon in the Lake Superior basin (Auer 2003).
- The waters off of Marble Point on the Bad River Indian Reservation are noted to be critical spawning and nursery habitat for fish and wildlife, including cisco and lake trout (WDNR 1999a).

Coastal Zone and Islands

- The Kakagon and Bad River Sloughs is a 4,355 ha largely undeveloped wetland complex, located at the mouth of the Bad River on Lake Superior (Ramsar & Wetlands International 2013). As of 2012, the Kakagon and Bad River Sloughs site is designated as a Ramsar Wetland of International Importance. The site is comprised of sloughs, bogs and coastal lagoons, and is located in the Bad-Montreal regional unit, to reflect locally adopted management boundaries. The area is located on the Bad River Band of Lake Superior Tribe of Chippewa Indians Reservation, and is a Conservation Area under tribal management through an Integrated Resource Management Plan (Ramsar & Wetlands International 2013, USDA NRCS No date d). The slough is the largest freshwater estuary remaining on Lake Superior and may also be in the most pristine condition; it is an important spawning area for the fish community of Lake Superior and it is the Great Lakes' largest remaining natural wild rice bed (USDA NRCS No date d, BRWA 2013a, Ramsar & Wetlands International 2013).
- Long Island is a site for the Piping Plover (*Charadrius melodus*), an endangered species (Ramsar & Wetlands International 2013).
- The Bad-Montreal regional unit contains several areas which are noted to be Important Habitat Sites, as well as an Important Habitat Area (Lake Superior Binational Program Habitat Committee 2006) (Table 14.3, Figure 14.3).
- The Wisconsin Department of Natural Resources (WDNR) has identified primary coastal wetlands deemed to be ecologically significant coastal wetlands. This assessment has been completed for both the Lake Superior and Lake Michigan basins. Site S-21 Bad River-Kakagon Sloughs is located in the Bad-Montreal regional unit (WDNR 2012d). A complete list of the Lake Superior ecologically significant wetlands and their specific site attributes is available on the WDNR website (WDNR 2012c, 2012d).



Wild rice in the Kakagon and Bad River Slough complex. Photo supplied by Cyrus Hester, Bad River Band of the Lake Superior Tribe of Chippewa Indians. Photo credit: Mike Wiggins Jr.

Tributaries and Watersheds

- Rivers and streams in this regional unit are known to be important to lake sturgeon, brook trout, walleye and introduced salmon, steelhead and brown trout (USDA NRCS No date d, BRWA 2013a, W. Blust, pers. comm., March 6 2013).
- The Bad River provides important spawning habitat for lake sturgeon and walleye (W. Blust, pers. comm., March 6 2013).
- The headwater streams and wetlands of the Bad River watershed are also critical to cold-water fisheries, climate resilience and downstream flow regimes (C. Hester and N. Tillison, pers. comm., March 25 2013, TNC No date b).
- Historically 21 tributaries in Lake Superior had lake sturgeon spawning runs. Three of these historical spawning tributaries, the Bad River, the White River (Wisconsin) and the Montreal River are in the Bad-Montreal regional unit. The Bad River and White River are now recognized as one lake sturgeon population, for which the population status is extant and the population trajectory is stable. The Montreal River population status is extirpated (Golder Associates Ltd. 2011).

- The Bad River is one of ten Lake Superior tributaries where lake sturgeon have currently been documented spawning (as of 2012); this is the same number as 2005, however the specific tributaries have changed (Lake Superior Lake Sturgeon Work Group 2012, unpublished data). The White River (Wisconsin) had been removed as it is a tributary to the Bad River and is not a separate spawning population (Lake Superior Lake Sturgeon Work Group 2012, unpublished data). Genetic studies support the viewpoint that the same fish use spawning locations in both the Bad and White rivers (Lake Superior Lake Sturgeon Work Group 2012, unpublished data).
- The lake sturgeon population in the Bad River is one of two Lake Superior populations which meets the criteria for self-sustaining, as defined in the Auer (2003) Lake Sturgeon Rehabilitation Plan for Lake Superior (Lake Superior Lake Sturgeon Work Group 2012, unpublished data)
- A Lake Sturgeon Rehabilitation Plan for Lake Superior (Auer 2003) identifies the Bad River as one of the seventeen tributaries to Lake Superior in which there should be a focus on lake sturgeon rehabilitation.
- The entire White River, from the headwaters to Lake Superior is a State of Wisconsin fishery, natural or wildlife area (S. Toshner, pers. comm., March 6 2013). Eighteen Mile Creek and Long Lake Branch are two White River tributaries with high ecological importance (WDNR 2013b).
- The Penokee Range is noted to have extensive forests and unusual features, including high-gradient, soft headwater streams and glades of open bedrock (WDNR 2005).

Figure 14.1: Bad-Montreal - Coastal and Watershed Features




TABLE 14.2: Bad-Montreal CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	
Nearshore ¹	C (0.50)	
Embayments and Inshore ^{1,2}	C (0.53)	
Coastal Wetlands ^{2,3}	B (0.674)	Local experts believe a condition score of B may be low for the coastal wetlands target, due to the presence of the Kakagon and Bad River Sloughs. At over 10,000 acres, this wetland complex is the largest, most pristine freshwater estuary and coastal wetland complex on the largest freshwater lake in the world (R. O'Connor, pers. comm., March 15 2013). The B score is driven by the assessment of watershed stresses and the condition of nearshore waters.
Islands ⁴	A	
Coastal Terrestrial ³	A+ (0.972)	Local experts believe a condition score of B may be accurate for the coastal terrestrial portion of this regional unit (C. Hagen et al., pers. comm., March 20 2013).
Tributaries and Watersheds ²	C (0.55)	

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 14.2: Bad-Montreal - Condition



Important Issues & Threats

- Non-native invasive species, including purple loosestrife, ruffe and sea lamprey have been noted as threat to the Lower Bad River Watershed (WDNR 1999b). Significant purple loosestrife infestations along rivers in the watershed were noted by the Great Lakes Indian Fish and Wildlife Commission. Common buckthorn and spotted knapweed have also been noted (WDNR 1999b).
- Invasive species are noted to be a large ecological threat to the Kakagon and Bad River Sloughs (Ramsar & Wetlands International 2013).
- Mining companies own a band of land approximately 22 miles long and 22,000 acres in size in the Penokee Range (TNC No date b). Mining in the Penokee-Gogebic Range could impact much of the headwaters in the Bad River watershed, and is viewed as a significant ecological threat to the Kakagon and Bad River Sloughs (Ramsar & Wetlands International 2013, TNC No date b). Impacts throughout the Bad River watershed will have downstream impacts on the sloughs and Lake Superior. In total, 71 miles of perennial and intermittent waters flow through the mining land, including a number of rivers and streams designated as Exceptional or Outstanding Resource Waters by the State of Wisconsin (TNC No date b).
- The Montreal River watershed was noted to be much influenced by the mining history of the area (WDNR 1999a).
- A number of waterbodies within the Bad-Montreal regional unit are listed as impaired. Reasons for impairment include mercury and PCBs. Affected uses include aquatic consumption (USDA NRCS No date c, U.S. EPA 2013k).
- Excessive sediments are a concern in the Bad River Watershed, a "flashy" system in the red clay plain of the Great Lakes Basin. Land use practices, such as silviculture and agricultural practices, have great potential to cause erosion and/or sedimentation problems, particularly if best management practices are not properly implemented and maintained (Bad River Band of Lake Superior Tribe of Chippewa Indians 2006).
- Erosion and slumping of streambanks, channels and gullies leading to sedimentation is the largest non-point resource concern in the Bad-Montreal regional unit. Sediments have a negative effect on fish spawning sites, fish movements, aquatic habitats and Lake Superior (USDA NRCS No date d). The presence of red clay soils interspersed with sands is one natural factor that contributes to this problem. Some water courses are also deeply entrenched, with high banks of up to seventy feet in some locations (USDA NRCS No date d). Other factors contributing to the erosion and slumping include the conversion of native forests to grass and aspen, and in some areas, overgrazing (USDA NRCS No date d). The strategy in place to address this issue is to "slow the flow" of runoff, minimizing channel degradation and erosion (C. Hagen et al., pers. comm., March 20 2013).
- The majority of soils in the regional unit (57%) are classified as poorly suited to most kinds of field crops, based on Land Capability Classification. Other classifications for the soil in the regional unit include moderately well suited (32%), well-suited (5%) and unsuited (5%). These classifications reflect generally how suitable the soils are for typical field crops (USDA NRCS No date d).
- Nutrients entering the watershed from private septic systems and barnyards are a concern (USDA NRCS No date d). Nutrients and bacteria entering the watershed from private septic systems, agriculture (e.g., livestock management, etc.), and municipal wastewater treatment are a concern, particularly in the Marengo River and Beartrap Creek Subwatersheds (Bad River Band of Lake Superior Tribe of Chippewa Indians 2006).
- Hydroelectric power production activities associated with the Gile Flowage may have a negative impact on the fisheries. The Gile Flowage is associated with the Montreal River watershed (USDA NRCS No date d).

- Forest fragmentation through real estate development is an emerging concern in the Upper Peninsula Lake Superior watersheds. Large tracts of forest lands owned by corporate land holders are being sold to companies which run real estate investment trusts; smaller parcels are then developed (W. Taft, pers. comm., February 25, 2013).

Conservation In Action

Parks & Protected Areas

- Bad River Band of the Lake Superior Tribe of Chippewa Indians' Reservation
- Chequamegon National Forest
- Ottawa National Forest
- Copper Falls State Park
- Iron County Forest

Existing Programs & Projects

- A number of waterbodies in the Wisconsin portion of this regional unit have been designated as high quality waters through the state Outstanding Resource Waters (ORWs) or Exceptional Resource Waters (ERWs)³ designations (USDA NRCS No date d). Over 60 waterbodies, including the Bad River and the Bad River Slough had received one of these designations at the time of the Rapid Watershed Assessment (USDA NRCS No date d).
- All of the waterbodies within the Bad River Indian Reservation portion of this regional unit have been designated as high quality waters through the tribal Outstanding Tribal Resource Waters (OTRWs), ORWs or ERWs designations. The Kakagon and Bad River Slough complex, along with the majority of the Bad River itself, are considered OTRWs, waters supporting wild rice and other sensitive and unique resources (Bad River Band of Lake Superior Tribe of Chippewa Indians 2011).
- The Bad River Band of Lake Superior Tribe of Chippewa Indians has a Non-point Source Management Program for the Bad River, through the Bad River Natural Resources Department (C. Hester and N. Tillison, pers. comm., March 25 2013).
- The WDNR has a Designated Waters designation for waterbodies with permit requirements. Designated Waters include Areas of Special Natural Resource Interest (ASNRI), Public Rights Features (PRF) and Priority Navigable Waters (PNW) (WDNR No date). These designations offer protection for various important waters, including Wild Rice Waters and Outstanding and Exceptional Resource Waters (C. Hagen et al., pers. comm., March 20 2013).
- Wisconsin's Wildlife Action Plan identified a number of Conservation Opportunity Areas for Wildlife Species of Greatest Conservation Need. In the Superior Coastal Plain Ecological Landscape several areas of State, Continental and Global Significance were identified, including some in the Bad-Montreal regional unit (WDNR 2008a, 2008b, 2008c).
- Road stream crossings which are barriers to fish passage and contribute to increased sedimentation have been identified as major concerns in the Bad River watershed. The Bad River Watershed Association (BRWA) and partners, including the US Fish and Wildlife Service, Wisconsin DNR, Ashland County, Iron County, Bayfield County, local towns and local landowners are working to address the issue through the Culvert Restoration Program. The program identifies and inventories all road/stream crossings in the Bad River watershed, and prioritizes crossings which need repair. Education and finding sources of funding are additional facets of the program (BRWA 2013b).

³ ORWs receive the highest protection standards Wisconsin offers, while ERWs receive the next highest protection. ORWs usually do not have any point source pollution, whereas waters with an existing point source are likely to be designated ERWs (WDNR 2012a).

- The Chequamegon Bay Area Partnership (CBAP) recently published a Strategic Priorities document that outlines important issues and threats to the Chequamegon Bay area (CBAP 2013).
- There are watershed partnership efforts underway to implement the Fish Creek Watershed Restoration and Management Plan and the Marengo River Watershed Action Plan. Fish Creek and the Marengo River quaternary watersheds are two of the largest sediment contributors to the Chequamegon Bay area (M. Hudson, pers. comm., March 20 2013, C. Hagen et al., pers. comm., March 20 2013).
- The Bayfield Regional Conservancy, a land trust working within the Beartrap-Nemadji and the Bad-Montreal regional units, has developed strategic conservation plans for areas in these regional units (M. Hudson, pers. comm., March 20 2013).
- The White River Properties Group (WRPG) Draft Master Plan and Environmental Assessment is currently in development. The final Master Plan should be finished in fall 2013 (S. Toshner, pers. comm., March 6 2013).
- The Wisconsin Wetlands Association has identified a set of representative high quality wetlands in different regions of Wisconsin. These are referred to as Wetland Gems, and were identified by building on existing conservation planning efforts (Wisconsin Wetlands Association No date a). Several Wetland Gems are in the Superior Region, including the Kakagon Sloughs in the Bad-Montreal regional unit (Wisconsin Wetlands Association No date b).
- A number of State Important Bird Areas (IBAs) are located in the Bad-Montreal regional unit. These IBAs are Bibon Swamp IBA, Camp Nine Pines IBA, Kakagon-Bad River Wetlands and Forest Corridor IBA, Moose Lake Old-Growth Forest-Muskeg IBA, Owen-Teal Forest IBA, Penokee Range IBA and St. Peter's Dome-North Country IBA (National Audubon Society 2013, 2012).
- The Natural Resources Conservation Service (NRCS) Performance Results System (PRS) provides support for reporting the development and delivery of conservation programs. From 1999 to 2007 plans were made for a total of 50,367 acres of Total Conservation Systems. From 1999 to 2007 the Total Conservation Systems Applied amounted to 17,080 acres. The activities which contributed the largest amount to the Total Conservation Systems Applied were Total Wildlife Habitat (9,920 acres), Erosion Control Total Soil Saved (5,739 tons/year), Riparian Forest Buffers (5,661 acres) and Total Nutrient Management (3,137 acres). Additional activities involved prescribed grazing, residue management, wetlands (created, restored or enhanced) and tree and shrub establishment (USDA NRCS No date c).
- The watershed assessment score is used to assess the agricultural non-point pollution potential of Wisconsin watersheds, relative to one another. Based on the criteria used in the model⁴, watershed assessment values ranged from 0.0 (lowest conservation need) to 24.4 (highest conservation need). The score for the Bad Montreal Watershed assessment was 1.6 (USDA NRCS No date d).
- A large portion of the Bad-Montreal watershed is a riparian project area of the Conservation Reserve Enhancement Program (CREP). Landowners who agree to fifteen year agreements may have filter strips, riparian buffers and grassed waterways installed, with annual payments available. In Bad-Montreal this project is located in three counties (northern Bayfield, Ashland and Iron) (USDA NRCS No date d).
- The "Slow the Flow" Program has been adopted by the Lake Superior Partner team, as a way to help manage water run-off in clay soils (University of Wisconsin Extension 2011).
- Trout Unlimited – Wild Rivers Chapter is very active in the Bad-Montreal watershed (S. Toshner, pers. comm., March 6 2013).

⁴ Factors used to calculate the watershed assessment score in the model included acres of cropland, acres of highly erodible land, and the number of animal units in the watershed (USDA No date d).


Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

- Friends of the White River is very active in the Bad-Montreal watershed (S. Toshner, pers. comm., March 6 2013).
- Key partners in the Bad-Montreal regional unit include the Chequamegon Bay Area Partnership and the Bad River Watershed Association (C. Hagen et al., pers. comm., March 20 2013). Additional partners in the Bad-Montreal regional unit are noted in the Rapid Watershed Assessment (USDA NRCS No date d).
- 25 Citizen-based Groups are noted to do work in the Bad-Montreal regional unit (U.S. EPA 2013k).
- The White River Property Group (WRPG) is a combination of state managed lands, including approximately 1,000 acres of State Wildlife Area (White River Wildlife Area), 4,698 acres of Fishery Area (White River Fishery Area, including the Sajdak Springs State Natural Area which is located within the Fishery Area boundary) and 9,263 acres of State Natural Area (Bibon Swamp State Natural Area). In total nearly 15,000 acres are protected as state managed lands, although in the middle segment of the White River Fishery Area project boundary 90% of land is privately owned (WNDR 2013b).

TABLE 14.3: Bad-Montreal IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
WI-001	Site	Montreal River Mouth	Great Lakes costal wetland, old growth white cedar forest
WI-002	Site	Saxon Harbor	Lake Superior beach, fish spawning area
WI-003	Site	Graveyard Creek	Coastal wetlands, spawning habitat for brook and rainbow trout and coho salmon
WI-004	Site	Marble Point	Spawning habitat for lake trout, exposed rocky cliff shore
WI-005	Area	Kakagon Sloughs/Bad River	Largest, healthiest fully-functioning estuarine system in the upper Great Lakes. Rare plant and animal habitat, high biodiversity
WI-006	Site	Honest John Lake	Coastal wetland includes bog communities, patches of open water, sedge meadow, low shrub and lowland coniferous forest
WI-007	Site	Oak Point	Coastal wetland. Bog communities are present as linear strips occupying swales between the forested ridges
WI-008	Site	Long Island	Lake dune landforms, sand beach. Emergent vegetation is common offshore. Rare plant and animal habitat
WI-009	Site	Bibon Swamp	Large wetland complex, largest cold water stream system
WI-034e	Area	Iron River Watershed	Fish spawning habitat
WI-054	Area	Copper Falls State Park	River gorge with falls, old growth mixed northern hardwoods forest
WI-056	Area	Rainbow Lake Wilderness Area	Representative plant communities, northern hardwood and mixed conifer/deciduous forest communities, old growth forest, diverse habitat types, rare plant and animal habitat.

Figure 14.3: Bad-Montreal - Important Habitat Sites and Areas



15. Black-Presque Isle /Ontonagon


HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	NA
NEARSHORE	C	COASTAL WETLANDS	B
EMBAYMENTS & INSHORE	C	COASTAL TERRESTRIAL	A-
TRIBUTARIES & WATERSHEDS	B	OVERALL B	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



Maple Creek. Photo credit: William Taft/ Michigan Department of Environmental Quality.

Summary/ Description

The Black-Presque Isle/ Ontonagon regional unit is located along the southern shore of Lake Superior, (HUC 04020101 and HUC 04020102) and extends from the Montreal River in the west to the community of Ontonagon in the east. The regional unit is 6,150.48 km² in size, including the associated nearshore waters. The Porcupine Mountains, Porcupine Mountains State Park and the Ottawa National Forest are all located in the Black-Presque-Isle/ Ontonagon region. The majority of the land-base of this regional unit is in Michigan), however the southernmost portion of the land-base of the regional unit extends into Wisconsin. It is part of Subregion 0402 – Southern Lake Superior-Lake Superior. The Black-Presque Isle/ Ontonagon regional unit combines two tertiary (HUC 8) watersheds, Black-Presque Isle and Ontonagon, and contains 8 quaternary (HUC 10) watersheds. The watersheds of this unit are dominated by forests. The coast is characterized by sand beaches, rocky shore and extensive coastal wetland in the east.

TABLE 15.1: Black-Presque Isle/ Ontonagon BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
Agriculture	32.98	0.54	1,287.56	
Developed	6.97	0.11	351.99	
Forest	5,109.44	83.07	96,789.30	
Associated Nearshore Waters	588.77	9.57	15,951.22	
Other	299.79	4.87	7,452.78	
Water (inland)	112.53	1.83	8,283.49	
Total Area	6,150.48	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
Coastline (km)	94.9	NA	1.67	Based on SOLEC shoreline
Sand Beaches (km)	24.00	25.29	3.81*	*% of Lake Superior Total Sand Beaches
Coastal Wetlands (km ²)	46.33	25.37*	4.71**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
Natural Cover in Coastal Zone	180.21	98.70*	3.26**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
Number of Islands	0	NA	0	
Condition	Region	Region %	% of Lake Superior Total	
Population Density (persons/km ²)	2.54	NA		
Road Density (km/km ²)	0.58	NA		
Number of Dams and Barriers	2,639	NA	11.2	
Artificial Shoreline (km)	1.05	0.58	0.47	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
Private	2,728.68	49.15	5,551.81	Regional area based on landmass
Public/Crown	2,601.59	46.86	5,551.81	
Tribes/ First Nations	0.54	0.01	5,551.81	
Parks & Protected Areas (total)	221.02	3.98	5,551.81	
Parks & Protected Areas (coast)	62.29	34.11*	182.59**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- The Black-Presque-Isle/Ontonagon region contains several sites noted to be Important Habitat for lake trout. This region also contains two Important Habitat Areas and one Important Habitat site (Lake Superior Binational Program Habitat Committee 2006) (Figure 15.1, Table 15.3, Figure 15.3).

Coastal Zone and Islands

- This region contains extensive sand beaches, including large protected coastal areas in the Porcupine Mountains State Park.

Tributaries and Watersheds

- Historically 21 tributaries in Lake Superior had lake sturgeon spawning runs. The Ontonagon River is one of these historical spawning tributaries, and is located in the Black-Presque Isle/ Ontonagon regional unit. The Ontonagon River population status is reintroduced and the population trajectory is unknown (Golder Associates Ltd. 2011).
- A Lake Sturgeon Rehabilitation Plan for Lake Superior (Auer 2003) identifies the Ontonagon River as one of the seventeen tributaries to Lake Superior in which there should be a focus on lake sturgeon rehabilitation. Habitat restoration is a priority in the Ontonagon River, due to the loss of all wetland habitats (Auer 2003).
- Segments of the Black River, Presque Isle River and Ontonagon River have all been designated as portions of the National Wild and Scenic Rivers System (Interagency Wild & Scenic Rivers Council 2012)

Figure 15.1: Black-Presque Isle/ Ontonagon - Coastal and Watershed Features




TABLE 15.2: Black-Presque Isle/ Ontonagon CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	
Nearshore ¹	C (0.48)	
Embayments and Inshore ^{1,2}	C (0.54)	
Coastal Wetlands ^{2,3}	B (0.627)	
Islands ⁴	NA	No islands have been documented
Coastal Terrestrial ³	A- (0.800)	
Tributaries and Watersheds ²	B (0.60)	

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 15.2: Black-Presque Isle/ Ontonagon - Condition



Important Issues & Threats

- A number of waterbodies within the Black-Presque Isle/ Ontonagon regional unit are listed as impaired. Reasons for impairment include mercury and PCBs in the water column, copper, E. coli and mercury and PCBs in fish tissues. Affected uses include aquatic consumption (U.S. EPA 2013I, 2013am).
- Forest fragmentation through real estate development is an emerging concern in the Upper Peninsula Lake Superior watersheds. Large tracts of forest lands owned by corporate land holders are being sold to companies which run real estate investment trusts; smaller parcels are then developed (W. Taft, pers. comm., February 25, 2013).
- The Copperwood Project of Orvana Resources US Corp. (a subsidiary of Orvana Minerals Corp.) is a mineral lease covering 712 hectares in the Black-Presque Isle/ Ontonagon regional unit. Proven and probable reserves of both copper and silver have been identified. A mining permit was received from the Michigan Department of Environmental Quality in April 2012, and additional permits related to wetlands, discharges and air-quality are under review by the Michigan DEQ. Production at the mine is expected to start in 2014 (Orvana Minerals Corp. 2013). The Copperwood Measured and Indicated Mineral Resource Area is located in close proximity to the Lake Superior shoreline.

Conservation In Action

Parks & Protected Areas

- Ottawa National Forest
- Agate Falls Scenic Site
- Bond Falls Scenic Site
- Lake Gogebic State Park
- Porcupine Mountains Wilderness State Park
- State Forest – Baraga Management Unit
- Twin Lakes State Park

Existing Programs & Projects


- Under the Michigan Water Quality Standards (WQS), portions of the Ontonagon River (Ontonagon County) are designated as Outstanding State Resource Waters (OSRW). Also under the Michigan WQS, all surface waters of the Lake Superior basin that are not identified as OSRWs are designated as Lake Superior basin - Outstanding International Resource Waters (LSB-OIRW). Under the above designations, additional anti-degradation controls are applied for new or increased pollutant loadings (Michigan DEQ 2013a).
- A 14.0 mile segment of the Black River, from the Ottawa National Forest boundary to Lake Superior, is designated as Scenic through the National Wild and Scenic Rivers System (Interagency Wild & Scenic Rivers Council 2012, USDA Forest Service 2007a).
- 72 miles of the Presque Isle River and its tributaries are designated as Recreational or Scenic through the National Wild and Scenic Rivers System. A total of 24.0 miles are designated as Scenic and 48.0 miles are designated as Recreational (Interagency Wild & Scenic Rivers Council 2012, USDA Forest Service 2007a).
- 170 miles of the Ontonagon River and its tributaries are designated as Recreational, Wild or Scenic through the National Wild and Scenic Rivers System. A total of 43.0 miles are designated as Wild, 35.0 miles are designated as Scenic, and 92.0 miles are designated as Recreational (Interagency Wild & Scenic Rivers Council 2012, USDA Forest Service 2007a).

- The Winegar Moraine and Iron Co. Northern Goshawk State Important Bird Area (IBA) is located in the Black-Presque Isle/ Ontonagon regional unit (National Audubon Society 2013, 2012).
- The Western Peninsula Invasives Coalition is the Cooperative Weed Management Area serving all of Gogebic, Iron, and Ontonagon Counties, including the areas covered by the Ottawa National Forest. The mission of the Western Peninsula Invasives Coalition is the prevention, containment and management of non-native invasive species (M. Preisser, pers. comm., May 31 2013).
- 25 Citizen-based Groups are noted to do work in the Black-Presque Isle/ Ontonagon regional unit (U.S. EPA 2013l, 2013m).

TABLE 15.3: Black-Presque Isle/ Ontonagon IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
MI-005	Site	Long Lake	Rare plant and animal habitats, proposed wilderness
MI-042	Area	Porcupine Mountains	Mesic Northern Forest, rare plant and animal habitat
MI-044	Area	Sylvania Wilderness Area	Large area of undisturbed presettlement vegetation, old growth forest, pristine lakes, diverse habitat types, rare plant and animal habitat.

Figure 15.3: Black-Presque Isle/ Ontonagon - Important Habitat Sites and Areas



16. Keweenaw Peninsula/ Sturgeon


HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	B	COASTAL WETLANDS	B
EMBAYMENTS & INSHORE	C	COASTAL TERRESTRIAL	A
TRIBUTARIES & WATERSHEDS	C	OVERALL B	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



Queen Anne's Falls on Slaughterhouse Creek, Houghton County. Photo credit: William Taft/ Michigan Department of Environmental Quality

Summary/ Description

The Keweenaw Peninsula/ Sturgeon regional unit (HUC 04020103 and HUC 04020104) extends from the community of Ontonagon in the west to Jacobsville in the east. This regional unit is 6,077 km² in size, including the associated nearshore waters and encompasses most of the Keweenaw Peninsula. A portion of the Keweenaw Bay Indian Community is located in this regional unit, while the rest of the community is in the Dead-Kelsey regional unit. The Keweenaw Peninsula/ Sturgeon regional unit is within the boundary of the 1842 Ceded Territory. It is part of Subregion 0402 – Southern Lake Superior-Lake Superior. Part of this region (HUC 04020104, Sturgeon) is entirely inland (with no Lake Superior shoreline), although a portion of this HUC extends up the Keweenaw Peninsula and comes quite close to Keweenaw Bay. The Keweenaw Peninsula/ Sturgeon regional unit combines two tertiary (HUC 8) watersheds, Keweenaw Peninsula and Sturgeon, and contains 7 quaternary (HUC 10) watersheds. The watersheds are characterized by forests with some agricultural lands. The coast includes rocky shores and sand beaches. Coastal wetlands are a common habitat along the eastern side of Keweenaw Peninsula.

TABLE 16.1: Keweenaw Peninsula/ Sturgeon BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
Agriculture	149.27	2.46	1,287.56	
Developed	9.25	0.15	351.99	
Forest	3,296.90	54.25	96,789.30	
Associated Nearshore Waters	1,808.62	29.76	15,951.22	
Other	723.30	11.90	7,452.78	
Water (inland)	89.58	1.47	8,283.49	
Total Area	6,076.91	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
Coastline (km)	281.6	NA	4.95	Based on SOLEC shoreline
Sand Beaches (km)	84.90	30.15	13.49*	*% of Lake Superior Total Sand Beaches
Coastal Wetlands (km ²)	187.44	39.80*	19.04**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
Natural Cover in Coastal Zone	404.69	85.93*	7.33**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
Number of Islands	44	NA	1.7	
Condition	Region	Region %	% of Lake Superior Total	
Population Density (persons/km ²)	9.70	NA		
Road Density (km/km ²)	0.72	NA		
Number of Dams and Barriers	1,835	NA	7.8	
Artificial Shoreline (km)	3.36	0.83	1.51	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
Private	3,410.71	80.05	4,260.56	Regional area based on landmass
Public/Crown	756.02	17.74	4,260.56	
Tribes/ First Nations	10.91	0.26	4,260.56	
Parks & Protected Areas (total)	82.99	1.95	4,260.56	
Parks & Protected Areas (coast)	9.48	2.01*	470.94**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- The Keweenaw Peninsula/ Sturgeon regional unit contains many sites of Important Habitat for lake trout and lake whitefish. The Important Habitat sites for both species are found along the shore throughout the peninsula, as well as in and around Grand Traverse and Little Traverse bays (Lake Superior Binational Program Habitat Committee 2006) (Figure 16.1).
- Misery Bay and Bete Grise Bay are noted as Lake Superior embayments which are important for lake sturgeon (Auer 2003). In the Keweenaw Peninsula/ Sturgeon regional unit these embayments and the nearshore zone, which provides corridors for movement, are identified as critical management areas for lake sturgeon in the Lake Superior basin (Auer 2003).
- Buffalo Reef is an important spawning reef for both lake trout and lake whitefish (LSBP 2008).

Coastal Zone and Islands

- This region contains Important Habitat Areas and Important Habitat Sites; a number of these areas and sites are concentrated at the tip of the Keweenaw Peninsula (Lake Superior Binational Program Habitat Committee 2006) (Table 16.3, Figure 16.4).

Tributaries and Watersheds

- The Sturgeon River contains a native stock of lake sturgeon (Auer and Baker No date). Historically 21 tributaries in Lake Superior had lake sturgeon spawning runs. The Sturgeon River is one of these historical spawning tributaries, and is located in the Keweenaw Peninsula/ Sturgeon regional unit. The Sturgeon River population status is extant and the population trajectory is stable (Golder Associates Ltd. 2011).
- The Sturgeon River is one of ten Lake Superior tributaries where lake sturgeon have currently been documented spawning (as of 2012). This is the same number of rivers as in 2005, however the specific tributaries have changed (Lake Superior Lake Sturgeon Work Group 2012, unpublished data).
- The lake sturgeon population in the Sturgeon River is one of two Lake Superior populations which meets the criteria for self-sustaining, as defined in the Auer (2003) Lake Sturgeon Rehabilitation Plan for Lake Superior (Lake Superior Lake Sturgeon Work Group 2012, unpublished data).
- A Lake Sturgeon Rehabilitation Plan for Lake Superior (Auer 2003) identifies the Sturgeon River as one of the seventeen tributaries to Lake Superior in which there should be a focus on lake sturgeon rehabilitation.

Figure 16.1: Keweenaw Peninsula/ Sturgeon - Coastal and Watershed Features




TABLE 16.2: Keweenaw Peninsula/ Sturgeon CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	
Nearshore ¹	B (0.66)	
Embayments and Inshore ^{1,2}	C (0.58)	
Coastal Wetlands ^{2,3}	B (0.705)	
Islands ⁴	A	
Coastal Terrestrial ³	A (0.966)	
Tributaries and Watersheds ²	C (0.49)	

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)


Figure 16.2: Keweenaw Peninsula/ Sturgeon - Condition



Important Issues & Threats

- Mining activities have impacted nearshore Lake Superior sediments in the Keweenaw Region; in the Keweenaw Peninsula area, the sediments contain copper enrichments (Kerfoot et al. 2009). Investigations have revealed a metal-rich area around the Keweenaw Peninsula, where copper tailings have drifted from their original sources (Kerfoot et al. 2012). Three large copper mills operated in the Keweenaw Bay area; two were located at Gay, Michigan, and one was north of Baraga, near Assinins (Kerfoot et al. 2012). The two mills at Gay were the Mohawk and Wolverine Mills, combined they discharged 22.7 teragrams (Tg) of stamp sands (coarse sand waste leftover from the processing of ore), amounting to approximately 89% of the total stamp sands discharged into the Keweenaw Bay area (Kerfoot et al. 2012). Over a century after their initial discharge, tailings on the east side of the Keweenaw Peninsula in the Grand Traverse Bay area have drifted and increased the total surface area of shoreline covered (by 178%), as well as the underwater surface area of the bay bottom covered (by 567%) (Kerfoot et al. 2012) (See Figure 16.3). The U.S. Environmental Protection Agency and the U.S. Army Corps of Engineers – Detroit District estimate that stamp sands cover 1,426 acres of the shoreline and lake bottom (Mattes No date). The most affected shoreline area is between Tobacco River and the Traverse River breakwall (Kerfoot et al. 2012, Mattes No date). Important spawning and nursery areas such as Buffalo Reef are expected to be impacted. Whitefish and lake trout use this reef, and these are important fisheries for the Keweenaw Bay tribal members (Kerfoot et al. 2012).
- The Torch Lake Area of Concern (AOC) is located in the Keweenaw Peninsula/ Sturgeon regional unit, in Houghton County. The 2,700 acre Torch Lake was the site of approximately half of the copper processing in the area between the mid 1840s and 1968 (U.S. EPA 2013n). Copper ore tailings totalling approximately 200 million tons were deposited into Torch Lake during this time, displacing 20% of the lake's original volume (Michigan DNR 1987). The lake was contaminated by other process wastes as well, including other heavy metals, polychlorinated biphenyls, and polycyclic aromatic hydrocarbons (Michigan DEQ 2013). The 1987 Remedial Action Plan (RAP) for the site identified three Beneficial Use Impairments (BUIs) for the Torch Lake AOC: fish tumors, degraded benthos and fish consumption advisories. The fish tumor BUI was removed in 2007. As of 2013, the two other BUIs remain. Federal, state, and local partners continue to investigate ways to restore the site.
- The Torch Lake Superfund Site is located in Houghton County, Michigan. The Superfund Site designation is due to the contamination of the sediments and shoreline of the lake due from copper mining activities (U.S. EPA 2013o).
- The presence of Emerald Ash Borer (EAB) has been detected in counties in the Keweenaw Peninsula/ Sturgeon regional unit. The area is now under a number of quarantine measures, including Federal EAB quarantine and Michigan state quarantine (Cooperative Emerald Ash Borer Project 2013).
- Forest fragmentation through real estate development is an emerging concern in the Upper Peninsula Lake Superior watersheds. Large tracts of forest lands owned by corporate land holders are being sold to companies which run real estate investment trusts; smaller parcels are then developed (W. Taft, pers. comm., February 25, 2013).

Figure 16.3: Location of Stamp Sands in the Keweenaw Peninsula area



Source for Stamp Sands data: U.S. Department of Agriculture, Natural Resources Conservation Service 2000.

Conservation In Action

Parks & Protected Areas

- Ottawa National Forest
- McLain State Park
- State Forest – Baraga Management Unit
- Sturgeon River Sloughs State Wildlife Management Area
- Twin Lakes State Park
- Keweenaw National Historical Park

Existing Programs & Projects

- The Sturgeon River (part of the Sturgeon unit) contains a native stock of lake sturgeon. Eggs from the Sturgeon River population are being used in a lake sturgeon reintroduction program in the Ontonagon River (Auer and Baker No date). The Sturgeon River population is the nearest native stock to the Ontonagon River.
- 28.0 miles of the Sturgeon River, from where it enters the Ottawa National Forest to the Ottawa National Forest northern boundary, are designated through the National Wild and Scenic Rivers System (Interagency Wild & Scenic Rivers Council 2012, USDA Forest Service 2007a). A 20.0 mile segment, portions of which are in the Sturgeon River Gorge Wilderness, has been classified as Wild, and the management plan states that it will be managed consistent with the wilderness values (USDA Forest Service 2007a). An 8.0 mile segment is designated as Scenic (Interagency Wild & Scenic Rivers Council 2012, USDA Forest Service 2007a).
- The Sturgeon River Gorge Wilderness area is located in the Keweenaw Peninsula/ Sturgeon regional unit. Wilderness is defined as Congressionally designated public land, managed in accordance with the Wilderness Act of 1964. Natural processes are given first priority, and management is undertaken in a way that human use does not substantially change nature (USDA Forest Service 2007b)
- Under the Michigan Water Quality Standards (WQS), portions of the Sturgeon River (Baraga and Houghton Counties) are designated as Outstanding State Resource Waters (OSRW). Also under the Michigan WQS, all surface waters of the Lake Superior basin that are not identified as OSRWs are designated as Lake Superior basin - Outstanding International Resource Waters (LSB-OIRW). Under the above designations, additional anti-degradation controls are applied for new or increased pollutant loadings (Michigan DEQ 2013a).
- The State of Michigan has identified exceptional areas of fish and wildlife habitat along its coastline, connecting waterways, and rivermouths. Designated as Environmental Areas (EAs), certain uses within these areas require state review and approval (Michigan DEQ 2013b). One of these EAs is located in the Township of Torch Lake in Houghton County, in the Keweenaw Peninsula /Sturgeon regional unit (Michigan DEQ 2013c, 2013d).
- In this region, several individual watersheds have state-approved watershed management plans, including Eagle River, Otter River, Pilgrim River and Trap Rock River. These plans serve as guides for communities to protect and improve water quality (M. Preisser, pers. comm., May 31 2013).
- The Houghton Keweenaw Conservation District (HKCD) recently expanded the Bete Grise Preserve. HKCD acquired 1,493 acres of property, including high-quality wetlands and approximately 3,300 feet of shoreline frontage on Lac La Belle along the southeastern coast of the Keweenaw. The property will be managed for conservation and passive recreation (M. Preisser, pers. comm., May 31 2013).
- The Keweenaw Invasive Species Management Area facilitates cooperation and education among federal, state, tribal, local groups and landowners in prevention and management of invasive


species across land ownership boundaries. It spans Houghton, Keweenaw and Baraga Counties, including the Ottawa National Forest (M. Preisser, pers. comm., May 31 2013).

- The Isle Royale & Keweenaw Parks Association is a non-profit educational organization that works with the National Park Service to promote public understanding of Isle Royale National Park and Keweenaw National Historical Park. The group also works to fund research projects (Isle Royale & Keweenaw Parks Association 2010).
- A number of State Important Bird Areas (IBAs) are located in the Keweenaw Peninsula/ Sturgeon regional unit. These IBAs are Sturgeon River Sloughs Wildlife Area IBA, Winegar Moraine and Iron Co. Northern Goshawk IBA, Peshekee Highlands LTA IBA, Brockway Mountain Hawk Watch IBA and Baraga Plains IBA (National Audubon Society 2013, 2012).
- The Keweenaw Land Trust has protected several coastal areas, including property on Manitou Island.

TABLE 16.3: Keweenaw Peninsula/ Sturgeon IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
MI-003	Site	Canyon Falls	Rare plant habitat, geographical feature
MI-006	Site	Silver Mountain	Rare plant habitat, geographical feature
MI-007	Site	Agate Harbor	Rare plant habitat
MI-009	Site	Cat Harbor	Hardwood-conifer swamp, rare plant habitat
MI-010	Site	Cliff Mine	Rare plant habitat
MI-013	Site	Estivant Pines	Dry-mesic northern forest
MI-014	Site	Fort Wilkins	Rare plant and animal habitats
MI-016	Site	Mount Brockway	Rare plant habitat
MI-040	Area	Portage River	Great Lakes marsh, geologic features, rare animal habitat
MI-041	Area	Keweenaw Peninsula	Rare plant and animal habitat geologic features, high biodiversity
MI-048	Area	Bete Grise	

Figure 16.4: Keweenaw Peninsula/ Sturgeon - Important Habitat Sites and Areas



17. Dead-Kelsey

HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	C	COASTAL WETLANDS	B
EMBAYMENTS & INSHORE	C	COASTAL TERRESTRIAL	A
TRIBUTARIES & WATERSHEDS	C	OVERALL B	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



The Huron Mountains are a dominant feature in the Dead/Kelsey watersheds. Photo credit: NOAA.

Summary/ Description

The Dead/Kelsey region (HUC 04020105) extends from Jacobsville in the west to just beyond Marquette in the east. This regional unit is 3,151 km² in size, including the associated nearshore waters. A portion of the Keweenaw Bay Indian Community is located in this regional unit, while the rest of the community is in the Keweenaw Peninsula/ Sturgeon regional unit. regional unit is part of Subregion 0402 – Southern Lake Superior – Lake Superior. The Huron Mountains are located within this region. Mount Arvon, the highest natural point in Michigan is part of the Huron Mountains. The Dead/ Kelsey regional unit contains one tertiary (HUC 8) watershed, Dead-Kelsey, and 6 quaternary (HUC 10) watersheds. The watershed in this unit are dominated by forests. Coastal habitats include rocky shore, sand beaches and coastal wetlands. This region contains some of the most extensive coastal wetlands in Lake Superior.

TABLE 17.1: Dead-Kelsey BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
Agriculture	40.89	1.30	1,287.56	
Developed	11.67	0.37	351.99	
Forest	1,861.73	59.08	96,789.30	
Associated Nearshore Waters	1,014.80	32.20	15,951.22	
Other	187.70	5.96	7,452.78	
Water (inland)	34.54	1.10	8,283.49	
Total Area	3,151.32	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
Coastline (km)	240.5	NA	4.23	Based on SOLEC shoreline
Sand Beaches (km)	18.60	7.73	2.95*	*% of Lake Superior Total Sand Beaches
Coastal Wetlands (km ²)	94.02	26.54*	9.55**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
Natural Cover in Coastal Zone	296.72	83.76*	5.37**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
Number of Islands	47	NA	1.8	
Condition	Region	Region %	% of Lake Superior Total	
Population Density (persons/km ²)	19.31	NA		
Road Density (km/km ²)	0.74	NA		
Number of Dams and Barriers	1,262	NA	5.3	
Artificial Shoreline (km)	15.38	5.18	6.93	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
Private	1,804.02	84.03	2,146.79	Regional area based on landmass
Public/Crown	124.87	5.82	2,146.79	
Tribes/ First Nations	247.86	11.55	2,146.79	
Parks & Protected Areas (total)	7.24	0.34	2,146.79	
Parks & Protected Areas (coast)	0.82	0.23*	354.24**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- The Dead-Kelsey regional unit contains a number of sites of Important Habitat for lake trout and lake whitefish. The Important Habitat sites for both species are found in Keweenaw Bay (L'Anse Bay, Pequaming Bay, Sand Bay) as well as on the eastern side of Point Abbaye and along the shoreline to Marquette (Lake Superior Binational Program Habitat Committee 2006) (Figure 17.1).
- Keweenaw Bay and Huron Bay are noted as Lake Superior embayments which are important for lake sturgeon (Auer 2003). In the Dead-Kelsey regional unit these embayments and the nearshore zone, which provides corridors for movement, are identified as critical management areas for lake sturgeon in the Lake Superior basin (Auer 2003).

Coastal Zone and Islands

- This region contains Important Habitat Areas and Important Habitat Sites (Lake Superior Binational Program Habitat Committee 2006) (Table 17.3, Figure 17.3).
- This region contains some of the most extensive coastal wetlands in Lake Superior.

Tributaries and Watersheds

Figure 17.1: Dead-Kelsey - Coastal and Watershed Features




TABLE 17.2: Dead-Kelsey CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	
Nearshore ¹	C (0.52)	
Embayments and Inshore ^{1,2}	C (0.47)	
Coastal Wetlands ^{2,3}	B (0.632)	
Islands ⁴	A	
Coastal Terrestrial ³	A (0.955)	
Tributaries and Watersheds ²	C (0.42)	

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 17.2: Dead-Kelsey - Condition



Important Issues & Threats

- The ports of Two Harbors, Minnesota, Superior, Wisconsin-Duluth, Minnesota and Presque Isle-Marquette, Michigan have been identified as sites at high risk for invasion by aquatic invasive species, due to ballast water from laker traffic (Rup et al. 2010 as cited in International Joint Commission (IJC) Work Group on Aquatic Invasive Species Rapid Response 2011).
- Mining activities have impacted Lake Superior sediments in the Keweenaw Region; in the Keweenaw Peninsula area, the sediments contain copper enrichments (Kerfoot et al. 2009). Three large copper mills operated in the Keweenaw Bay area; two were located at Gay, Michigan, and one was north of Baraga, near Assinins, Michigan (Kerfoot et al. 2012). The Mass Mill discharged 2.7 teragrams (Tg) of stamp sands, equal to approximately 11% of the total stamp sands discharged into the Keweenaw Bay area (Kerfoot et al. 2012). Investigations have revealed a metal-rich area around the Keweenaw Peninsula, and in the Gay area the copper tailings have drifted from their original sources (Kerfoot et al. 2012). Stamp sands in the Dead-Kelsey region are only found in western lower Keweenaw Bay, north of Baraga (W. Taft, pers. comm., February 25, 2013), which is consistent with the location of the Mass Mill operation.
- The Eagle Mine, a nickel and copper mining operation, is located near the headwaters of the Salmon-Trout River.
- The Deer Lake Area of Concern (AOC) is located in the Dead-Kelsey regional unit, in Marquette County. Two known industrial sources of mercury and untreated wastewater contributed to negative environmental impacts in the AOC. The three beneficial use impairments identified in this AOC are Restrictions on Fish and Wildlife Consumption, Bird or Animal Deformities or Reproductive Problems, and Eutrophication or Undesirable Algae. Two of these BUIs have been removed (See Existing Programs and Projects section) (U.S. EPA 2013s, Michigan DEQ 2013).
- Forest fragmentation through real estate development is an emerging concern in the Upper Peninsula Lake Superior watersheds. Large tracts of forest lands owned by corporate land holders are being sold to companies which run real estate investment trusts; smaller parcels are then developed (W. Taft, pers. comm., February 25, 2013).

Conservation In Action

Parks & Protected Areas

- Ottawa National Forest
- Baraga Plains State Waterfowl Management Area
- Baraga State Park
- Little Presque Isle Natural Area
- Rocking Chair Lakes Natural Area
- State Forest – Baraga & Gwinn Management Units
- Sugarloaf Mountain Natural Area
- Huron Mountain Club (Private)
- McCormick Wilderness Area

Existing Programs & Projects


- Under the Michigan Water Quality Standards (WQS), portions of the Yellow Dog River (Marquette County) are designated as Outstanding State Resource Waters (OSRW). Surface waters of the Lake Superior basin that are not identified as OSRWs are designated as Lake Superior basin - Outstanding International Resource Waters (LSB-OIRW). Under the above designations, additional anti-degradation controls are applied for new or increased pollutant loadings (Michigan DEQ 2013a).

- A four mile segment of the Yellow Dog River from its source at the Bulldog Lake Dam to the Ottawa National Forest boundary is designated as Wild through the National Wild and Scenic Rivers System (Interagency Wild & Scenic Rivers Council 2012, USDA Forest Service 2007a). This four mile section is entirely within the McCormick Wilderness, and the management plan states that it will be managed consistent with the wilderness values (USDA Forest Service 2007a).
- The State of Michigan has identified exceptional areas of fish and wildlife habitat along its coastline, connecting waterways, and rivermouths. Designated as Environmental Areas (EAs), certain uses within these areas require state review and approval (Michigan DEQ 2013b). Five of these EAs are located in Baraga and Marquette counties, in the Dead-Kelsey regional unit. The three Baraga County EAs are located in Arvon and L'Anse Townships, and the two Marquette County EAs are located in Powell Township (Michigan DEQ 2013c, 2013d).
- A number of State Important Bird Areas (IBAs) are located in the Dead-Kelsey regional unit. These IBAs are Yellow Dog Plains IBA, Peshekee Highlands LTA IBA and Baraga Plains IBA (National Audubon Society 2013, 2012).
- A significant amount of work has been done in the Deer Lake Area of Concern (AOC), such that fish have seen a decline in mercury levels and there are no longer regular algal blooms on Deer Lake (S. Swart, pers. comm., March 6 2013). Two of the original three Beneficial Use Impairments (BUIs), reproductive problems and eutrophication, have been addressed and were removed in September 2011 (Michigan DEQ 2013). To address the third BUI, restrictions on fish and wildlife consumption due to mercury contamination, a Great Lakes Restoration Initiative project with the City of Ishpeming is diverting Partridge Creek from old mine workings below the city and back into the storm water system (City of Ishpeming 2011). This project is the last management action for the Area of Concern and has the potential to remove 21% of the mercury flowing into Deer Lake as well as create new habitat areas for brown trout, which used to inhabit Partridge Creek (S. Swart, pers. comm., March 6 2013).
- In this region, several individual watersheds have state-approved watershed management plans, including the Lower Dead River, Partridge Creek, Salmon Trout River, and the Whetstone Brook and Orianna Creek watershed. These plans serve as guides for communities to protect and improve water quality (M. Preisser, pers. comm., May 31 2013).
- Two Cooperative Weed Management Areas (CWMA), the Keweenaw Invasive Species Management Area and the Central Upper Peninsula CWMA, cover the counties of this region, including the Ottawa and Hiawatha National Forests. These groups facilitate cooperation and education among federal, state, tribal, local groups and landowners in prevention and management of invasive species (M. Preisser, pers. comm., May 31 2013).
- The private Huron Mountain Club owns 13,000 acres of lands in the Huron Mountains. Approximately 10,000 acres of these lands are old-growth forest, and several lakes are also located in the land holdings. The club has a limited membership of 50 partners, and in 1938 Aldo Leopold created a preservation plan for the area (Wikipedia contributors No Date).
- The Huron Mountain Wildlife Foundation research center is located in the Huron Mountains and supports the scientific study of the Lake Superior region ecosystems. Established in 1955, sponsorship by this group has produced over 200 publications, including theses and peer-reviewed publications. Approved researchers are able to access the private Huron Mountain Club reserve (Huron Mountain Wildlife Foundation No date).

TABLE 17.3: Dead-Kelsey IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
MI-027	Site	Mulligan Creek	Rare plant habitat
MI-035	Site	Presque Isle	Bedrock beach, dry-mesic northern forest, rare plant habitat, geographical features
MI-036	Site	McCormick	Mesic northern forest, rare plant habitat
MI-037	Area	Lake Independence	Great Lakes marsh, geologic features, rare animal habitat, high biodiversity
MI-038	Area	Huron Mountain	Rare plant and animal habitat, mesic northern forest, dry-mesic northern forest, bedrock glade
MI-039	Site	Pequaming Marsh	Great Lakes marsh, rare animal habitat, geologic feature
MI-049	Site	Huron National Wildlife Refuge	The refuge was established for the protection of migratory birds, specifically, a large nesting colony of herring gulls.

Figure 17.3: Dead-Kelsey - Important Habitat Sites and Areas



18. Betsy-Chocolay

HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	C	COASTAL WETLANDS	B
EMBAYMENTS & INSHORE	C	COASTAL TERRESTRIAL	A-
TRIBUTARIES & WATERSHEDS	C	OVERALL B	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



The Grand Sable Dunes are part of the longest sand beach on Lake Superior. Photo credit: National Park Service photo by Gregg Bruff.

Summary/ Description

The Betsy-Chocolay region (HUC 04020201) extends from near Marquette in the west to Emerson in the east, south of Whitefish Point. The regional unit is 5,031.48 km² in size, including the associated nearshore waters. It is part of Subregion 0402 – Southern Lake Superior-Lake Superior.

The Betsy-Chocolay regional unit contains one tertiary (HUC 8) watershed, Betsy-Chocolay, and 4 quaternary (HUC 10) watersheds. The watersheds are within a forest dominated landscape. The 300 km coast includes over 20% of all the sand beached found in Lake Superior. Other coastal habitats include rocky shores, sandstone cliffs and coastal wetlands. Extensive coastal wetland occur at Whitefish Point.

TABLE 18.1: Betsy-Chocolay BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
Agriculture	21.86	0.43	1,287.56	
Developed	4.07	0.08	351.99	
Forest	2,382.81	47.36	96,789.30	
Associated Nearshore Waters	2,288.31	45.48	15,951.22	
Other	290.34	5.77	7,452.78	
Water (inland)	44.09	0.88	8,283.49	
Total Area	5,031.48	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
Coastline (km)	299.4	NA	5.26	Based on SOLEC shoreline
Sand Beaches (km)	137.50	45.93	21.84*	*% of Lake Superior Total Sand Beaches
Coastal Wetlands (km ²)	205.66	40.79*	20.89**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
Natural Cover in Coastal Zone	470.31	93.27*	8.52**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
Number of Islands	4	NA	0.2	
Condition	Region	Region %	% of Lake Superior Total	
Population Density (persons/km ²)	4.85	NA		
Road Density (km/km ²)	0.61	NA		
Number of Dams and Barriers	846	NA	3.6	
Artificial Shoreline (km)	11.10	2.36	5.00	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
Private	1,250.17	45.58	2,742.82	Regional area based on landmass
Public/Crown	1,165.97	42.51	2,742.82	
Tribes/ First Nations	0.00	0.00	2,742.82	
Parks & Protected Areas (total)	328.13	11.96	2,742.82	
Parks & Protected Areas (coast)	112.17	22.25*	504.23**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- The Betsy-Chocolay regional unit contains numerous sites of Important Habitat for lake trout and lake whitefish. The Important Habitat sites for both species are found in at various points along the shore, and with a slight concentration of Important Habitat sites for both species in the area of Grand Island, Au Train Bay, and South Bay, near Munising Michigan, and Pictured Rocks National Lakeshore (Lake Superior Binational Program Habitat Committee 2006) (Figure 18.1).
- Munising Bay is noted as a Lake Superior embayment important for lake sturgeon (Auer 2003). In the Betsy-Chocolay regional unit this embayment and the nearshore zone, which provides corridors for movement, are identified as critical management areas for lake sturgeon in the Lake Superior basin (Auer 2003).

Coastal Zone and Islands

- This region includes some of the most extensive sand beaches on Lake Superior, including the Grand Sable Dunes.

Tributaries and Watersheds

- The Two Hearted watershed covers approximately 269,154 acres (108,922 hectares) (Chippewa/East Mackinac Conservation District 2008).
- The Betsy-Chocolay regional unit contains many Important Habitat Areas and Important Habitat Sites (Lake Superior Binational Program Habitat Committee 2006) (Table 18.3, Figure 18.3).

LAKE SUPERIOR BIODIVERSITY CONSERVATION ASSESSMENT

18. Betsy-Chocolay

Coastal and Watershed Features

LEGEND

Coastal and Aquatic Habitats

- Lake Trout Important Habitat
- Whitefish Important Habitat
- Other Important Fish Habitat
- Lake Trout Spawning Reefs
- Whitefish Spawning Areas
- Sand Beach
- Coarse Beach
- Rocky Shore / Bluff
- Coastal Wetland (2km)

Land Cover

- Wetland
- Natural
- Agricultural
- Developed
- Unknown / Cloud

Other Features

- Road Network
- HUC10 Watershed Boundary
- Coastal Terrestrial Zone
- Unit Boundary
- Inshore Zone / Lake / River
- Nearshore Zone
- Open Water

Scale: 5,000 0 5,000 m / 20,000 0 20,000 ft

PROJECTION: North American Lambert Conformal Conic

TABLE 18.2: Betsy-Chocolay CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	
Nearshore ¹	C (0.56)	
Embayments and Inshore ^{1,2}	C (0.54)	
Coastal Wetlands ^{2,3}	B (0.673)	
Islands ⁴	A	
Coastal Terrestrial ³	A- (0.938)	
Tributaries and Watersheds ²	C (0.52)	

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 18.2: Betsy-Chocolay - Condition



Important Issues & Threats

- The Duck Lake Fire of 2012 burned approximately 21,069 acres in the Betsy-Chocoday regional unit, in Luce County (Michigan DNR 2012).
- The presence of Emerald Ash Borer (EAB) has been detected in counties in the Betsy-Chocoday regional unit. The area is now under a number of quarantine measures, including Federal EAB quarantine and Michigan state quarantine (Cooperative Emerald Ash Borer Project 2013).
- An outbreak of Beech Bark Disease has occurred in the Betsy/ Chocoday regional unit (W. Taft, pers. comm., February 25, 2013). Beech scale and beech bark disease were first discovered in Michigan in 2000; affected beech trees were first found in the northwestern Lower Peninsula and the eastern Upper Peninsula (McCullough, Heyd and O'Brien 2002).
- An Emergency Prevention and Response Plan for Viral Hemorrhagic Septicemia has been developed for Isle Royale National Park, Pictured Rocks National Lakeshore, Apostle Islands National Lakeshore and the Grand Portage Band of the Lake Superior Chippewa Reservation (within which is the Grand Portage National Monument) (NPS 2013a).
- Forest fragmentation through real estate development is an emerging concern in the Upper Peninsula Lake Superior watersheds. Large tracts of forest lands owned by corporate land holders are being sold to companies which run real estate investment trusts; smaller parcels are then developed (W. Taft, pers. comm., February 25, 2013).

Conservation In Action

Parks & Protected Areas

- Pictured Rocks National Lakeshore
- Au Train Basin State Waterfowl (Wildlife) Management Area
- Cusino State Wildlife Research Area
- Deer Park Natural Area
- Hiawatha National Forest
- Laughing Whitefish Falls Natural Area
- McMahon Lake Strangmoor Natural Area
- Muskallonge Lake State Park
- State Forest – Gwinn, Newberry & Shingleton Management Units
- Tahquamenon Falls State Park
- Wagner Falls State Park

Existing Programs & Projects

- The Two Hearted River is a Michigan Natural River System, a designation given to the finest river systems in the state (Michigan DNR 2013a). It is the only one of Michigan's 16 Natural River Systems that flows to Lake Superior.
- Under the Michigan Water Quality Standards (WQS), portions of the Two-Hearted River and Dawson Creek (Luce County) and all waterbodies within the designated boundary of Pictured Rocks National Lakeshore are designated as Outstanding State Resource Waters (OSRW). Also under the Michigan WQS, all surface waters of the Lake Superior basin that are not identified as OSRWs are designated as Lake Superior basin - Outstanding International Resource Waters (LSB-OIRW). Under the above designations, additional anti-degradation controls are applied for new or increased pollutant loadings (Michigan DEQ 2013a).


Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

- The State of Michigan has identified exceptional areas of fish and wildlife habitat along its coastline, connecting waterways, and rivermouths. Designated as Environmental Areas (EAs), certain uses within these areas require state review and approval (Michigan DEQ 2013b). Two of these EAs are located in Grand Island Township in Alger County, in the Betsy-Chocolay regional unit (Michigan DEQ 2013c, 2013d).
- A number of State Important Bird Areas (IBAs) are located in the Betsy-Chocolay regional unit. These IBAs are Sleeper Lake burn IBA, Grand Marais to Whitefish Point beaches (Piping Plover) IBA and Tahquamenon Falls State Park and Munising Moraine IV LTA IBA (National Audubon Society 2013, 2012).
- The Whitefish Point Bird Observatory is located in the Betsy/ Chocolay regional unit. Whitefish Point is a concentration area for migrating birds, and the Whitefish Point Bird Observatory has been monitoring and documenting annual spring and fall migrations for over 30 years (Whitefish Point Bird Observatory No date).
- In this region, several individual watersheds have state-approved watershed management plans, including the Chocolay River, Two Hearted River, and the Munising Bay watershed. These plans serve as guides for communities to protect and improve water quality (M. Preisser, pers. comm., May 31 2013).
- The Michigan Department of Natural Resources maintains a fish hatchery along Cherry Creek, just south of Harvey, Michigan on the far west end of this region. It is the primary broodstock and rearing facility for brook and lake trout that are used in both inland and Great Lakes waters. The facility also rears splake (a brook trout - lake trout hybrid) for both Great Lakes and inland waters (M. Preisser, pers. comm., May 31 2013).
- Two Cooperative Weed Management Areas (CWMA), the Central Upper Peninsula CWMA and the Eastern Upper Peninsula CWMA, cover the counties of this region, including the Hiawatha National Forest. These groups facilitate cooperation and education among federal, state, tribal, local groups and landowners in prevention and management of invasive species (M. Preisser, pers. comm., May 31 2013).

TABLE 18.3: Betsy-Chocolay IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
MI-001	Site	Laughing Whitefish Falls/ Whitefish National Wildlife Reserve	Mesic Northern Forest, rare plant habitat
MI-001	Area	Laughing Whitefish Falls/ Whitefish National Wildlife Reserve	Mesic Northern Forest, rare plant habitat
MI-002	Area	Rock River Canyon Wilderness Area	Moist non-acid cliff, mesic northern forest, geographical features, rare plant and animal habitats
MI-004	Site	Whitefish Point	Rare plant and animal habitat
MI-017	Site	Barclay Lake	Bog, dry northern forest, rare plant habitat
MI-018	Site	Barfield Lakes	Bog, dry northern forest, rich conifer swamp, mesic northern forest
MI-019	Site	Beavertown Lakes	Dry northern forest, hardwood-conifer swamp, rich conifer swamp, muskeg, dry-mesic northern forest
MI-020	Site	Blind Sucker River	Rare plant and animal habitat
MI-021	Site	Crisp Point	Rare plant and animal habitat
MI-022	Site	Deer Park	Rare plant and animal habitat
MI-023	Site	Little Lake	Rare plant and animal habitat
MI-024	Site	McMahon Lake	Patterned fen, rare plant habitat
MI-025	Site	Swamp Lakes	Intermittent wetland, dry-mesic northern forest, rare plant habitat
MI-026	Site	Dukes RNA	Rich conifer swamp, mesic northern forest
MI-029	Area	Vermilion	Lake Superior beach community, rare plant and animal habitat
MI-030	Area	Two-Hearted River	Representative landscape complex, old growth red/white pine forest, old growth cedar forest, hemlock and white pine forest, rare plant habitat, migrant bird habitat, relatively undisturbed wetland communities, coastal plain marsh, patterned fen, muskeg,
MI-031	Site	Grand Marais	Rare plant and animal habitats
MI-032	Area	Grand Sable Dunes	Perched dunes, open dunes, geologic feature, rare plant habitat
MI-033	Area	Grand Island	Great Lakes marsh, mesic northern forest, rare plant and animal habitat
MI-034	Area	Au Train	Large Great Lakes marsh, fresh water estuary, high biodiversity
MI-046	Area	Pictured Rocks	Rare plant habitat, geomorphic features
MI-047	Area	Tahquamenon Falls State Park	Representative natural plant communities

Figure 18.3: Betsy-Chocolay - Important Habitat Sites and Areas



19. Waiska/ Tahquamenon

HEALTHY WATERS REPORT CARD

OFFSHORE	NA	ISLANDS	A
NEARSHORE	C	COASTAL WETLANDS	C
EMBAYMENTS & INSHORE	C	COASTAL TERRESTRIAL	A
TRIBUTARIES & WATERSHEDS	C	OVERALL B-	

Report card denotes general condition/health of each biodiversity target in the region based on condition/stress indices. See introduction to the regional summaries.



A Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>



The mouth of the Tahquamenon River. Photo provided by William Taft. Photo credit: David Kenyon/ Michigan Department of Natural Resources

Summary/ Description

The Waiska/ Tahquamenon region (HUC 04020202, HUC 04020203 and HUC 04070001) extends from Emerson to Sault Ste Marie and the Michigan/Ontario border at the St. Marys River. It is 2,939.00 km² in size, including the associated nearshore waters. It is part of Subregion 0402 – Southern Lake Superior-Lake Superior. The Bay Mills Indian Community is located in this regional unit. The reservation of the Sault tribe of Chippewa Indians is also located in this regional unit. These watersheds are dominated by forest cover. Coastal habitats include sand beaches and coastal wetlands, with scattered rocky shores.

TABLE 19.1: Waиска/ Tahquamenon BY THE NUMBERS

Land and Water Cover	Region (km²)	Region %	Lake Superior Total (km²)	Notes
Agriculture	38.23	1.30	1,287.56	
Developed	2.80	0.10	351.99	
Forest	2,285.52	77.77	96,789.30	
Associated Nearshore Waters	390.0	13.27	15,951.22	
Other	206.22	7.02	7,452.78	
Water (inland)	16.21	0.55	8,283.49	
Total Area	2,939.00	100	130,116.37	
Coastal Features	Region	Region %	% of Lake Superior Total for Coastal Feature	
Coastline (km)	113.3	NA	1.99	Based on SOLEC shoreline
Sand Beaches (km)	38.60	34.07	6.13*	*% of Lake Superior Total Sand Beaches
Coastal Wetlands (km ²)	79.91	53.82*	8.12**	*% of Regional Coastal Area ** % of Lake Superior Total Coastal Wetlands
Natural Cover in Coastal Zone	139.46	93.94*	2.53**	*% of Regional Coastal Area ** % of Lake Superior Total Natural Cover in Coastal Area
Number of Islands	16	NA	0.6	
Condition	Region	Region %	% of Lake Superior Total	
Population Density (persons/km ²)	4.89	NA		
Road Density (km/km ²)	0.60	NA		
Number of Dams and Barriers	305	NA	1.3	
Artificial Shoreline (km)	27.91	20.01	12.57	
Land Ownership & Protection	Region (km²)	Region %	Regional Area (km²)	
Private	1,137.55	44.63	2,548.88	Regional area based on landmass
Public/Crown	1,316.43	51.65	2,548.88	
Tribes/ First Nations	5.83	0.23	2,548.88	
Parks & Protected Areas (total)	89.17	3.50	2,548.88	
Parks & Protected Areas (coast)	12.25	8.25*	148.47**	*% of Regional Coastal Area **Regional Coastal Area (km ²)

Important Biodiversity Features

Nearshore and Inshore Waters

- The Waиска/ Tahquamenon regional unit contains sites of Important Habitat for lake trout and lake whitefish (Lake Superior Binational Program Habitat Committee 2006) (Figure 19.1).
- Whitefish Bay is noted as a Lake Superior embayment important for lake sturgeon (Auer 2003). In the Waиска/ Tahquamenon regional unit this embayment and the nearshore zone, which provides corridors for movement, are identified as critical management areas for lake sturgeon in the Lake Superior basin (Auer 2003).

Coastal Zone and Islands

- This region has extensive coastal wetlands in Whitefish Bay and the mouth of the St. Marys River.

Tributaries and Watersheds

- Historically 21 tributaries in Lake Superior had lake sturgeon spawning runs. The Tahquamenon River is one of these historical spawning tributaries, and is located in the Waiska/ Tahquamenon regional unit. The Tahquamenon River population status is extirpated (Golder Associates Ltd. 2011).
- A Lake Sturgeon Rehabilitation Plan for Lake Superior (Auer 2003) identifies the Tahquamenon River as one of the seventeen tributaries to Lake Superior in which there should be a focus on lake sturgeon rehabilitation.
- This regional unit contains two Important Habitat Areas and one Important Habitat Site (Lake Superior Binational Program Habitat Committee 2006) (Table 19.3, Figure 19.3).

Figure 19.1: Waika/ Tahquamenon - Coastal and Watershed Features




TABLE 19.2: Waiska/ Tahquamenon CONDITION AND TRENDS

Target (Data Source)	Condition	Trends
Offshore ¹	NA	
Nearshore ¹	C (0.41)	
Embayments and Inshore ^{1,2}	C (0.47)	
Coastal Wetlands ^{2,3}	C (0.590)	
Islands ⁴	A	
Coastal Terrestrial ³	A (0.840)	
Tributaries and Watersheds ²	C (0.52)	

A: Very Good	<i>Ecologically desirable status; requires little intervention for maintenance</i>
B: Good	<i>Within acceptable range of variation; may require some intervention for maintenance.</i>
C: Fair	<i>Outside of the range of acceptable variation and requires management. If unchecked, the biodiversity target may be vulnerable to serious degradation.</i>
D: Poor	<i>Allowing the biodiversity target to remain in this condition for an extended period will make restoration or preventing extirpation practically impossible.</i>
Unknown	<i>Insufficient information.</i>


1: Great Lakes Cumulative Stress (GLEAM 2012, Allan et al. 2013)

2: Watershed Stress Index (GLEI 2013)

3: Coastal Condition Index (developed for this report)

4 : Island Condition Score (Henson et al. 2010)

Figure 19.2: Waika/ Tahquamenon - Condition



Important Issues & Threats

- The presence of Emerald Ash Borer (EAB) has been detected in counties in the Waiska/ Tahquamenon regional unit. The area is now under a number of quarantine measures, including Federal EAB quarantine and Michigan state quarantine (Cooperative Emerald Ash Borer Project 2013).
- Forest fragmentation through real estate development is an emerging concern in the Upper Peninsula Lake Superior watersheds. Large tracts of forest lands owned by corporate land holders are being sold to companies which run real estate investment trusts; smaller parcels are then developed (W. Taft, pers. comm., February 25, 2013).
- The St. Marys River Area of Concern (AOC) is located between two regional units, the Goulais and Waiska/ Tahquamenon regional units. A number of point and nonpoint sources have contributed to beneficial use impairments, and sediments are contaminated with arsenic, cadmium, chromium, copper, cyanide and lead. Ten beneficial use impairments were identified in the St. Marys River AOC (U.S. EPA 2013a).
- The Cannelton Industries Inc. Superfund site is located just inside the boundaries of the Waiska/ Tahquamenon regional unit. This site consists of 75 acres along the St. Marys River. Soils, sediments and surface water in the river were contaminated with heavy metals from tannery operations and general wastes. A number of cleanup activities have been undertaken and the EPA plans to remove the site from the National Priorities List within the next several years. A restrictive covenant and easement for the property was signed in March 2013, to ensure the continued management of contaminated materials on the property (U.S. EPA 2013q).

Conservation In Action

Parks & Protected Areas

- Brimley State Park
- Hiawatha National Forest
- State Forest – Sault Ste. Marie Management Unit
- Tahquamenon Falls State Park

Existing Programs & Projects


- Segments of the East Branch of the Tahquamenon River have been designated as portions of the National Wild and Scenic Rivers System (Interagency Wild & Scenic Rivers Council 2012)
- Under the Michigan Water Quality Standards (WQS), portions of the Tahquamenon River (Chippewa County) are designated as Outstanding State Resource Waters (OSRW). Also under the Michigan WQS, all surface waters of the Lake Superior basin that are not identified as OSRWs are designated as Lake Superior basin - Outstanding International Resource Waters (LSB-OIRW). Under the above designations, additional anti-degradation controls are applied for new or increased pollutant loadings (Michigan DEQ 2013a).
- The State of Michigan has identified exceptional areas of fish and wildlife habitat along its coastline, connecting waterways, and rivermouths. Designated as Environmental Areas (EAs), certain uses within these areas require state review and approval (Michigan DEQ 2013b). One EA is located in the Lake Superior waters of Chippewa County (at Tahquamenon Island), in the Waiska/ Tahquamenon regional unit (Michigan DEQ 2013c, 2013d).
- The Kirtland's Warbler Management Units & Guide's Rest IBA is a Global Important Bird Area located in the Waiska/ Tahquamenon regional unit (National Audubon Society 2013, 2012).

- A number of State Important Bird Areas (IBAs) are located in the Waiska/ Tahquamenon regional unit. These IBAs are Sault Sainte Marie Clay Plain IBA, Sleeper Lake burn IBA, Raco Plains & Wilwin Wetland IBA, Dollarville Flooding IBA and Tahquamenon Falls State Park and Munising Moraine IV LTA IBA (National Audubon Society 2013, 2012).
- The Eastern Upper Peninsula Cooperative Weed Management Area is a partnership of city, county, state, federal, and tribal officials who have joined together with local citizens, landowners, and not-for-profit groups to share invasive plant management resources. Their goal is to facilitate cooperation and coordination networking across jurisdictional boundaries (M. Preisser, pers. comm., May 31 2013).

TABLE 19.3: Waiska/ Tahquamenon IMPORTANT HABITAT SITES AND AREAS

<i>Code</i>	<i>Site/ Area</i>	<i>Important Habitat Site/Area Name</i>	<i>Key Features</i>
MI-028	Site	Tahquamenon Bay	Rare plant habitat, rare animal habitat, wooded dune and swale complex
MI-045	Area	Delirium Wilderness Area	Representative natural plant communities
MI-047	Area	Tahquamenon Falls State Park	Representative natural plant communities
ON-128	Area	St. Mary's River	Fish spawning area.


Figure 19.3: Waiska/ Tahquamenon - Important Habitat Sites and Areas



20. Lake Superior Open Waters

Summary/ Description


The final regional unit includes all of the open, offshore waters of Lake Superior. Classifying the open waters as a single unit was recommended by the Aquatic Community Committee/Lake Superior Technical Committee.



Important Biodiversity Features


Offshore Waters

- The historic offshore fish community was a coldwater fish community dominated by lake trout, lake whitefish and cisco (OMNR 2013d).
- A detailed description of these offshore waters can be found in Volume One of this report in the biodiversity target descriptions.




Lake Superior sunset. Taken in the Black Presque Isle Ontonagon regional unit. Photo provided by William Taft. Photo credit: Mat Wesener/ Michigan Department of Environmental Quality.

Figure 20.1: Open Waters - Coastal and Watershed Features



Note: Detailed information on the features and condition of nearshore and inshore waters are included in the other regional summaries.

Figure 20.2: Open Waters – Condition



Important Issues & Threats

- The rehabilitation of lean lake trout in nearshore waters of the east end of the lake has not progressed to the same extent as the remainder of the lake. The Ontario Ministry of Natural Resources (OMNR) continues to make efforts to negotiate an agreement with First Nation communities in order to reduce harvests. In the interim, rehabilitation efforts through fish stocking have been discontinued (S. Greenwood, pers. comm., May 27 2013).
- Invasive species and habitat loss have negatively affected the historic fish communities in FMZ 9. The focus of fisheries management in this zone is now rehabilitation of the fish communities (OMNR 2013d).

Conservation In Action


Parks & Protected Areas

- The Lake Superior National Marine Conservation Area (LSNMCA) of Canada extends from the water's edge, beyond the nearshore water boundaries associated with the Little Pic, Jack Pine/ Nipigon and Black Sturgeon regional units, into the offshore waters and to the international boundary with the United States.

Existing Programs & Projects

- Cooperative management of Lake Superior and FMZ 9 is coordinated through the Great Lakes Fisheries Commission. Provincial, state, federal and tribal agencies all play a role in this collaborative management (OMNR 2013d).
- The Canadian waters of Lake Superior are contained within Fisheries Management Zone 9 (FMZ 9), and are managed by the Upper Great Lakes Management Unit of the Ontario Ministry of Natural Resources (OMNR 2013d, M. Chase, pers. comm., June 3 2013). This includes the nearshore and inshore waters, including embayments, and the offshore waters. Fisheries Management Zone 9 also includes all of the islands in the zone, except for St. Ignace Island, Simpson Island and Michipicoten Island (OMNR 2013d). FMZ 9 is managed to “rehabilitate and maintain a diverse, healthy and self-regulating fish community, dominated by indigenous species and supporting sustainable fisheries” (OMNR 2013d: paragraph 4).

Figure 20.3: Open Waters - Important Habitat Sites and Areas



References

- Allan, J.D., McIntyre, P.B., Smith, S.D.P., Halpern, B.S., Boyer, G.L., Buchsbaum, A., ... Steinman, A.D. 2013. Joint analysis of stressors and ecosystem services to enhance restoration effectiveness. *Proceedings of the National Academy of Sciences of the United States of America (PNAS)* 110(1): 372-377. doi:10.1073/pnas.1213841110
- Allen, B. 2013. The Nature Conservancy (TNC), Wisconsin: Clough Island: Heart of the St. Louis River Estuary. Available at <http://www.nature.org/ourinitiatives/regions/northamerica/unitedstates/wisconsin/howwework/clough-island.xml> Accessed 14 January 2013.
- Annells, R.N. 1974. Keweenaw Volcanic Rocks of Michipicoten Island, Lake Superior, Ontario: An Eruptive Centre of Proterozoic Age, Geological Survey of Canada, Bulletin 218, Department of Energy, Mines and Resources Canada.
- Apex Realty Services Ltd. 2013. File 73 – Own a part of the magic of Michipicoten Island. Available at http://www.recreationland.net/recreation_land_for_sale/ontario/Islands_For_Sale/File_73_-_OWN_A_PART_OF_THE_MAGIC_OF_MICHIPICOTEN_ISLAND/index.html Accessed 22 February 2013.
- Auer, N.A. [Ed.]. 2003. A lake sturgeon rehabilitation plan for Lake Superior. Great Lakes Fish. Comm. Misc. Publ. 2003-02. Available at <http://www.glfc.org/pubs/pub.htm> Accessed 10 May 2012.
- Auer, N.A. and E.A. Baker. No Date. Streamside lake sturgeon culture for the Ontonagon River, Michigan. Revised Final Report. Available at <http://www.nero.noaa.gov/StateFedOff/grantfactsheets/MI/FINAL%20REPORTS/FINAL%20NA04NMF4050265.pdf> Accessed 16 January 2013.
- Bad River Band of Lake Superior Tribe of Chippewa Indians. 2011. Water Quality Standards. Available at: <http://badriver-nnsn.gov/natural-resources/announcement-a-alerts-natural-resources> Modified 6 July 2011.
- Bad River Band of Lake Superior Tribe of Chippewa Indians. 2006. Non-Point Source Assessment Report.
- Bad River Watershed Association (BRWA). 2013a. Bad River Watershed Description [Homepage]. Available at <http://www.badriverwatershed.org/index.php/bad-river-watershed> Accessed 12 February 2013.
- Bad River Watershed Association (BRWA). 2013b. Restore: Culvert Restoration Program. Available at <http://www.badriverwatershed.org/index.php/restore> Accessed 17 April 2013.
- Baird & Associates. 2000. Nemadji River Basin Sediment Transport Modeling for Two Subwatersheds. Prepared For: USACE, Detroit District. Available at <http://glc.org/tributary/models/nemadji.html> Accessed 11 February 2013.
- Brookfield Renewable Energy Partners L.P. 2013. Development Projects Available at http://www.brookfieldrenewable.com/content/portfolio/development_projects-30490.html Accessed 21 March 2013.
- Brookfield Renewable Power Inc. No Date. Coldwell Wind Energy Project. Available at <http://www.coldwellwind.com/> Accessed 21 March 2013.
- Canadian Food Inspection Agency (CFIA). 2012. Emerald Ash Borer Infested Places Order. Available at <http://www.inspection.gc.ca/plants/plant-protection/insects/emerald-ash-borer/infested-places-order/eng/1337705116683/1337705207346> Accessed 14 February 2013.
- Canadian Wind Energy Association. 2008. Prince Wind farm. Available at http://www.canwea.ca/featuredWindFarm_e.php?farmId=36 Accessed 21 March 2013.
- Chequamegon Bay Area Partnership (CBAP). Strategic Priorities Mapping a Path to the Future. Available at <http://www.northland.edu/assets/files/SOEI/CBAP.Strategic.Priorities.2013.Final.pdf> Accessed 23 May 2013.
- Chikoski, J. and L. Nyman. 2011. The 2010 Ontario Peregrine Falcon Survey – A Summary Report. Unpublished report, Ontario Ministry of Natural Resources, Thunder Bay. 36 pp. Available at http://www.mnr.gov.on.ca/stdprodconsume/groups/lr/@mnr/@species/documents/document/stdprod_101845.pdf Accessed 20 March 2013.
- Chippewa/East Mackinac Conservation District. 2008. Eastern Upper Peninsula Rapid Watershed Assessment Summary. 16p. Available at <http://www.mi.nrcs.usda.gov/programs/RWA.html> Accessed 21 February 2013.
- City of Duluth. 2013a. Park Point Trail. Available at <http://www.duluthmn.gov/parks/parkpointtrail.cfm> Accessed 22 May 2013.
- City of Duluth. 2013b. Duluth Natural Areas Program. Available at <http://www.duluthmn.gov/planning/dnap/dnap.cfm> Accessed 24 April 2013.
- City of Ishpeming. 2011. Ishpeming's Partridge Creek Diversion. Brochure. Available at http://ishpemingcity.org/download/city_projects/CreekBrochureProof7.FINAL_.pdf Accessed 13 March 2013.
- City of Superior. No date a. Superior Municipal Forest. Available at <http://www.ci.superior.wi.us/index.aspx?NID=224> Accessed 22 May 2013.
- City of Superior. No date b. Wisconsin Point. Available at <http://www.ci.superior.wi.us/index.aspx?NID=226> Accessed 22 May 2013.
- Cooperative Emerald Ash Borer Project. 2013. Cooperative Emerald Ash Borer Project Map. Available at http://www.emeraldashborer.info/files/MultiState_EABpos.pdf Accessed 13 February 2013.

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

- COSEWIC Committee on the Status of Endangered Wildlife in Canada. 2011a. Wildlife species search. Falcon anatum/tundrius, Peregrine | Falco peregrines anatum/tundrius. Available at http://www.cosewic.gc.ca/eng/sct1/searchdetail_e.cfm Updated 11 July 2011. Accessed 21 March 2013.
- COSEWIC Committee on the Status of Endangered Wildlife in Canada. 2011b. Wildlife species search. Caribou, Woodland | Rangifer tarandus caribou | Boreal population. Available at http://www.cosewic.gc.ca/eng/sct1/searchdetail_e.cfm Updated 7 November 2011. Accessed 4 April 2013.
- Eason, G. 2002, 2003. Personal communication. Area Biologist, Wawa/White River, OMNR.
- Ecclestone, A. 2013. Lake Sturgeon Population Characteristics and Habitat Utilization in the White River, Ontario, from 2011 to 2012. Anishinabek/Ontario Fisheries Resource Centre, North Bay, Ontario.
- Ecclestone, A., T. Pratt, T. Whilans and C. Wilson. 2013. Lake Sturgeon Population Characteristics and Movement Patterns in the Pic River, Ontario DRAFT. Anishinabek/Ontario Fisheries Resource Centre, North Bay, Ontario.
- Egan, A. T. 2009. Breeding bird monitoring at Isle Royale National Park, Michigan: 1996-2008. Natural Resource Technical Report NPS/GLKN/NRTR—2009/160. National Park Service, Fort Collins, Colorado. Available at http://www.nps.gov/isro/naturescience/upload/ISRO_BBS_Egan_96_08_NRTR%5B1%5D.pdf Accessed 19 February 2013.
- Environment Canada 2010a. Peninsula Harbour Area of Concern. Available at <http://www.ec.gc.ca/raps-pas/default.asp?lang=En&n=FD4765B0-1&toc=show> Modified 12 January 2010. Accessed 28 June 2013.
- Environment Canada 2010b. Jackfish Bay Area of Concern. Available at <http://www.ec.gc.ca/raps-pas/default.asp?lang=En&n=7B2CB149-1&toc=show> Modified 12 January 2010. Accessed 28 June 2013.
- Friends of Pukaskwa National Park. No date. Homepage. Available at <http://www.friendsofpukaskwa.ca/> Accessed 20 February 2013.
- Gogan, P.J.P. and Cochrane, J.F. 1994. Restoration of Woodland Caribou to the Lake Superior Region. U.S. National Park Service Publications and Papers. Paper 11. Available at <http://digitalcommons.unl.edu/natlpark/11> Accessed 2 April 2013.
- Golder Associates Ltd. 2011. Recovery Strategy for Lake Sturgeon (*Acipenser fulvescens*) – Northwestern Ontario, Great Lakes-Upper St. Lawrence River and Southern Hudson Bay-James Bay populations in Ontario. Ontario Recovery Strategy Series. Prepared for the Ontario Ministry of Natural Resources, Peterborough, Ontario. vii + 77 pp. Available at http://www.mnr.gov.on.ca/stdprodconsume/groups/lr/@mnr/@species/documents/document/stdprod_086034.pdf Accessed 10 May 2013.
- Goodier, J.L. 1982 The Fish and Fisheries of Canadian Lake Superior Inst. for Env. Studies Univ. of Toronto 176 pp.
- Gorman, O.T., S.A. Moore, A.J. Carlson, and H.R. Quinlan. 2008. Nearshore habitat and fish community associations of coaster brook trout in Isle Royale, Lake Superior. Transactions of the American Fisheries Society. 137:1252-1267.
- Grand Portage Band of Lake Superior Chippewa. 2006. Grand Portage Reservation Water Quality Standards. Accessed 23 May 2013.
- Grand Portage. No date. Our Heritage. Available at <http://www.grandportage.com/heritage.php> Accessed 24 April 2013.
- Greenwood S. 2000. Summary Of Fourteen Years Of Chinook Salmon Fall Spawning Index Counts In The Michipicoten River. Upper Great Lakes Management Unit - Lake Superior Quik Report 00 - 01 Ontario Ministry of Natural Resources.
- Henson, B.L., D.T. Kraus, M.J. McMurtry, and D. N. Ewert. 2010. Islands of life: a biodiversity and conservation atlas of the Great Lakes islands. Nature Conservancy of Canada. 154pp. Available at http://nhic.mnr.gov.on.ca/MNR/nhic/projects/Islands_of_Life/Islands_of_Life_Final.pdf Accessed 12 November 2012.
- Hoff, M.H. [ED.] 2002. A rehabilitation plan for walleye populations and habitats in Lake Superior. Great Lakes Fish. Comm. Misc. Pub. 2003-01. Available at http://www.glfc.org/pubs/SpecialPubs/2003_01.pdf Accessed 21 March 2013.
- Huron Mountain Wildlife Foundation. No date. Huron Mountain Wildlife Foundation Home Page. Available at <http://www.hmwf.org/index.php> Accessed 12 May 2013.
- Interagency Wild & Scenic Rivers Council. 2012. River Mileage Classifications for Components of the National Wild and Scenic Rivers System. Available at <http://www.rivers.gov/rivers/publications.php> Accessed 13 February 2013.
- International Joint Commission (IJC) Work Group on Aquatic Invasive Species Rapid Response. 2011. An Assessment of Early Detection Monitoring and Risk Assessments for Aquatic Invasive Species in the Great Lakes-St. Lawrence Basin. Available at <http://meeting.ijc.org/sites/default/files/workgroups/RAandMonitoringJuly29.pdf> Accessed 21 February 2013.
- Isle Royale & Keweenaw Parks Association. 2010. About the Isle Royale & Keweenaw Parks Association. Available at http://www.irkpa.org/the.store?cart_id=1135152.9736&page=/about/index.html Accessed 21 February 2013.
- Jackfish Bay Remedial Action Plan Team. 1998. Jackfish Bay Remedial Action Plan Stage 2: Remedial Strategies for Ecosystem Restoration. Available at <http://www.northshorerap.ca/publications> Accessed 28 June 2013.
- Kerfoot, W.C., F. Yousef, S.A. Green, R. Regis, R. Shuchman, C.N. Brooks, M. Sayers, B. Sabol, and M. Graveset. 2012. Light detection and ranging (LiDAR) and multispectral studies of disturbed Lake Superior coastal environments. Limnology and Oceanography. 57(3): 749-771. DOI: 10.4319/lo.2012.57.3.0749

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

- Kerfoot, W.C., J. Jeong, and J.A. Robbins. 2009. Lake Superior mining and the proposed mercury zero-discharge region. Pages 153-216 in Munawar, M. and I.F. Munawar (Eds.) (2009) State of Lake Superior. Aquatic Ecosystem Health and Management Society, Ecovision World Monograph Series, Canada.
- Lake Superior Binational Program (LSBP) 2008. Lakewide Management Plan (LaMP) 2008. Available at http://www.epa.gov/glnpo/lamp/ls_2008/index.html. Accessed 23 May 2013.
- Lake Superior Binational Program Habitat Committee. 2006. Important Habitat in the Lake Superior Basin. Available at <http://www.nrri.umn.edu/lsgis/importanthabitat.htm> Accessed 5 February 2013.
- LRCA Lakehead Region Conservation Authority. No date. Programs and Services of the Lakehead Region Conservation Authority. Available at <http://www.lakeheadca.com/services.htm> Accessed 21 February 2013.
- Liebermann, R.J. 1998. Comparative phytogeography and conservation of the Caribou Islands, Lake Superior, Canada and the Ushkanii Islands, Lake Baikal, Russia. MA thesis research, Western Michigan University, 1995-1998.
- The Manitou Collaborative. No Date. The Manitou Collaborative. Home. Available at <https://sites.google.com/site/manitoucollab/> Accessed 23 April 2012.
- Mattes, B. No date. Great Lakes Indian Fish and Wildlife Commission (GLIFWC) Ceded territories mining update. Available at http://www.glifwc.org/Accordian_Stories/stampsands.html Accessed 7 March 2013.
- McCullough, D.G., R.L. Heyd, and J.G. O'Brien. 2002. Michigan State University Extension Bulletin E-2746: Biology and Management of Beech Bark Disease. Michigan State University Extension, East Lansing, MI. Reprinted March 2005. Available at <http://www.baycounty-mi.gov/Docs/Health/GypsyMoth/BeechBark.pdf> Accessed 5 March 2013.
- Michigan DEQ (Michigan Department of Environmental Quality). 2013a. National Pollutant Discharge Elimination System (NPDES) Permits, Applicable Rules & Regulations, Part 4 Rules. Available at <http://www.mi.gov/deqnpdes> Accessed 30 March 2013.
- Michigan DEQ (Michigan Department of Environmental Quality). 2013b. Environmental Areas. Available at http://www.michigan.gov/deq/0,4561,7-135-3313_3677_3700-10863--,00.html Accessed 30 March 2013.
- Michigan DEQ (Michigan Department of Environmental Quality). 2013c. Environmental Areas, listed by County. Available at http://www.michigan.gov/deq/0,4561,7-135-3313_3677_3700-10863--,00.html Accessed 30 March 2013.
- Michigan DEQ (Michigan Department of Environmental Quality). 2013d. Michigan Townships Containing State Designated Environmental Areas. Available at http://www.michigan.gov/deq/0,4561,7-135-3313_3677_3700-10863--,00.html Accessed 30 March 2013.
- Michigan DEQ (Michigan Department of Environmental Quality). 2013z. Areas of Concern. Available at www.michigan.gov/deqaocprogram Accessed 21 June 2013.
- Michigan DNR (Michigan Department of Natural Resources). 2013a. Overview Michigan's Natural Rivers Program. Available at: http://www.michigan.gov/dnr/0,4570,7-153-10364_52259_31442--,00.html Accessed 13 February 2013.
- Michigan DNR (Michigan Department of Natural Resources). 2012. Press Releases: DNR reports U.P.'s Duck lake Fire is 100 percent contained. Available at http://www.michigan.gov/dnr/0,4570,7-153-10371_10402-280414--,00.html Accessed 23 May 2013.
- Michigan DNR (Michigan Department of Natural Resources). 1987. Remedial Action Plan for Torch Lake Area of Concern. Available at http://www.epa.gov/greatlakes/aoc/torchlake/pdfs/RAP_Torch-Lake_1987.pdf Accessed 5 June 2013.
- Minnesota DNR (Minnesota Department of Natural Resources). 2013a. Cloquet River. Available at: <http://www.dnr.state.mn.us/watertrails/cloquetriver/more.html> Accessed 11 January 2013.
- Minnesota DNR (Minnesota Department of Natural Resources). 2013b. North Shore Peregrine Falcon Eyries IBA: Minnesota DNR. Available at http://www.dnr.state.mn.us/north_shore_iba.html Accessed 4 March 2013.
- Minnesota DNR (Minnesota Department of Natural Resources). 2013c. Hawk Ridge Nature Reserve IBA: Minnesota DNR. Available at http://www.dnr.state.mn.us/hawk_ridge_iba.html Accessed 4 March 2013.
- Minnesota DNR (Minnesota Department of Natural Resources). 2013d. Sax-Zim Bog IBA: Minnesota DNR. Available at http://www.dnr.state.mn.us/sax_zim_bog_iba.html Accessed 4 March 2013.
- Minnesota DNR (Minnesota Department of Natural Resources). 2013e. Minnesota Biological Survey. Available at <http://www.dnr.state.mn.us/mbs/index.html> Accessed 23 April 2013.
- Minnesota DNR (Minnesota Department of Natural Resources). 2012a. A State Water Trail Guide to the Upper St. Louis River [Map]. Available at <http://www.dnr.state.mn.us/watertrails/stlouisriver/index.html> Accessed 11 January 2013.
- Minnesota DNR (Minnesota Department of Natural Resources). 2012b. A State Water Trail Guide to the Lower St. Louis River [Map]. Available at <http://www.dnr.state.mn.us/watertrails/stlouisriver/index.html> Accessed 11 January 2013.
- Minnesota DNR (Minnesota Department of Natural Resources). 2008. An Evaluation of the Ecological Significance of the Art Lake Hardwood Ridges Lake County, Minnesota. Minnesota Department of Natural Resources, Minnesota County Biological Survey.
- Minnesota DNR (Minnesota Department of Natural Resources). 1984. Natural Heritage Program Project Evaluation: Susie Island. Minnesota Department of Natural Resources, Minnesota Natural Heritage Program.

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

- Minnesota PCA (Minnesota Pollution Control Agency). 2012a. Lake Superior - North Watershed. Available at <http://www.pca.state.mn.us/index.php/water/water-types-and-programs/watersheds/lake-superior-north.html> Accessed 28 January 2013.
- Minnesota PCA (Minnesota Pollution Control Agency). 2012b. St. Louis River Watershed. Available at http://www.pca.state.mn.us/index.php?option=com_k2&layout=item&view=item&Itemid=2893&id=2786 Accessed 13 January 2013.
- Minnesota PCA (Minnesota Pollution Control Agency). 2012c. Contaminated Sediment Studies: St. Louis River Area of Concern. Available at http://www.pca.state.mn.us/index.php?option=com_k2&Itemid=104&id=741&layout=item&view=item Accessed 13 January 2013.
- Minnesota PCA (Minnesota Pollution Control Agency). 2012d. Lake Superior – South Watershed. Available at http://www.pca.state.mn.us/index.php?option=com_k2&layout=item&view=item&Itemid=2893&id=2792 Accessed 4 February 2013.
- Minnesota PCA (Minnesota Pollution Control Agency). 2012e. Water quality standards. Available at <http://www.pca.state.mn.us/index.php/water/water-permits-and-rules/water-rulemaking/water-quality-standards.html#elements-of-standards> Modified 11 December 2012. Accessed 22 May 2013.
- Minnesota PCA (Minnesota Pollution Control Agency). 2011a. Cloquet River Watershed. Available at <http://www.pca.state.mn.us/index.php/water/water-types-and-programs/watersheds/cloquet-river.html> Modified 28 November 2011. Accessed 13 January 2013.
- National Audubon Society. 2013. Important Bird Areas Database, Boundary Digital Data Set. Ivyland, Pennsylvania.
- National Audubon Society. 2012. Important Bird Areas in the U.S. Available at <http://www.audubon.org/bird/iba>. Accessed 4 March 2012
- NERRS (National Estuarine Research Reserve System). No date. NERRS Reserves, Lake Superior, WI. Available at <http://www.nerrs.noaa.gov/Reserve.aspx?ResID=LKS> Accessed 14 January 2013.
- NERRS (National Estuarine Research Reserve System). 2010. Lake Superior National Estuarine Research Reserve. Available at <http://www.nerrs.noaa.gov/NERRSReserve.aspx?ID=616&ResID=LKS> Accessed 14 January 2013.
- National Park Service (NPS). U.S. Department of the Interior. 2013a. Apostle Islands National Lakeshore Wisconsin. Viral Hemorrhagic Septicemia. Available at <http://www.nps.gov/apis/parkmgmt/vhs.htm> Modified 12 February 2013. Accessed 12 February 2013.
- National Park Service (NPS). U.S. Department of the Interior. 2013b. Apostle Islands National Lakeshore Wisconsin. Available at <http://www.nps.gov/apis/index.htm> Modified 12 February 2013. Accessed 12 February 2013.
- National Park Service (NPS). U.S. Department of the Interior. 2013c. Isle Royale Michigan. Available at <http://www.nps.gov/isro/index.htm> Modified 5 February 2013. Accessed 19 February 2013.
- Newman, L.E., R.B. DuBois, and T. N. Halpern (EDS.). 2003. A brook trout rehabilitation plan for Lake Superior. Great Lakes Fish. Comm. Misc. Publ. 2003-03. Available at <http://www.glfc.org/lakecom/lsc/lsc/home.php> Accessed 29 January 2013.
- Nipigon Bay Remedial Action Plan Team. 2011. Nipigon Bay Area of Concern Update 2011. Available at <http://www.northshorerap.ca/publications> Accessed 28 June 2013.
- Noble, T.W. 1984. Life Science Inventory: Life Science Inventory Check-sheet, Michipicoten Island. OMNR Northeast Region.
- North Shore Forest Collaborative. No date. Home. Available at <https://sites.google.com/site/northshoreforestcollaborative/home> Accessed 23 April 2013.
- OMNR Ontario Ministry of Natural Resources. No date a. Hydropower Peaking and River Ecology: A Tale of Two Rivers. Available at http://www.mnr.gov.on.ca/stdprodconsume/groups/lr/@mnr/@aquatics/documents/document/stdprod_102727.pdf Accessed 19 February 2013.
- OMNR Ontario Ministry of Natural Resources. 2013a. Species at Risk in Ontario (SARO) List. Available at <http://www.mnr.gov.on.ca/en/Business/Species/2ColumnSubPage/276722.html> Updated 24 January 2013. Modified 28 January 2013. Accessed 21 March 2013.
- OMNR Ontario Ministry of Natural Resources. 2013b. Species at Risk in Ontario (SARO) List. Available at <http://www.mnr.gov.on.ca/en/Business/Species/2ColumnSubPage/276722.html> Updated 22 April 2013. Accessed 19 June 2013.
- OMNR Ontario Ministry of Natural Resources. 2013c. Wanted: Information about the Lake Sturgeon. Available at http://www.mnr.gov.on.ca/en/Business/GreatLakes/1ColumnSubPage/STDPROD_083640.html Modified 18 July 2011. Accessed 21 March 2013.
- OMNR Ontario Ministry of Natural Resources. 2013d. Fish Ontario Fisheries Management Zone 9 (Lake Superior). Available at http://www.mnr.gov.on.ca/en/Business/LetsFish/2ColumnSubPage/STDPROD_101354.html Modified 9 January 2013. Accessed 15 February 2013.
- OMNR Ontario Ministry of Natural Resources. Northwest Region Planning Unit. 2012. Black Bay & Black Sturgeon River Native Fisheries Rehabilitation – Decommissioning of the Camp 43 dam and construction of a multi-purpose sea lamprey barrier at Eskwanonwatin Lake. Project Description – Summary. Available at

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

- http://www.mnr.gov.on.ca/en/Business/LetsFish/2ColumnSubPage/STDPROD_101354.html Accessed 20 February 2013.
- OMNR Ontario Ministry of Natural Resources. 2007a. Crown Land Use Policy Atlas Policy Report E39g-7: Great Lakes Coast – Sault Ste. Marie. Available at <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=E39g-7> Updated 1 June 2007. Accessed 18 June 2013.
- OMNR Ontario Ministry of Natural Resources. 2006a. Crown Land Use Policy Atlas Policy Report P282: Batchawana River. Available at <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=P282> Updated 31 January 2006. Accessed 19 February 2013.
- OMNR Ontario Ministry of Natural Resources. 2006b. Crown Land Use Policy Atlas Policy Report P253: Goulais River. Available at <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=P253> Updated 31 January 2006. Accessed 19 February 2013.
- OMNR Ontario Ministry of Natural Resources. 2006c. Crown Land Use Policy Atlas Policy Report P292: Lake Superior Additions. Available at <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=P292> Updated 31 January 2006. Accessed 19 February 2013.
- OMNR Ontario Ministry of Natural Resources. 2006d. Crown Land Use Policy Atlas Policy Report P292e: Lake Superior. <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=P292e> Updated 31 January 2006. Accessed 20 February 2013.
- OMNR Ontario Ministry of Natural Resources. 2006e. Crown Land Use Policy Atlas Policy Report P1877: Montreal River. <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=P1877> Updated 31 January 2006. Accessed 20 February 2013.
- OMNR Ontario Ministry of Natural Resources. 2006f. Crown Land Use Policy Atlas Policy Report C1517: South Michipicoten River – Superior Shoreline. <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=C1517> Updated 31 January 2006. Accessed 20 February 2013.
- OMNR Ontario Ministry of Natural Resources. 2006g. Crown Land Use Policy Atlas Policy Report P1768: Michipicoten. <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=P1768> Updated 31 January 2006. Accessed 20 February 2013.
- OMNR Ontario Ministry of Natural Resources. 2006h. Crown Land Use Policy Atlas Policy Report C1519: Lake Superior Highlands. <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=C1519> Updated 31 January 2006. Accessed 20 February 2013.
- OMNR Ontario Ministry of Natural Resources. 2006i. Crown Land Use Policy Atlas Policy Report P1511: Nimoosh (Waterway Class). <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=P1511> Updated 31 January 2006. Accessed 20 February 2013.
- OMNR Ontario Ministry of Natural Resources. 2006j. Crown Land Use Policy Atlas Policy Report P1513: Pukaskwa River. <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=P1513> Updated 31 January 2006. Accessed 20 February 2013.
- OMNR Ontario Ministry of Natural Resources. 2006k. Crown Land Use Policy Atlas Policy Report P1514: Pokei Lake/ White River Wetlands. <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=P1514> Updated 31 January 2006. Accessed 20 February 2013.
- OMNR Ontario Ministry of Natural Resources. 2006l. Crown Land Use Policy Atlas Policy Report P1501e: Craig's Pit. <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=P1501e> Updated 31 January 2006. Accessed 20 February 2013.
- OMNR Ontario Ministry of Natural Resources. 2005a. Crown Land Use Policy Atlas Policy Report C307: Ile Parisienne. <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=C307> Updated 14 November 2005. Accessed 18 June 2013.
- OMNR Ontario Ministry of Natural Resources. 2005b. Crown Land Use Policy Atlas Policy Report C291: Goulais River Beach Ridges. <http://www.lio.ontario.ca/imf-ows/sites/clupa/xmlReader.jsp?xsl=XML/web-primary.xsl&polid=C291> Updated 14 November 2005. Accessed 1 April 2013.
- OMNR Ontario Ministry of Natural Resources / Ontario Parks. 2004a. Michipicoten Post and Michipicoten Island Background Information. Queen's Printer of Ontario. Available at http://www.ontarioparks.com/english/planning_pdf/michipicoten_background_info2.pdf Accessed 15 February 2013.
- Ontario Peregrine Falcon Recovery Team. 2010. Recovery strategy for the Peregrine Falcon (*Falco peregrines*) in Ontario. Ontario Recovery Strategy Series. Prepared for the Ontario Ministry of Natural Resources, Peterborough, Ontario. vi + 36pp. Available at <http://www.mnr.gov.on.ca/stdprodconsume/groups/lr/@mnr/@species/documents/document/286970.pdf> Accessed 20 March 2012.
- Ontario Woodland Caribou Recovery Team. 2008. Woodland Caribou (*Rangifer tarandus caribou*) (Forest-dwelling, Boreal Population) in Ontario. Prepared for the Ontario Ministry of Natural Resources, Peterborough, Ontario. 93 pp.

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

- Available at <http://www.mnr.gov.on.ca/stdprodconsume/groups/lr/@mnr/@species/documents/document/251755.pdf> Accessed 29 November 2012.
- Orvana Minerals Corp. 2013. Projects: Copperwood. Available at <http://www.orvana.com/projects/copperwood/index.html> Accessed 16 April 2013.
- Parks Canada. 2013. Pukaskwa National Park of Canada. Available at <http://www.pc.gc.ca/eng/pn-np/on/pukaskwa/index.aspx> Modified 18 February 2013. Accessed 20 February 2013.
- Parks Canada. 2012. Pukaskwa National Park of Canada Ecosystem Conservation Plan Executive Summary. Available at <http://www.pc.gc.ca/eng/pn-np/on/pukaskwa/natcul/natcul6.aspx> Modified 11 April 2012. Accessed 20 February 2013.
- Parks Canada. 2009. Lake Superior National Marine Conservation Area of Canada. Map. Available at <http://www.pc.gc.ca/eng/amnc-nmca/on/super/visit/visit3a.aspx?m=1> Accessed 20 February 2013.
- Peninsula Harbour Remedial Action Plan Team. 1991. Stage 1: Environmental Conditions and Problem Definition. Available at <http://www.northshorerap.ca/publications> Accessed 28 June 2013.
- Petzold, M. 2003. Personal Communication. Fisheries Biologist, Sault Ste. Marie, OMNR.
- Ramsar and Wetlands International. 2013. Ramsar Sites Information Service Summary Description: United States of America 4US031. Available at <http://ramsar.wetlands.org/Database/SearchforRamsarsites/tabid/765/Default.aspx> Accessed 22 May 2013.
- Redig, P.T., J.S. Castrale and A. Burnette. 2010. Midwest Peregrine Falcon Restoration, 2010 Report. Univ. of Minnesota, St. Paul. 75 pp. <http://midwestperegrine.org/>
- Rup, M.P., S.A. Bailey, C.J. Wiley, M.S. Minton, A.W. Miller, G.M. Ruiz, and H.J. MacIsaac. 2010. Domestic ballast operations on the Great Lakes: potential importance of Lakers as a vector for introduction and spread of nonindigenous species. *Can. J. Fish. Aquat. Sci.* 67: 256-268. Available at <http://www.caisn.ca/publications/Importance.pdf> Accessed 21 February 2013.
- SLRT (St. Louis River Trail Association). 2012. St. Louis River Trail. Available at <http://www.slrt.org/> Accessed 13 January 2013.
- Sugarloaf: The North Shore Stewardship Association. No date. Home. Available at <http://www.sugarloafnorthshore.org/> Accessed 23 April 2013.
- TNC (The Nature Conservancy). No date a. Clough Island Questions & Answers. Available at <http://www.nature.org/ourinitiatives/regions/northamerica/unitedstates/wisconsin/howwework/clough-island.xml> Accessed 14 January 2013.
- TNC (The Nature Conservancy). No date b. Iron Mining In the Penokee Range – What’s at Risk [Report]. Available at <http://www.nature.org/ourinitiatives/regions/northamerica/unitedstates/wisconsin/mining-in-the-penokee-gogebic-range-whats-at-risk.xml> Accessed 22 May 2013.
- TNC (The Nature Conservancy). 2010. News Release: The Nature Conservancy Preserves Island Neat Lake Superior. Available at <http://www.nature.org/ourinitiatives/regions/northamerica/unitedstates/wisconsin/howwework/clough-island.xml> Accessed 14 January 2013.
- Thompson, J.E. 1994. Life Science and Human History Features of Michipicoten Post Provincial Park. OMNR.
- Thunder Bay Field Naturalists. No date. Project Peregrine. Available at www.tbfn.net/project-peregrine Accessed 20 March 2013.
- Thunder Bay RAP. 2013. DRAFT Thunder Bay Remedial Action Plan Update 2009-2012. 38pp.
- UNESCO United Nations Educational, Scientific and Cultural Organization, MAB Biosphere Reserves Directory. 2005. Isle Royale. Available at <http://www.unesco.org/mabdb/br/brdir/directory/biores.asp?code=USA+33&mode=all> Updated 11 March 2005. Accessed 19 February 2013.
- USDA Forest Service (United States Department of Agriculture Forest Service). No date a. Superior National Forest, The Manitou Collaborative. Available at <http://www.fs.usda.gov/detail/superior/landmanagement/resourcemanagement/?cid=stelprdb5209873> Accessed 23 April 2013.
- USDA Forest Service (United States Department of Agriculture Forest Service). 2007a. Ottawa National Forest, Wild and Scenic River Comprehensive River Management Plan. Available at www.rivers.gov/rivers/documents/plans/ottawa-nf-plan.pdf Accessed 12 May 2013.
- USDA Forest Service (United States Department of Agriculture Forest Service). 2007b. McCormick Wilderness Ottawa National Forest [Map]. Available at <http://www.fs.usda.gov/main/ottawa/maps-pubs> Accessed 12 May 2013.
- USDA NRCS (United States Department of Agriculture Natural Resources Conservation Service). No date a. Rapid Watershed Assessment Baptism-Brule (MN) HUC: 04010101. Available at <http://www.mn.nrcs.usda.gov/technical/rwa/Assessments/04010101.html> Accessed 27 January 2013.
- USDA NRCS (United States Department of Agriculture Natural Resources Conservation Service). No date b. Rapid Watershed Assessment Beaver-Lester (MN) HUC: 04010102. Available at <http://www.mn.nrcs.usda.gov/technical/rwa/Assessments/04010102.html> Accessed 4 February 2013.

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

- USDA NRCS (United States Department of Agriculture Natural Resources Conservation Service). No date c. Rapid Watershed Assessment Resource Profile Beartrap – Nemadji (MN/WI) HUC: 04010301. Available at <http://www.mn.nrcs.usda.gov/technical/rwa/Assessments/04010301.html> Accessed 10 February 2013.
- USDA NRCS (United States Department of Agriculture Natural Resources Conservation Service). No date d. Rapid Watershed Assessment Bad-Montreal River (WI) HUC: 04010302. Available at <http://www.wi.nrcs.usda.gov/technical/rwa.html> Accessed 12 February 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013a. Great Lakes Areas of Concern: St. Marys River. Available at <http://www.epa.gov/greatlakes/aoc/stmarys/index.html> Modified 30 January 2013. Accessed 7 June 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013b. Baptism-Brule Watershed –04010101. Available at http://cfpub.epa.gov/surf/huc.cfm?huc_code=04010101 Accessed 19 February 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013c. Great Lakes Areas of Concern: St. Louis River. Available at <http://www.epa.gov/greatlakes/aoc/stlouis.html> Updated 30 January 2013. Accessed 7 June 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013d. Beaver-Lester Watershed –04010102. Available at http://cfpub.epa.gov/surf/huc.cfm?huc_code=04010102 Updated 19 February 2013. Accessed 19 February 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013e. St. Louis River Beneficial Use Impairments. Available at http://www.epa.gov/greatlakes/aoc/stlouis/StLouis-AOC_BUIs.pdf Accessed 12 January 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013f. Region 5 Superfund: Arrowhead Refinery Co. Available at <http://www.epa.gov/region5/superfund/npl/minnesota/MND980823975.html> Modified 13 June 2013. Accessed 27 June 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013g. Region 5 Superfund: St. Louis River Site. Available at <http://www.epa.gov/region5/superfund/npl/minnesota/MND039045430.html> Modified 13 June 2013. Accessed 27 June 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013h. St. Louis Watershed –04010201. Available at http://cfpub.epa.gov/surf/huc.cfm?huc_code=04010201 Updated 19 February 2013. Accessed 19 February 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013i. Cloquet Watershed –04010202. Available at http://cfpub.epa.gov/surf/huc.cfm?huc_code=04010202 Updated 19 February 2013. Accessed 19 February 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013j. Region 5 Cleanup Sites: Ashland/Northern States Power Lakefront Site. Available at <http://www.epa.gov/region5/cleanup/ashland/> Modified 9 May 2013. Accessed 23 May 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013k. Bad-Montreal Watershed –04010302. Available at http://cfpub.epa.gov/surf/huc.cfm?huc_code=04010302 Accessed 12 February 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013l. Black-Presque Isle Watershed –04020101. Available at http://cfpub.epa.gov/surf/huc.cfm?huc_code=04020101 Accessed 12 February 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013m. Ontonagon Watershed –04020102. Available at http://cfpub.epa.gov/surf/huc.cfm?huc_code=04020102 Accessed 12 February 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013n. Great Lakes Areas of Concern: Torch Lake. Available at <http://www.epa.gov/greatlakes/aoc/torchlake/index.html> Modified 6 June 2013. Accessed 6 June 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013o. Region 5 Cleanup Sites: Torch Lake Superfund Site. Available at <http://www.epa.gov/region5/cleanup/torchlake/> Modified 15 April 2013. Accessed 10 June 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013q. Region 5 Superfund: Cannerton Industries, Inc. Available at <http://www.epa.gov/region5/superfund/npl/michigan/MID980678627.html> Modified 13 June 2013. Accessed 27 June 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013r. Beartrap Nemadji Watershed –04010301. Available at http://cfpub.epa.gov/surf/huc.cfm?huc_code=04010301 Accessed 11 February 2013.
- U.S. EPA (United States Environmental Protection Agency). 2013s. Great Lakes Areas of Concern: Deer Lake. Available at <http://www.epa.gov/greatlakes/aoc/deerlake/index.html> Modified 3 June 2013. Accessed 7 June 2013.
- U.S. EPA (United States Environmental Protection Agency). 2012a. National Priorities List (NPL) NPL Site Narrative for Arrowhead Refinery Co. Available at <http://www.epa.gov/superfund/sites/npl/nar642.htm> Modified 27 November 2012. Accessed 27 June 2013. U.S. EPA (United States Environmental Protection Agency). 2011. Region 5 Cleanup Sites: DuPont Barksdale Explosives Plant Site. Available at <http://www.epa.gov/region05/cleanup/duPont/index.htm> Modified 29 June 2011. Accessed 23 May 2013.
- USGS. 2012a. Nonindigenous Aquatic Species (NAS Data) for the Baptism Brule – HUC 04010101 Available at <http://water.usgs.gov/lookup/getwatershed?04010101> Modified 10 December 2012. Accessed 09 January 2013.
- USGS. 2012b. Nonindigenous Aquatic Species (NAS Data) for the Beaver Lester – HUC 04010102 Available at <http://nas.er.usgs.gov/queries/SpeciesList.aspx?Group=&Status=0&FMB=0&pathway=0&Sortby=1&Size=50&HUCNumber=4010102> Modified 10 December 2012. Accessed 21 February 2013.
- University of Wisconsin-Extension. 2011. Lake Superior Basin in Wisconsin Programs Slow the Flow. Available at <http://basineducation.uwex.edu/lakesuperior/programs.html#flow> Accessed 23 May 2012.

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

- University of Wisconsin-Extension. 2010. Lake Superior National Estuarine Research Reserve Management Plan 2010-2015. Available at http://www.nerrs.noaa.gov/Doc/PDF/Reserve/LKS_MgmtPlan.pdf Accessed 5 February 2013.
- Vucetich, J.A. 2012. Wolves & Moose of Isle Royale. About the Project: Overview. Available at http://isleroyalewolf.org/overview/overview/at_a_glance.html Accessed 19 February 2013.
- Vucetich, J.A. and R.O. Peterson. 2013. Ecological Studies of Wolves on Isle Royale. Annual Report 2012-13. School of Forest Resources and Environmental Science. Michigan Technological University. Available at <http://www.mtu.edu/news/stories/2013/march/story87157.html> Accessed 28 May 2013.
- White, D.J. 2000. Life Science Study of Michipicoten Island Provincial Park. Ontario Ministry of Natural Resources, Northeastern Region, Sudbury
- Whitefish Point Bird Observatory. No date. Whitefish Point Bird Observatory. Available at <http://www.wpbo.org/> Accessed 5 March 2013.
- Wikipedia contributors. "Huron Mountain Club". Wikipedia, The Free Encyclopedia, Available at http://en.wikipedia.org/w/index.php?title=Huron_Mountain_Club&oldid=529585343 Accessed 12 May 2013.
- WDNR Wisconsin Department of Natural Resources. No date. Surface Water Data Viewer – Designated Waters. Available at <http://dnrmaps.wi.gov/imf/imf.jsp?site=SurfaceWaterViewer.deswaters> Accessed 22 May 2013.
- WDNR Wisconsin Department of Natural Resources. 2013a. Nemadji River Floodplain Forest (No. 320). Available at <http://dnr.wi.gov/topic/Lands/naturalareas/index.asp?SNA=320> Modified 4 January 2013. Accessed 11 February 2013.
- WDNR Wisconsin Department of Natural Resources. 2013b. [DRAFT] White River Property Group (Bayfield & Ashland Counties) Draft Master Plan and Environmental Assessment.
- WDNR Wisconsin Department of Natural Resources. 2012a. Outstanding and Exceptional Resource Waters. Available at <http://dnr.wi.gov/topic/SurfaceWater/orwerw.html> Modified 24 April 2012. Accessed 12 February 2013.
- WDNR Wisconsin Department of Natural Resources. 2012b. DNR Fisheries Areas St. Louis/Red River Stream Bank Protection Area. Available at <http://dnr.wi.gov/topic/Lands/fisheriesareas/2460stlouisriver.html> Modified 29 August 2012. Accessed 17 March 2013.
- WDNR Wisconsin Department of Natural Resources. 2012c. Primary Sites – Western Lake Superior. Available at <http://dnr.wi.gov/topic/wetlands/cw/WLSup/> Modified 20 June 2012. Accessed 21 May 2013.
- WDNR Wisconsin Department of Natural Resources. 2012d. Primary Sites – Eastern Lake Superior. Available at <http://dnr.wi.gov/topic/wetlands/cw/ELSup/> Modified 20 June 2012. Accessed 21 May 2013.
- WDNR Wisconsin Department of Natural Resources. 2008a. Wisconsin's Priority Conservation Opportunity Areas for Wildlife Species of Greatest Conservation Need 2008-2015 [Map]. Available at <http://dnr.wi.gov/topic/WildlifeHabitat/COA.html> Modified 20 June 2012. Accessed 22 May 2013.
- WDNR Wisconsin Department of Natural Resources. 2008b. Wisconsin's Priority Conservation Opportunity Areas for Wildlife Species of Greatest Conservation Need 2008-2015 Superior Coastal Plain Ecological Landscape [Map]. Available at <http://dnr.wi.gov/topic/WildlifeHabitat/COA.html> Modified 20 June 2012. Accessed 22 May 2013.
- WDNR Wisconsin Department of Natural Resources. 2008c. Wisconsin's Wildlife Action Plan (2005-2015) Priority Conservation Actions & Conservation Opportunity Areas Superior Coastal Plain Ecological Landscape [Report]. Available at <http://dnr.wi.gov/topic/WildlifeHabitat/COA.html> Modified 24 June 2012. Accessed 22 May 2013.
- WDNR Wisconsin Department of Natural Resources. 2005. Wisconsin's Strategy for Wildlife Species of Greatest Conservation Need. Madison, WI. Available at <http://dnr.wi.gov/topic/WildlifeHabitat/documents/WWAP.pdf>. Accessed 22 May 2013.
- WDNR Wisconsin Department of Natural Resources. 1999a. Watershed – Montreal River (LS15). Available at <http://dnr.wi.gov/water/watershedDetail.aspx?code=LS15&Name=Montreal%20River> Accessed 12 February 2013.
- WDNR Wisconsin Department of Natural Resources. 1999b. Watershed – Lower Bad River (LS09). Available at <http://dnr.wi.gov/water/watershedDetail.aspx?key=924673> Accessed 22 May 2013.
- Wisconsin Wetlands Association. No date a. Overview of the Wetlands Gems™ Program: Wisconsin's Wetland Gems. Available at <http://www.wisconsinwetlands.org/Gems/WetlandGemsIntroFolder.pdf> Accessed 21 May 2013.
- Wisconsin Wetlands Association. No date b. Wisconsin's Wetland Gems Regional Wetland Gems Lists Superior Region. Available at <http://www.wisconsinwetlands.org/gemsSUregion.htm> Accessed 21 May 2013.

Personal Communications

- Blust, W. (Wisconsin Department of Natural Resources). March 6 2013. Personal communication.
- Carlson, B. (Minnesota Department of Natural Resources). March 20 2013. Personal communication.
- Chase, M. (Ontario Ministry of Natural Resources). June 3 2013. Personal communication.
- Czypinski, G. (United States Department of the Interior, Fish and Wildlife Service). March 20 2013. Personal communication.
- Hagen C. et al. Includes input from Frank Koshore, Cherie Hagen, Lynelle Hanson, Steve LaValley, Ryan Magana, and Paul Piszczek. (Wisconsin Department of Natural Resources). March 20 2013. Personal communication.

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

Hester, C. and Tillison, N. (Bad River Band of the Lake Superior Tribe of Chippewa Indians). March 25 2013. Personal communication.

Jennings, M. (Wisconsin Department of Natural Resources). March 13 2013. Personal communication.

Kaspar, T. (1854 Treaty Authority). March 14 2013. Personal communication

O'Connor, R. (Wisconsin Department of Natural Resources). March 15 2013. Personal communication.

Gerdes, L. (Minnesota Department of Natural Resources). March 18 2013. Personal communication.

Greenwood, S. (Ontario Ministry of Natural Resources). February 3 2013. Personal communication.

Greenwood, S. (Ontario Ministry of Natural Resources). March 11 2013. Personal communication.

Greenwood, S. (Ontario Ministry of Natural Resources). March 12 2013. Personal communication.

Greenwood, S. (Ontario Ministry of Natural Resources). May 27 2013. Personal communication.

Lebel, S. and G. Eason. (Ontario Ministry of Natural Resources). June 18 2013. Personal communication.

McCammon Soltis, A. (Great Lakes Indian Fish and Wildlife Commission). March 19 2013. Personal communication.

Moore, S. (Grand Portage Band of Chippewa). May 14 2013. Personal communication.

O'Connor, R. (Wisconsin Department of Natural Resources). March 15, 2012. Personal communication.

Perry, E. (Minnesota Department of Natural Resources). February 26 2013. Personal communication.

Preisser, M. (Michigan Department of Environmental Quality). May 31 2013. Personal communication.

Schuldt, N. (Fond du Lac Environmental Program). March 19 2013. Personal communication.

Swart, S. (Michigan Department of Environmental Quality). March 6 2013. Personal communication.

Taft, W. (Michigan Department of Environmental Quality). February 25, 2013. Personal communication.

Tillison, N. (Bad River Band of Lake Superior Tribe of Chippewa Indians). April 26 2013. Personal communication.

Toshner, S. (Wisconsin Department of Natural Resources). March 6 2013. Personal communication.

Vis, C. (Parks Canada). May 21 2013. Personal communication.

Appendix A

Spatial Data Catalogue and Methods

Dataset	Vintage	Scale/ Resolution	Source	Notes
COASTAL AND WATERSHED FEATURES				
Atlas of the Spawning and Nursery Areas of Great Lakes Fishes	1982	-	Goodyear, C. S., T. A. Edsall, D. M. Ormsby Dempsey, G. D. Moss, and P. E. Polanski	Digitized the maps in report for Lake Trout and Lake Whitefish http://www.glsc.usgs.gov/main.php?content=products_publications_atlas&title=Publications0&menu=products
Important Habitat Sites and Areas	2006		Lake Superior Binational Program's Habitat Committee	Available at
SOLEC Shoreline Data (CAN)	2008	-	Environment Canada	Used for Ontario Also used for Artificial Shoreline (Threat).
Great Lakes and St. Lawrence River Medium Resolution Vector Shoreline Data	1997		Great Lakes Environmental Research Laboratory, NOAA	Used for the United States. Also used for Artificial Shoreline (Threat). – below just artificial??
Continually Updated Shoreline Product (CUSP)	2011	5,000 – 20,000	U.S. Department of Commerce (DOC), National Oceanic and Atmospheric Administration (NOAA), National Ocean Service (NOS), National Geodetic Survey (NGS)	Data Download: http://www.ngs.noaa.gov/CUSP/ Portal: http://shoreline.noaa.gov/data/datasheets/
Wetland Units (Ontario)	2012	1:10,000-1:50,000	Ontario Ministry of Natural Resources	Version also identifies whether wetland evaluated and provincial/other significance
Great Lakes Coastal Wetland Inventory	2004	1:10,000	Publisher: Great Lakes Wetland Consortium Originator: Environment Canada US Geological Service Michigan Natural Features Inventory Ontario Ministry of Natural Resources	Albert, D.A., J. Ingram, T. Thompson, and D. Wilcox http://www.glc.org/wetlands/inventory.html
National Wetlands Inventory Polygons (US)	2003	1:24,000	DNR Minnesota	http://deli.dnr.state.mn.us/metadata.html?id=L260000162101
DNR 100k Wetlands	2002	1:100,000	DNR Minnesota	http://deli.dnr.state.mn.us
Final Wetland Inventory (US)	2007	1:24,000	DNR Michigan/Michigan Department of Technology, Management and Budget	http://www.mcgi.state.mi.us/mgdl/?rel=thext&action=thmname&cid=3&cat=Final%5FWetland%5FInventory Available by County, assembled together
Wetlands	1994	1:24,000	DNR Wisconsin	Available by County, assembled together
Land Use / Land Cover	1999	200 m	Natural Resources Research Institute, University of Minnesota Duluth	Covers entire basin. Used for analysis. http://www.nrri.umn.edu/lsgis2/data/landuse.html Subset from classified Landsat MSS of Great Lakes Basin by Peter Wolter. US land cover was derived from the National Land Cover Database (Vogelmann <i>et al.</i> 1998)

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

Dataset	Vintage	Scale/ Resolution	Source	Notes
				and Canadian land cover from the Ontario Land Cover Database (Spectralanalysis 2004). Both of these land cover datasets were derived from 30 m Landsat Thematic Mapper satellite data, and use similar land classification schemes
Roads	2011	1:10,000	Ontario Ministry of Natural Resources	Ontario Road Network
	2001	1:24,000	MN Department of Transportation	DOT Basemap Roads - All Types
	2012	1:24,000	Michigan Center for Geographic Information	MI Geographic Framework All Roads (v12b)
	2010	1:24,000	DNR Wisconsin	WI Roads 2010 (TIGER Lines)
CONDITION				
Areas of Concern	2008	1:5,000- 1:10,000	Environment Canada	
Cumulative Stress, Laurentian Great Lakes, 2000-2009	2013	921 m	Dr. J.D. Allan, School of Natural Resources and Environment, University of Michigan	Great Lakes Environmental Assessment and Mapping (GLEAM) Project http://www.greatlakemapping.org
Watershed Stress Index	2011	-	Host et al., Natural Resources Research Institute, University of Minnesota Duluth	Great Lakes Environmental Indicators (GLEI) Project
Lake Superior Bathymetry	1998	1000 m	Natural Resource Research Institute	http://www.nrri.umn.edu/lsgis2/ - Generated 5-m contours from model for mapping bathymetry on maps
IMPORTANT HABITAT SITES AND AREAS				
Important Habitat Sites and Areas	2006		Lake Superior Binational Program's Habitat Committee	Available http://www.nrri.umn.edu/lsgis/importanthabitat.htm
Federal Marine Conservation Areas (Ontario)	2010	50,000	Parks Canada	
National Parks (Ontario)	2003	1:50,000	Ontario Ministry of Natural Resources	
National Parks (US)	2001	1:24,000	Michigan GAP Land Stewardship Coverage	Isle Royal Island geometry: (http://www.mcgi.state.mi.us/mgdl/?rel=thext&action=thmname&cid=4&cat=GAP+Land+Stewardship)
	2008	?	National Park Service	Policy Area (Buffer around island)
National Wildlife Areas	2002	~1:50,000	Canadian Wildlife Service	
Crown Game Preserves	-	-	Ontario Ministry of Natural Resources	
Provincial Parks	2011	1:10,000	Ontario Ministry of Natural Resources	
Recommended Provincial Parks	2007	1:10,000	Ontario Parks	
Conservation Reserves	2007	1:10,000	Ontario Parks	
Conservation Authority Properties	2006 – 2012	1:10,000	Ontario Ministry of Natural Resources with supplemental information added by NCC	Dataset is not regularly maintained by MNR. Some areas have been manually added by NCC
NCC Properties	2012		Nature Conservancy of Canada	
NGO	2006		Ontario Ministry of Natural Resources	Dataset is not regularly maintained by MNR. Some areas have been manually added by NCC
Municipal Park	2006		Ontario Ministry of Natural Resources	Dataset is not regularly maintained by MNR.

Lake Superior Biodiversity Conservation Assessment - Volume 2: Regional Unit Summaries

Dataset	Vintage	Scale/ Resolution	Source	Notes
Conservation and Recreation Lands (Michigan)	2008		Ducks Unlimited and The Nature Conservancy	Michigan
Protected Areas Database of the United States Protected Lands Database (US)	2011	1:24,000-1:100,000	USGS	Version 1.2 - http://www.protectedlands.net/padus/preview.php Public land ownership, management and conservation lands nationally, including voluntarily provided privately protected areas
Protected Areas Database of the United States Protected Lands Database (US)	2010		CBI Edition	Version 1.1 - http://www.protectedlands.net/padus/preview.php Public and private protected areas across US at variety of scales
BASE MAP DATA				
Land Use / Land Cover	1999	200 m	Natural Resources Research Institute, University of Minnesota Duluth	Covers entire basin. Used for analysis. http://www.nrri.umn.edu/lsgis2/data/landuse.html Subset from classified Landsat MSS of Great Lakes Basin by Peter Wolter. US land cover was derived from the National Land Cover Database (Vogelmann <i>et al.</i> 1998) and Canadian land cover from the Ontario Land Cover Database (Spectralanalysis 2004). Both of these land cover datasets were derived from 30 m Landsat Thematic Mapper satellite data, and use similar land classification schemes
Great Lakes				
International Boundaries				
State Boundaries				
Detailed Population Centres				
Basin Names				
Lake Superior Basin HYDRO	2006	1:100,000	The Lake Superior Decision Support Project	http://www.nrri.umn.edu/lsgis/databases.htm
WRIP Water Flow Network (CAN)	2005	1:10,000	Ontario Ministry of Natural Resources	